

1st Putnam 1938

Problem A1


A solid in Euclidean 3-space extends from $z = -h/2$ to $z = +h/2$ and the area of the section $z = k$ is a polynomial in k of degree at most 3. Show that the volume of the solid is $h(B + 4M + T)/6$, where B is the area of the bottom ($z = -h/2$), M is the area of the middle section ($z = 0$), and T is the area of the top ($z = h/2$). Derive the formulae for the volumes of a cone and a sphere.

Solution

Let the polynomial be $az^3 + bz^2 + cz + d$. Then the volume is $\int_{-h/2}^{h/2} (az^3 + bz^2 + cz + d) dz = bh^3/12 + dh$. But $B + T = bh^2/2 + 2d$, $M = d$, so $h(B + 4M + T)/6 = bh^3/12 + dh$, which proves the formula. For a sphere radius R , we have $h = 2R$, $B + T = 0$ and $M = \pi R^2$, so the formula gives $4/3 \pi R^3$, as usual. For a cone height h , base area A , we have $B = A$, $T = 0$, $M = A/4$, so the volume is $hA/3$, as usual.

Problem A2

A solid has a cylindrical middle with a conical cap at each end. The height of each cap equals the length of the middle. For a given surface area, what shape maximizes the volume?



Solution

Let the radius be R and the height H . The area is $2\pi RH + 2\pi R\sqrt{R^2 + H^2}$. The volume is $5/3 \pi R^2 H$.

The area is fixed, so for some fixed k , we have $R (H + \sqrt{R^2 + H^2}) = k$. This gives $H = (k^2 - R^4)/(2kR)$. We must now choose R to maximise $f(R) = R^2 H = R (k^2 - R^4)/2k$. Evidently the allowed range for R is from $R = 0$ up to \sqrt{k} (corresponding to $H = 0$). But $f(0) = 0$ and $f(\sqrt{k}) = 0$, so the maximum is at some interior point of the interval. Differentiating, we find it is at $R_{\max} = (k^2/5)^{1/4}$. In terms of the area A , we have $A = 2\pi k$, so $R_{\max} = (A/(\pi 2\sqrt{5}))^{1/2}$.

Problem A3

A particle moves in the Euclidean plane. At time t (taking all real values) its coordinates are $x = t^3 - t$ and $y = t^4 + t$. Show that its velocity has a maximum at $t = 0$, and that its path has an inflection at $t = 0$.

Solution

The speed squared is $(dx/dt)^2 + (dy/dt)^2 = 16t^6 + 9t^4 + 8t^3 - 6t^2 + 2$. Let this be $f(t)$. We have $f'(t) = 12t(8t^4 + 3t^2 + 2t - 1)$. So $f'(t) = 0$ at $t = 0$. Also for t small (positive or negative), $8t^4 + 3t^2 + 2t - 1$ is close to -1 and hence negative, so $f'(t)$ is positive for t just less than 0 and negative for t just greater than 0 . Hence $f(t)$ has a maximum at $t = 0$. Hence the speed does also.

The gradient $dy/dx = (4t^3 + 1)/(3t^2 - 1)$. Let this be $g(t)$. Then $g'(t) = 6t(2t^3 - 2t - 1)/(3t^2 - 1)^2$. Hence $g'(0) = 0$. Also $g'(t)$ is positive for t just less than 0 and negative for t just greater than 0 , so it is a point of inflection.

Problem A4

A notch is cut in a cylindrical vertical tree trunk. The notch penetrates to the axis of the cylinder and is bounded by two half-planes. Each half-plane is bounded by a horizontal line passing through the axis of the cylinder. The angle between the two half-planes is θ . Prove that the volume of the notch is minimized (for given tree and θ) by taking the bounding planes at equal angles to the horizontal plane.

Solution

We find the volume of the notch above the horizontal plane. Suppose that the upper bounding half-plane is at an angle ϕ to the horizontal. We may take the radius of the tree to be 1 . A vertical section through the notch at a distance x from its widest extent is a right-angled triangle with base $\sqrt{1 - x^2}$ and area $1/2 (1 - x^2) \tan \phi$. Hence the volume is $2/3 \tan \phi$. So the total volume of the notch is $2/3 (\tan \phi + \tan(\theta - \phi))$. So we have to find the angle ϕ which minimises $(\tan \phi + \tan(\theta - \phi))$. Differentiating, or otherwise, we easily find that the minimum is at $\phi/2$.

Problem A5

- (1) Find $\lim_{x \rightarrow \infty} x^2/e^x$.
- (2) Find $\lim_{k \rightarrow 0} 1/k \int_0^k (1 + \sin 2x)^{1/x} dx$.

Solution

(1) Let $f(x) = x^3 e^{-x}$. Then $f'(x) = (3x^2 - x^3) e^{-x} < 0$ for $x > 3$. Hence $f(x) < f(3)$ for $x > 3$, so $x^2 e^{-x} < f(3)/x$ for $x > 3$. Hence $x^2 e^{-x}$ tends to zero.

(2) We use L'Hôpital's rule $\lim f(x)/g(x) = \lim f'(x)/g'(x)$. Applied to the expression given it gives $\lim (1 + \sin 2x)^{1/x}$. Write $(1 + \sin 2x)^{1/x} = \exp(1/x \ln(1 + \sin 2x))$. So apply the rule again to $1/x \ln(1 + \sin 2x)$ to get $2 \cos 2x/(1 + \sin 2x)$ which tends to 2. Hence $(1 + \sin 2x)^{1/x}$ tends to e^2 and so does the original expression.

Problem A6

A swimmer is standing at a corner of a square swimming pool. She swims at a fixed speed and runs at a fixed speed (possibly different). No time is taken entering or leaving the pool. What path should she follow to reach the opposite corner of the pool in the shortest possible time?

Solution

Answer: let k be the running speed divided by the swimming speed. For $k > \sqrt{2}$, the unique solution is to run round the outside. For $k < \sqrt{2}$, the unique solution is to swim direct. For $k = \sqrt{2}$ there is no unique solution. Run along a side to X , swim to Y equidistant from the corner between X and Y , then run from Y . The time taken is independent of X .

We may take the side of the square to be 1, the swimming speed to be 1 and the running speed to be k . Let the square be $ABCD$. Suppose the start is at A and the finish at C . Possible routes are (1) run to X on AB , swim to Y on BC , run to C , (2) run to X on AD , swim to Y on CD , run to C , (3) run to X on AB , swim to Y on CD , run to C . We start by considering case (1). Take BX to be x , BY to be y . Then the time taken is $(2 - x - y)/k + \sqrt{(x^2 + y^2)}$. Note that this includes the extreme cases of running all the way ($x = y = 0$) and swimming all the way ($x = y = 1$).

Now $(x - y)^2 \geq 0$, with equality iff $x = y$, so $(x + y)^2 \leq 2(x^2 + y^2)$ and hence $(x + y) \leq \sqrt{2} \sqrt{(x^2 + y^2)}$, with equality iff $x = y$. So if $k > \sqrt{2}$, then $(x + y) < k \sqrt{(x^2 + y^2)}$ and hence $2/k < (2 - x - y)/k + \sqrt{(x^2 + y^2)}$ unless $x = y = 0$ (when we have equality). Hence for $k > \sqrt{2}$, the unique solution is to run all the way.

If $k < \sqrt{2}$, then $(x + y) \leq \sqrt{2} \sqrt{(x^2 + y^2)}$ implies $\sqrt{2} (\sqrt{2} - \sqrt{(x^2 + y^2)}) \leq 2 - x - y$ and hence $k (\sqrt{2} - \sqrt{(x^2 + y^2)}) < 2 - x - y$ unless $x = y = 1$ (when we have equality). So $\sqrt{2} < (2 - x - y)/k + \sqrt{(x^2 + y^2)}$ unless $x = y = 1$. In other words, for $k < \sqrt{2}$ the unique solution is to swim all the way.

For $k = \sqrt{2}$ we have equality (in both the previous paragraphs) iff $x = y$. So any solution with $x = y$ is optimal in this case.

Problem A7

Do either (1) or (2)

(1) S is a thin spherical shell of constant thickness and density with total mass M and center O . P is a point outside S . Prove that the gravitational attraction of S at P is the same as the gravitational attraction of a point mass M at O .

(2) K is the surface $z = xy$ in Euclidean 3-space. Find all straight lines lying in S . Draw a diagram to illustrate them.

Solution

(1) Let Q be a point on S . The obvious coordinate is the angle $\theta = \text{angle } QOP$. The density is $\rho = M/4\pi r^2$. By symmetry the attraction on P is towards O . Let the distance PO be d and the radius of the sphere be r . Let the gravitational constant be G . The component of the attraction towards O (per unit mass at P) is $G \int_0^\pi r \, d\theta \, 2\pi r \sin \theta \rho (d - r \cos \theta) (d^2 + r^2 - 2dr \cos \theta)^{-3/2}$. Note that the factor $(d - r \cos \theta) (d^2 + r^2 - 2dr \cos \theta)^{-1/2}$ is needed to resolve the force in the direction PO . Writing $x = \cos \theta$, this becomes $2G \pi r^2 \rho \int_{-1}^1 (d - r x)/(d^2 + r^2 - 2dr x)^{3/2} dx$.

This is not as bad as it looks. It is just the sum of a $(1 - x)^{-1/2}$ and a $(1 - x)^{-3/2}$ integral, both of which are straightforward. Moreover, when we come to substitute $x = \pm 1$, the factor $(d^2 + r^2 - 2dr x)^{1/2}$ becomes just $d - r$ or $d + r$. So we get after a little simplification $4 \pi r^2 \rho / d^2 = MG/d^2$, which is the same result as if all the mass was concentrated at O .

(2) We can write a general line as $x = at + b$, $y = ct + d$, $z = et + f$, for some constants a, b, c, d, e, f and a parameter t which takes all real values. If this lies in $z = xy$, then $et + f = ac t^2 + (bc + ad) t + bd$ for all t . Hence a or c must be

zero. If a is 0, then $z = by$, so the line can be written as $x = b$, $z = by$. Similarly, if $c = 0$, then the line can be written as $y = d$, $z = dx$. Conversely, it is easy to see that these two families of lines lie in the surface.

Problem B1

Do either (1) or (2)

(1) Let A be matrix (a_{ij}) , $1 \leq i, j \leq 4$. Let $d = \det(A)$, and let A_{ij} be the cofactor of a_{ij} , that is, the determinant of the 3×3 matrix formed from A by deleting a_{ij} and other elements in the same row and column. Let B be the 4×4 matrix (A_{ij}) and let $D = \det B$. Prove $D = d^3$.

(2) Let $P(x)$ be the quadratic $Ax^2 + Bx + C$. Suppose that $P(x) = x$ has unequal real roots. Show that the roots are also roots of $P(P(x)) = x$. Find a quadratic equation for the other two roots of this equation. Hence solve $(y^2 - 3y + 2)^2 - 3(y^2 - 3y + 2) + 2 - y = 0$.

Solution

Answer: (2) The quadratic is $A^2x^2 + (AB + A)x + (AC + B + 1) = 0$. The quartic in y has roots 0, 1, 2, 2.

(1) We have $a_{i1}A_{i1} + a_{i2}A_{i2} + a_{i3}A_{i3} + a_{i4}A_{i4} = d$. But $a_{i1}A_{j1} + a_{i2}A_{j2} + a_{i3}A_{j3} + a_{i4}A_{j4} = 0$ for i not equal to j (because it can be considered as an expansion of the determinant for the matrix derived from A by replacing row i by row j - the resulting matrix has two identical rows and hence zero determinant). So if we multiply the transpose of A by the matrix (A_{ij}) then we get d down the diagonal and zeros elsewhere. Hence $dD = d^4$, so $D = d^3$.

(2) It is obvious that if $P(x) = x$, then $P(P(x)) = x$.

$P(P(x)) = x$ is $A(Ax^2 + Bx + C)^2 + B(Ax^2 + Bx + C) + C = x$. This is evidently a quartic and two of its roots are those of $Ax^2 + (B - 1)x + C = 0$. We could obtain the quadratic for the other two roots by multiplying out $P(P(x)) - x$ and factorising it. But it is sufficient to obtain the coefficients of x^4 , x^3 and x^0 . This gives us the sum of the four roots as $-2B/A$ and their product as $(AC + B + 1)C/A^3$. The sum and product of the two known roots are $-B/A - 1/A$ and C/A . Hence the sum and product of the other two roots are $-B/A + 1/A$ and $(AC + B + 1)/A^2$, so the roots are the roots of the quadratic $A^2x^2 + (AB + A)x + (AC + B + 1) = 0$.

$y^2 - 3y + 2 = 0$ has roots 1 and 2. So these values are also roots of $(y^2 - 3y + 2)^2 - 3(y^2 - 3y + 2) + 2 - y = 0$. The other two roots are also the roots of $x^2 + (-3 + 1)x + (2 - 3 + 1) = 0$. These are obviously 0 and 2.

Problem B2

Find all solutions of the differential equation $zz'' - 2z'z' = 0$ which pass through the point $x=1$, $z=1$.

Solution

Answer: $z = 1/(A(x - 1) + 1)$.

We have $z''/z' = 2z'/z$. Integrating, $\ln z' = 2 \ln z + \text{const}$, so $z' = -A/z^2$. Integrating again: $1/z = Ax + B$. But $z(1) = 1$, so $B = 1 - A$.

Problem B3

A horizontal disk diameter 3 inches rotates once every 15 seconds. An insect starts at the southernmost point of the disk facing due north. Always facing due north, it crawls over the disk at 1 inch per second. Where does it again reach the edge of the disk?

Solution

Answer: at the northernmost point of the disk.

Take polar coordinates with $r = 3/2$, $\theta = 0$ at the start. The equations of motion are $dr/dt = -\cos \theta$, $r d\theta/dt = 2\pi/15 + \sin \theta$.

Differentiating the second equation: $(dr/dt)(d\theta/dt) + r d^2\theta/dt^2 = (2\pi/15) dr/dt + (d\theta/dt) \cos \theta$. Substituting from the first equation, $2(dr/dt)(d\theta/dt) + r d^2\theta/dt^2 = (2\pi/15) dr/dt$. Multiplying through by r and integrating wrt t , we get $r^2 d\theta/dt = (\pi/15)r^2 + C$, for some constant C . At $t = 0$, $r = 3/2$ and $d\theta/dt = 2\pi/15$, so $C = 3\pi/20$. Thus $r^2 d\theta/dt = (\pi/15)r^2 + 3\pi/20$. But the original equation gives $r^2 d\theta/dt = 2r^2\pi/15 + r \sin \theta$. Hence $\pi r^2/15 + r \sin \theta = 3\pi/20$ (**). Hence if $r = \pm 3/2$, we have $\sin \theta = 0$ and hence $\theta = 0$ or π .

That is not quite enough to show that the insect reaches the edge again at $\theta = \pi$. But we can treat (**) as a quadratic in r and solve to get $r^2 = (\sin^2\theta + (\pi/5)^2)^{1/2} - \sin\theta$. This shows that r first decreases, but then increases again to ± 1 at $\theta = \pi$. We can rule out -1 because r^2 is always positive, and r starts positive. So by continuity it must always remain positive. Thus the insect next reaches the edge at the northernmost point of the disk.

Problem B4

The parabola P has focus a distance m from the directrix. The chord AB is normal to P at A . What is the minimum length for AB ?

Solution

Answer: $3\sqrt{3}m$.

We may take the equation of P as $2my = x^2$. The gradient at the point $A(a, a^2/2m)$ is a/m , so the normal at (a, b) is $(y - a^2/2m) = -m/a(x - a)$. Substituting in $2my = x^2$, it meets P at (x, y) where $x^2 + 2m^2/a x - (2m^2 + a^2) = 0$, so the other point B has $x = -(2m^2/a + a)$.

Thus $AB^2 = (2a + 2m^2/a)^2 + 4m^2(1 + m^2/a^2)^2 = 4a^2(1 + m^2/a^2)^3$. Differentiating, we find the minimum is at $a^2 = 2m^2$ and is $AB^2 = 27m^2$.

Problem B5

Find the locus of the foot of the perpendicular from the center of a rectangular hyperbola to a tangent. Obtain its equation in polar coordinates and sketch it.

Solution

Answer: $r^2 = 2k^2 \sin 2\theta$. It is a figure of 8 with its axis along the line $y = x$ and touching the x -axis and y -axis at the origin.

Take the hyperbola as $xy = k^2$. Then the tangent at $(a, k^2/a)$ is $(y - k^2/a) = -k^2/a^2(x - a)$. The perpendicular line through the origin is $y = a^2/k^2 x$. They intersect at $x = 2k^2/(a(a^2/k^2 + k^2/a^2))$, $y = 2a/(a^2/k^2 + k^2/a^2)$. So the polar coordinates r, θ satisfy $\tan\theta = y/x = a^2/k^2$, $r^2 = x^2 + y^2 = 4a^2(k^4/a^4 + 1)/(k^2/a^2 + a^2/k^2) = 4a^2(\cot^2\theta + 1)/(\tan\theta + \cot\theta)^2 = 4a^2 \cos^2\theta/(\sin^2\theta + \cos^2\theta)^2 = 4a^2 \cos^2\theta = 4k^2 \sin\theta \cos\theta = 2k^2 \sin 2\theta$. Thus the polar equation of the locus is $r^2 = 2k^2 \sin 2\theta$.

Problem B6

What is the shortest distance between the plane $Ax + By + Cz + 1 = 0$ and the ellipsoid $x^2/a^2 + y^2/b^2 + z^2/c^2 = 1$. You may find it convenient to use the notation $h = (A^2 + B^2 + C^2)^{-1/2}$, $m = (a^2A^2 + b^2B^2 + c^2C^2)^{1/2}$. What is the algebraic condition for the plane not to intersect the ellipsoid?

Solution

The tangent plane to the ellipsoid at (X, Y, Z) is $Xx/a^2 + Yy/b^2 + Zz/c^2 = 1$. It is parallel to $Ax + By + Cz + 1 = 0$ iff $X/a^2 = kA$, $Y/b^2 = kB$, $Z/c^2 = kC$ for some k . But $1 = X^2/a^2 + Y^2/b^2 + Z^2/c^2 = k^2(a^2A^2 + b^2B^2 + c^2C^2) = k^2m^2$, so $k = \pm 1/m$. There are two values corresponding to two parallel tangent planes (one on either side of the ellipse). The equation of the tangent plane is $k(Ax + By + Cz) = 1$.

The distance of the origin from the plane $Ax + By + Cz + 1 = 0$ is $1/(A^2 + B^2 + C^2)^{1/2} = h$. The distance of the origin from the tangent plane $k(Ax + By + Cz) = 1$ is $h/|k| = hm$. So if $m \geq 1$, the plane $Ax + By + Cz + 1 = 0$ lies between the two tangent planes and hence intersects the ellipse. So in this case the minimum distance is zero. If $m < 1$, then the distance between the plane $Ax + By + Cz + 1 = 0$ and the nearer tangent plane is $h(1 - m)$ and that is the required shortest distance.

2nd Putnam 1939

Problem A1

Let C be the curve $y^2 = x^3$ (where x takes all non-negative real values). Let O be the origin, and A be the point where the gradient is 1. Find the length of the curve from O to A .

Solution

Ans: $8/27 (2\sqrt{2} - 1)$. Trivial integration.

Problem A2

Let C be the curve $y = x^3$ (where x takes all real values). The tangent at A meets the curve again at B . Prove that the gradient at B is 4 times the gradient at A .

Solution

Trivial. [Take the point as (a, a^3) . Write down the equation of the tangent. Write down its point of intersection with the curve: $(x^3 - a^3) = 3a^2(x - a)$. We know this has a repeated root $x = a$. The sum of the roots is zero, so the third root is $x = -2a$. Finally, $3(-2a)^2 = 4$ times $3a^2$.]

Problem A3

The roots of $x^3 + ax^2 + bx + c = 0$ are α, β and γ . Find the cubic whose roots are $\alpha^3, \beta^3, \gamma^3$.

Solution

$$x^3 + (a^3 - 3ab + 3c)x^2 + (b^3 - 3abc + 3c^2)x + c^3 = 0.$$

A routine manipulation. Suppose the roots are α, β, γ . Then $\alpha + \beta + \gamma = -a$, $\alpha\beta + \beta\gamma + \gamma\alpha = b$, $\alpha\beta\gamma = -c$. So to get the coefficients of the desired polynomial we have to find the corresponding expressions in the cubes: $\alpha^3 + \beta^3 + \gamma^3$ etc. You obviously start with $(\alpha + \beta + \gamma)^3$ etc and then add additional terms to get the desired expressions.

Problem A4

Given 4 lines in Euclidean 3-space:

$$L_1: x = 1, y = 0;$$

$$L_2: y = 1, z = 0;$$

$$L_3: x = 0, z = 1;$$

$$L_4: x = y, y = -6z.$$

Find the equations of the two lines which both meet all of the L_i .

Solution

A routine computation. Assume the line meets L_1 at $(1, 0, a)$ and L_2 at $(b, 1, 0)$. Then it is $(x - 1) = t(x - b)$, $y = t(y - 1)$, $(z - a) = tz$. So it can only cut L_3 if $1/b = 1 - a$, and L_4 if $6a = 6ab - 1$. This gives a quadratic for a , which we can solve to get $a = -1/2$ or $1/3$. Hence the possible lines are $(1, 0, -1/2) + t(-1/3, 1, 1/2)$ and $(1, 0, 1/3) + t(1/2, 1, -1/3)$.

Problem A5

Do either (1) or (2)

(1) x and y are functions of t . Solve $x' = x + y - 3$, $y' = -2x + 3y + 1$, given that $x(0) = y(0) = 0$.

(2) A weightless rod is hinged at O so that it can rotate without friction in a vertical plane. A mass m is attached to the end of the rod A , which is balanced vertically above O . At time $t = 0$, the rod moves away from the vertical with negligible initial angular velocity. Prove that the mass first reaches the position under O at $t = \sqrt{(OA/g)} \ln(1 + \sqrt{2})$.

Solution

(1) Differentiate the first equation, use the second equation to eliminate y' , and then the (undifferentiated) first equation to eliminate y , giving:

$x'' - 4x' + 5x = 10$. Solving: $x = 2 + A e^{2t} \sin t + B e^{2t} \cos t$. But $x(0) = 0$, so $B = -2$. The first equation now gives $y = x' - x + 3 = 1 + (A + 2) e^{2t} \sin t + (A - 2) e^{2t} \cos t$. But $y(0) = 1$, so $A = 1$. The final solution is thus: $x = 2 + e^{2t} \sin t - 2 e^{2t} \cos t$; $y = 1 + 3 e^{2t} \sin t - e^{2t} \cos t$.

(2) Trivial, except for the integral, which is moderately hard, unless you happen to know it.

Let the angle the rod makes with the (upward) vertical be θ . Conservation of energy gives immediately: $\frac{1}{2} OA^2(d\theta/dt)^2 = OA \cdot g(1 - \cos \theta)$.

Now for the first time in this exam we come up against something that is not completely obvious. How do we do the integral?

You need the half-angle formulae, eg $(1 - \cos \theta) = 2 \sin^2 \theta/2$. Now if you can remember the integral for $1/\sin z$ (eg $\ln \sin z - \ln(1 + \cos z)$, or equivalently $\ln(\operatorname{cosec} z - \cot z)$), then you are home.

If not, use the half-angle formulae again: $\sin \theta/2 = 2 \sin \theta/4 \cos \theta/4$. Putting $c = \cos \theta/4$, we have to integrate $1/(c(1 - c^2))$. Expand using partial fractions and now the integral is just a sum of logs.

Problem A6

Do either (1) or (2):

(1) A circle radius r rolls around the inside of a circle radius $3r$, so that a point on its circumference traces out a curvilinear triangle. Find the area inside this figure.

(2) A frictionless shell is fired from the ground with speed v at an unknown angle to the vertical. It hits a plane at a height h . Show that the gun must be sited within a radius $v/g(v^2 - 2gh)^{1/2}$ of the point directly below the point of impact.

Solution

(1) This is moderately difficult. It is not immediately obvious what coordinates to use (or at least, after meeting an integral I did not immediately recognize, I started worrying that there might be a better choice of coordinates), and it is not immediately obvious how to do the resulting integral.

Let C be the center of the large circle and let O be the initial point of contact between the two circles. Take O as the origin and OC as the x -axis, take the y -axis so that P the point of contact gets a positive y -coordinate just after rolling starts. The easiest parameter is to take angle $OCP = \theta$. Then it is not hard to see that $x/r = 3 - 2 \cos \theta - \cos 2\theta$, $y/r = 2 \sin \theta - \sin 2\theta$.

Evidently we need something like $\int y dx$. We need a little care on the limits of integration. Let A, B be the other vertices of the curvilinear triangle (A corresponding to $\theta = 2\pi/3$, B to $\theta = 4\pi/3$). Let X be the point where the curve AB cuts the x -axis and Y the point where the line AB cuts the x -axis. \int_O^A gives the area under the curve OA , in other words the area OAX plus the area AXY . Then \int_A^X gives minus the area AXY (because x is decreasing, so dx is negative). \int_X^B gives minus area BXY (x increasing, but y negative), and \int_B^O gives plus area BXY plus area OBX (x decreasing and y negative). So the entire integral $\int_{\theta=0}^{2\pi}$ gives the required area inside the curvilinear triangle OAB . In other words we need:

$$2r^2 \int_{\theta=0}^{2\pi} (2 \sin \theta - \sin 2\theta)(\sin \theta + \sin 2\theta) d\theta.$$

This is the point at which we are likely to get stuck. Changing variable to $z = \cos \theta$ does not apparently help. The trick is to put things in terms of $\sin nx$ or $\cos nx$. We may remember that $\sin^2 z = (1 - \cos 2z)/2$, and $\cos(w \pm z) = \cos w \cos z \mp \sin w \sin z$, so that $\sin z \sin 2z = (\cos z - \cos 3z)/2$.

Expanding the integrand gives: $2 \sin^2 \theta + \sin \theta \sin 2\theta - \sin^2 2\theta$. Using the two formulae above transforms this to: $1 - \cos 2\theta + (\cos \theta - \cos 3\theta)/2 - 1/2 + 1/2 \cos 4\theta$. The \cos terms all integrate to zero and the constant term $1/2$ integrates to π , so the final answer is $2\pi r^2$.

(2) There is a slight trap here. We may be tempted to argue that the extreme case is where the shell reaches the plane with zero vertical velocity. The horizontal velocity does not change during the trajectory, so taking u as the horizontal velocity and w as the vertical, we can write down immediately that $w^2 = 2gh$ (energy) and hence the radius $r = w/g(v^2 - 2gh)^{1/2}$, which is unfortunately wrong. Notice that it is smaller than the answer sought. The reason is that impact at zero vertical velocity is not the extreme case. We can usually do better by having the shell peak before the plane, so that it hits it on the way down - the extra time to travel horizontally outweighs the loss of horizontal velocity.

So we have to write down the equations: $r = tu$, $h = tw - gt^2/2$. Squaring to eliminate u, w in favor of v , gives:

$$g^2 t^4/4 + (gh - v^2)t^2 + (h^2 + r^2) = 0. (*)$$

For this to have a real root we require $(gh - v^2)^2 \geq g^2(h^2 + r^2)$ and hence $r \leq v/g (v^2 - 2gh)^{1/2}$.

We are not asked to prove that the plane can be hit from anywhere within this radius, so we could stop here. But $v^2 \geq 2gh > gh$, so (*) is a quadratic of the form $a z^2 - b z + c = 0$, with a, b, c positive. Hence if r satisfies the condition, it has two *positive* roots for z and (*) has two real (positive) roots for t . Having solved for t , we can then solve for the angle (or equivalently for u, w), which shows that we can hit the plane from anywhere inside the radius.

Problem A7

Do either (1) or (2):

(1) Let C_a be the curve $(y - a^2)^2 = x^2(a^2 - x^2)$. Find the curve which touches all C_a for $a > 0$. Sketch the solution and at least two of the C_a .

(2) Given that $(1 - hx)^{-1}(1 - kx)^{-1} = \sum_{i \geq 0} a_i x^i$, prove that $(1 + hkx)(1 - hkx)^{-1}(1 - h^2x)^{-1}(1 - k^2x)^{-1} = \sum_{i \geq 0} a_i^2 x^i$.

Solution

(1) It is not hard to see that C_1 is a figure 8 lying on its side with the double point at $(1,0)$, vertical tangents at $(1,1)$ and $(-1,1)$ and horizontal tangents at $(1/\sqrt{2}, 1/2)$, $(1/\sqrt{2}, 3/2)$, $(-1/\sqrt{2}, 1/2)$, $(-1/\sqrt{2}, 3/2)$. C_a is obtained from C_1 by transforming (x,y) to (ax, a^2y) (so stretching by a factor a in the x -direction and a factor a^2 in the y -direction). So we get a sequence of ever-larger horizontal 8s centered ever-further up the y -axis. This suggests that the envelope is something like a parabola $y = k x^2$.

At this point it is distinctly helpful to know more classical differential geometry than today's undergraduate. The classical method for finding the envelope for a 1-parameter family of curves (which *usually* works) is to differentiate wrt the parameter. Thus we eliminate a from:

$$(y - a^2)^2 - x^2(a^2 - x^2) = 0, \text{ and } 4a(y - a^2) + 2a x^2 = 0$$

giving $y = 3/4 x^2$ (or the y -axis, which is presumably a spurious solution arising from the double points).

Alternatively, we might guess that the envelope is a curve $y = k x^2$ for some k . The intersection of this with C_a is given by:

$$(k^2 + 1)x^4 - (2k + 1) a^2 x^2 + a^4 = 0, \text{ or } x^2 = (2k + 1)a^2 / (2k^2 + 2) \pm a^2(4k - 3)^{1/2} / (2k^2 + 2).$$

For this to be a tangent we need double roots and hence $k = 3/4$. It is now easy to check that this parabola meets C_a at $(2a/\sqrt{5}, 3a^2/5)$, $(-2a/\sqrt{5}, 3a^2/5)$ and to check that the gradients match.

(2) Fairly easy.

This is obviously not a general result (whereby we derive $\sum a_i^2 x^i$ from $\sum a_i x^i$), so we need to evaluate the a_i . In fact, it is easily seen that $a_i = (h^{i+1} - k^{i+1}) / (h - k)$. [For example, multiply the expansions of $(1 - hx)^{-1}$ and $(1 - kx)^{-1}$ to get $a_i = h^i + h^{i-1}k + \dots + h k^{i-1} + k^i = (h^{i+1} - k^{i+1}) / (h - k)$.]

Multiplying across to try to show that higher powers of $(1 - hkx)(1 - h^2x)(1 - k^2x) \sum a_i^2 x^i$ have zero coefficients is a mistake (doable, but much algebra). It is better to evaluate $\sum a_i^2 x^i$ directly. After substituting for a_i , we are going to get terms of the form $\sum z^i$ which evaluates immediately to $(1 - z)^{-1}$, giving the expression in the question in partial fraction form.

Problem B1

The points $P(a,b)$ and $Q(0,c)$ are on the curve $y/c = \cosh(x/c)$. The line through Q parallel to the normal at P cuts the x -axis at R . Prove that $QR = b$.

Solution

Trivial. [Let O be the origin. Then $OR/OQ = \sinh a/c$, so $QR^2 = c^2(1 + \sinh^2 a/c)$, so $QR = b$.]

Problem B2

Evaluate $\int_1^3 ((x-1)(3-x))^{-1/2} dx$ and $\int_1^{\infty} (e^{x+1} + e^{3-x})^{-1} dx$.

Solution

(1) π . Trivial. [Write $(x - 1)(3 - x) = 1 - (x - 2)^2$. So we have a standard $\sin^{-1}z$ integral.]

(2) $\pi/(4e^2)$. Trivial. [Multiply top and bottom by e^x . Change variable to $y = e^{x-1}$. We now have a standard $\tan^{-1}z$ integral.]

Problem B3

Given $a_n = (n^2 + 1)3^n$, find a recurrence relation $a_n + p a_{n+1} + q a_{n+2} + r a_{n+3} = 0$. Hence evaluate $\sum_{n \geq 0} a_n x^n$

Solution

We can solve formally to get the recurrence relation, but it is quicker to get there informally. We look for a relation between $b_n = a_n/3^n$, $b_{n+1} = a_{n+1}/3^{n+1}$, $b_{n+2} = a_{n+2}/3^{n+2}$, $b_{n+3} = a_{n+3}/3^{n+3}$, because that takes care of the powers of 3. So, ignoring the 3^n , we are looking at:

$$\begin{aligned} n^2 + 1 \\ n^2 + 2n + 2 \\ n^2 + 4n + 5 \\ n^2 + 6n + 10 \end{aligned}$$

We try to get a linear combination of the first three which is constant. But that is easy: subtracting twice the second from the third gets rid of the n term, then adding the first gets rid of the n^2 term. So, $b_{n+2} - 2b_{n+1} + b_n = 2.3^n$. But $b_{n+3} - 2b_{n+2} + b_{n+1}$ has the *same* value, so subtracting:

$$a_{n+3} - 9a_{n+2} + 27a_{n+1} - 27a_n = 0, \text{ which is the required recurrence relation.}$$

Let the power series sum to y . Then taking $y - 9x + 27x^2y - 27x^3y$ will give $a_{n+3} - 9a_{n+2} + 27a_{n+1} - 27a_n$ as the coefficient of x^{n+3} , so we need only worry about the early terms: $a_0 + (a_1 - 9a_0)x + (a_2 - 9a_1 + 27a_0)x^2 = (1 - 3x + 18x^2)/(1 - 9x + 27x^2 - 27x^3)$.

Using the ratio test, the original series evidently converges for $|x| < 1/3$, which may prompt us to notice that $1 - 9x + 27x^2 - 27x^3 = (1 - 3x)^3$.

That in turn may prompt us to try solving the problem backwards. We know that:

$$1/(1 - z) = \sum z^n; 1/(1 - z)^2 = \sum (n+1)z^n; 1/(1 - z)^3 = \sum (n+1)(n+2)/2 z^n.$$

Hence $2/(1 - z)^3 - 3/(1 - z)^2 + 2/(1 - z) = \sum (n^2 + 1)z^n$. Replacing z by $3x$ gives $\sum a_n x^n = (1 - 3x + 18x^2)/(1 - 3x)^3$. Multiplying across by $(1 - 3x)^3$ now gives the required recurrence relation.

Problem B4

The *axis* of a parabola is its axis of symmetry and its *vertex* is its point of intersection with its axis. Find: the equation of the parabola which touches $y = 0$ at $(1,0)$ and $x = 0$ at $(0,2)$; the equation of its axis; and its vertex.

Solution

The general equation of the a parabola is: $(ax + by)^2 + cx + dy + e = 0$. Its intersection with $y = 0$ is given by $a^2x^2 + cx + e = 0$. This must have a double root, so $c = -2a^2$, $e = -a^2$. Considering the other tangent, we find: $d = -4b^2$, $e = 4b^2$. So (up to an irrelevant constant factor) we have: $a = 2$, $b = 1$, $c = -8$, $d = -4$, $e = 4$; or $a = 2$, $b = -1$, $c = -8$, $d = -4$, $e = 4$. But in the first case the equation can be written as $(2x + y - 2)^2 = 0$, which is a double line. It is debatable whether this qualifies as a parabola, but it would not normally be said to *touch* the points $(1,0)$ and $(0,2)$. So we are left with the parabola: $(2x - y)^2 - 8x - 4y + 4 = 0$.

We want to put this in the form $u = kv^2$. The line $x + 2y = 0$ is perpendicular to the line $2x - y = 0$, so we change variables to $X = 2x - y$, $Y = x + 2y$, giving: $X^2 - 16/5 Y - 12/5 X + 4 = 0$, or $16/5 (Y - 4/5) = (X - 6/5)^2$, which is the equation of a parabola with vertex $X = 6/5$, $Y = 4/5$, axis $X = 6/5$. Changing back to the original coordinates, $x = (2X + Y)/5$, $y = (2Y - X)/5$, the vertex is $(16/25, 2/25)$ and the axis is $10x - 5y = 6$.

Problem B5

Do either (1) or (2):

(1) Prove that $\int_1^k [x] f'(x) dx = [k] f(k) - \sum_1^{[k]} f(n)$, where $k > 1$, and $[z]$ denotes the greatest integer $\leq z$. Find a similar expression for: $\int_1^k [x^2] f'(x) dx$.

(2) A particle moves freely in a straight line except for a resistive force proportional to its speed. Its speed falls from 1,000 ft/s to 900 ft/s over 1,200 ft. Find the time taken to the nearest 0.01 s. [No calculators or log tables allowed!]

Solution

(1) $[x]$ is constant over the interval $[i, i+1)$ for i an integer, so we split the range of integration to get $\int_1^k = \int_{[k]}^k + \int_1^2 + \int_2^3 + \dots + \int_{[k]-1}^{[k]}$. We can write down each of these integrals, collect terms and get the result.

The same idea works the second integral, except that we divide at $\sqrt{2}, \sqrt{3}, \sqrt{4}, \dots, \sqrt{[k^2]}$, giving the result: $[k^2] f(k) - (f(1) + f(\sqrt{2}) + f(\sqrt{3}) + \dots + f(\sqrt{[k^2]}))$.

(2) Easy. 1.26 s.

The equation of motion is $x'' = -kx'$. Integrating: $x' = kA e^{-kt}$. Integrating again, and putting $x(0) = 0$, $x = A(1 - e^{-kt})$. Suppose T is the required time. Then from the speed $e^{-kT} = 0.9$. $x(T) = 1200$, so $A = 12000$. The initial speed is 1000, so $k = 1/12$ and $T = -12 \ln 0.9$. The only slight snag is that in the exam calculators did not exist and log tables were not allowed. So we have to use the expansion $\ln(1+x) = x - x^2/2 + x^3/3 - \dots$, or more usefully: $-\ln(1-x) = x + x^2/2 + x^3/3 + \dots$, giving $T = 1.2 + 0.06 + 0.004 + 0.0003 + \dots$ or 1.26 s.

Problem B6

Do either (1) or (2):

(1) f is continuous on the closed interval $[a, b]$ and twice differentiable on the open interval (a, b) . Given $x_0 \in (a, b)$, prove that we can find $\xi \in (a, b)$ such that $((f(x_0) - f(a))/(x_0 - a) - (f(b) - f(a))/(b - a))/(x_0 - b) = f''(\xi)/2$.

(2) AB and CD are identical uniform rods, each with mass m and length $2a$. They are placed a distance b apart, so that $ABCD$ is a rectangle. Calculate the gravitational attraction between them. What is the limiting value as a tends to zero?

Solution

(1) We obviously have to use the mean value theorem. So we need to construct a suitable auxiliary function to apply it to. It is usually easiest to apply the MVT in cases where the function has equal values at the two ends of the interval. So we want to find some function g , such that g' has equal values at two different points. Let the value of the expression given, $((f(x_0) - f(a))/(x_0 - a) - (f(b) - f(a))/(b - a))/(x_0 - b)$, be y_0 . Then we are looking for $g''(x)$ to be something like $1/2 f''(x) - y_0$.

Let us start by rearranging the expression for y_0 to give $f(x_0) = f(a) + ((f(b) - f(a))(x_0 - a)/(b - a) + y_0(x_0 - a)(x_0 - b))$. After a little experimentation we may try looking at:

$$g(x) = f(x) - f(a) - ((f(b) - f(a))(x - a)/(b - a) - y_0(x - a)(x - b)).$$

We notice that $g(a) = 0$, $g(b) = 0$, $g(x_0) = 0$, $g'(x) = f'(x) - ((f(b) - f(a))/(b - a) - y_0(2x - a - b))$, $g''(x) = f''(x) - 2y_0$. At this point we should realize that we are home, because we have to show that we can find ξ such that $g''(\xi) = 0$. But the mean value theorem gives us a value in the interval (a, x_0) at which g' is zero and another in (x_0, b) . Hence there must be a value between the two (and *a fortiori* in (a, b)) at which g'' is zero.

(2) Straightforward, apart from an awkward integral. Answer: $Gm^2(1 - (1 + 4a^2/b^2)^{1/2})/(2a^2)$, which tends to Gm^2/b^2 as the rods shorten to become point masses.

By symmetry the net force must be perpendicular to the rods, so we just calculate the perpendicular component. Take coordinates x along one rod and y along the other. Then we can write down immediately that the perpendicular component is:

$$Gm^2/(4a^2) \int (b^2 + (x - y)^2)^{-1} b (b^2 + (x - y)^2)^{-1/2} dx dy.$$

The integrand is of the form $(1+z^2)^{-3/2}$. It helps to know that this integrates to $z/(1+z^2)^{1/2}$ (as is easily checked). People used to mess around with trigonometric substitutions (eg $\tan \theta$ reduces it to $\cos \theta$, which integrates immediately to $\sin \theta$, which one has to remember to think of as $\tan \theta / \sec \theta$). But it was always easier simply to learn a large number of integrals by rote. Nowadays, of course, one tends to look them up or use Mathematica/Maple.

So integrating first by y , we get $Gm^2/(4a^2) \int dx/b \{ (x-2a)/b (1+(x-2a)^2/b^2)^{-1/2} - x/b (1+x^2/b^2)^{-1/2} \}$. This is much less horrible than it looks, because we are just faced with integrating $z/(1+z^2)^{1/2}$ which is obviously $(1+z^2)^{1/2}$. So we get the expression above. For the limit, just expand the square root as a power series $1 + 2a^2/b^2 - a^4/(4b^4) + \dots$.

Problem B7

Do either (1) or (2):

(1) Let $a_i = \sum_{n=0}^{\infty} x^{3n+i}/(3n+i)!$ Prove that $a_0^3 + a_1^3 + a_2^3 - 3a_0a_1a_2 = 1$.

(2) Let O be the origin, λ a positive real number, C be the conic $ax^2 + by^2 + cx + dy + e = 0$, and C_λ the conic $ax^2 + by^2 + \lambda cx + \lambda dy + \lambda^2 e = 0$. Given a point P and a non-zero real number k , define the transformation $D(P,k)$ as follows. Take coordinates (x',y') with P as the origin. Then $D(P,k)$ takes (x',y') to (kx',ky') . Show that $D(O,\lambda)$ and $D(A,-\lambda)$ both take C into C_λ , where A is the point $(-c\lambda/(a(1+\lambda)), -d\lambda/(b(1+\lambda)))$. Comment on the case $\lambda = 1$.

Solution

(1) Moderately hard, unless differentiating is a reflex, in which case it is easy.

The series are all absolutely convergent for all x , so we can carry out whatever operations we want. But it is not at all obvious what to do. Multiplying out the power series to get a complicated sum for the coefficient of x^n is offputting. Much scope for algebraic error, and no guarantee that the eventual simplification will be obvious. So we look for some trick. The a_i are closely related to the exponential series, eg $a_0 + a_1 + a_2 = e^x$, and the prevalence of 3 in the question may eventually suggest looking at ω , the cube root of 1. Indeed, $a_0 + \omega a_1 + \omega^2 a_2 = e^{\omega x}$. Since ω^2 is also a root, we also have $a_0 + \omega^2 a_1 + \omega a_2 = e^{\omega^2 x}$. Remembering that $1 + \omega + \omega^2 = 0$ might prompt us to multiply these three expressions together, but it helps to remember the product of the three left hand sides: $(a_0 + a_1 + a_2)(a_0 + \omega a_1 + \omega^2 a_2)(a_0 + \omega^2 a_1 + \omega a_2) = a_0^3 + a_1^3 + a_2^3 - 3a_0a_1a_2$. Otherwise, you are faced with multiplying this out the hard way: after collecting terms, you get $(a_0^3 + a_1^3 + a_2^3)$, then 6 expressions of the type $a_0^2(1 + \omega + \omega^2)$, which are all zero, and $3a_0a_1a_2(\omega + \omega^2)$, which is $-3a_0a_1a_2$.

A more general approach, which is more likely to work, is to differentiate the expression $a_0^3 + a_1^3 + a_2^3 - 3a_0a_1a_2$. Provided you notice that $a_0' = a_2$ etc, this gives the result almost immediately (the derivative is zero, so the expression must be constant, but its value for $x = 0$ is 1).

(2) Trivial. [$D(O,\lambda)$ takes (x,y) to $(\lambda x, \lambda y)$. So if (x,y) satisfies the equation for C , just check that $(\lambda x, \lambda y)$ satisfies the equation for C_λ . Similarly, $D(A,-\lambda)$ takes (x,y) to $(-\lambda x - \lambda c/a, -\lambda y - \lambda d/b)$. Again, just check by substituting this into the equation for C_λ and using the fact that (x,y) satisfies the equation for C . If $\lambda = 1$, then $C_\lambda = C$, $D(O,\lambda)$ is the identity transformation, and $D(A,-\lambda)$ the central symmetry.]

3rd Putnam 1940

Problem A1

$p(x)$ is a polynomial with integer coefficients. For some positive integer c , none of $p(1), p(2), \dots, p(c)$ are divisible by c . Prove that $p(b)$ is not zero for any integer b .

Solution

Suppose $p(b) = 0$. Then $p(x) = (x - b)q(x)$, where q is a polynomial with integer coefficients. Put $b = cd + r$, where $1 \leq r \leq c$ (note that this is different from the conventional $0 \leq r < c$, but still possible because $1, 2, \dots, c$ are a complete set of residues mod c). Then $p(r) = p(b - cd) = -cd q(r)$ which is divisible by c . Contradiction.

Problem A2

$y = f(x)$ is continuous with continuous derivative. The arc PQ is concave to the chord PQ . X is a point on the arc PQ for which $PX + XQ$ is a maximum. Prove that XP and XQ are equally inclined to the tangent at X .

Solution

Take s to be the arc-length PX and z to be $PX + XQ$. Suppose the tangent is WXY , so that angle $WXP = \theta$ and angle $YXQ = \phi$. Then to first order if we vary X , the change $\delta z = \delta s (\cos \theta - \cos \phi)$, which is can be made positive (by choosing the sign of δs appropriately), thus contradicting the maximality of X , unless $\theta = \phi$.

The official solution. If X is a point of S such that $PX + XQ \geq PY + YQ$ for all points Y of S , then we can easily show that S lies in the half-plane bounded by external angle bisector of PXQ . If X lies on PQ , then S must be a subset of the segment PQ and the result is trivial. So assume it does not. Now reflect Q in the bisector to get Q' , with PXQ' a straight line. Then if Y is any point in the same half-plane as Q' , we have $YQ' < YQ$ and hence $PY + YQ > PY + YQ' \geq PQ' = PX + XQ' = PX + XQ$, so Y is not in S .

This argument depends upon X achieving a global maximum. The original wording of the question, "a point ... for which ... is a maximum" (which on this point I quoted exactly), is somewhat ambiguous. Does it mean a local or a global maximum? If it means a local, then we have to take S to be a small arc for which the maximum is global. Then that arc lies on one side of the line. Since it also has a point in common (X) and is differentiable, the line must be the tangent to the arc at X .

This solution is somewhat harder and less obvious (unless you have seen it before), so I prefer the simpler solution.

Problem A3

α is a fixed real number. Find all functions $f: \mathbb{R} \rightarrow \mathbb{R}$ (where \mathbb{R} is the reals) which are continuous, have a continuous derivative, and satisfy $\int_b^y f^\alpha(x) dx = (\int_b^y f(x) dx)^\alpha$ for all y and some b .

Solution

Straightforward to get the basic idea, but care is needed with the details. It is quite hard to get the answer exactly right (the official solution gives a spurious solution for the case (2) - overlooking the problem with the integration limits).

Answer:

- (1) No solutions for $\alpha = 0$;
- (2) No solutions for $\alpha < 0$;
- (3) Any continuous f with continuous derivative is a solution for $\alpha = 1$;
- (4) For $\alpha > 0$ and not of the form p/q , with p and q odd positive integers, $f = A e^{kx}$, where A is any real, and k is the positive real value of $\alpha^{1/(\alpha-1)}$;
- (5) For $\alpha = p/q$, with p and q odd positive integers (but not both 1), $f = A e^{kx}$ or $A e^{-kx}$, where A is any real, and k is the positive real value of $\alpha^{1/(\alpha-1)}$.

Case (3) is obvious. Case (1) is almost obvious: if $\alpha = 0$, then the lhs varies with y ($= y - b$), but the rhs does not ($= 1$), so there are no solutions. Assume now that α is not 0 or 1.

Differentiate wrt y and put $g(y) = \int_b^y f(x) dx$. Then $f = g'$ and we have $f^\alpha = \alpha g^{\alpha-1} f$. So $f = k g$, where $k = \alpha^{1/(\alpha-1)}$. Since $f = g'$, we can integrate immediately to get $f(x) = A e^{kx}$ (*).

However, we have to consider how many real values k can have. If $\alpha > 0$, then k certainly has a positive value, but we can also take the corresponding negative value if $1/(\alpha - 1)$ involves an even root, in other words if $\alpha = p/q$ with p and q both odd. Finally, (*) is clearly necessary, but not necessarily sufficient, so we have to substitute (*) back into the original equation. For $k > 0$, we find the solution works provided $b = -\infty$. If $k < 0$, then it works with $b = \infty$. This gives cases (4) and (5). [Note, however, that we can only allow A negative for α not of the form p/q with p and q odd, provided we are content for both sides of the original equation to have complex values (even though f is real valued).]

For $\alpha < 0$, all values of $\alpha^{1/(\alpha-1)}$ are complex unless $\alpha = -p/q$ with p an even positive integer and q an odd positive integer. If α has that form, then we may take $k = -1/|\alpha|^{1/(\alpha+1)}$. But now there is a problem with b . Taking $A = 1$ for simplicity, the $f^\alpha(x) = e^{|\alpha k| x}$, so the lhs = const $(e^{|\alpha k| y} - e^{|\alpha k| b})$ and the rhs = const $/(e^{-|k| y} - e^{-|k| b})^{|\alpha|}$. The constants are the same, but we need $b = -\infty$ to get rid of the b term on the lhs and $b = \infty$ to get rid of the term on the rhs. So there are no solutions in case (2).

Problem A4

p is a positive constant. Let R is the curve $y^2 = 4px$. Let S be the mirror image of R in the y -axis ($y^2 = -4px$). R remains fixed and S rolls around it without slipping. O is the point of S initially at the origin. Find the equation for the locus of O as S rolls.

Solution

Answer: $x(x^2 + y^2) + 2p y^2 = 0$.

Take the point of contact as (X, Y) , so $Y^2 = 4pX$. The tangent at to R at this point has gradient $2p/Y$, and hence has equation $(y - Y) = 2p/Y (x - X)$. The perpendicular to the tangent through the origin has equation $x = -2p/Y y$. If (x, y) is their point of intersection, then $(2x, 2y)$ is the point O (since S in its new position is the reflection of R in the tangent).

Solving for x, y : $y = Y^3/(2(Y^2 + 4p^2))$, $x = -2p/Y = -pY/(Y^2 + 4p^2)$. So the point O is $(-2pY^2/(Y^2 + 4p^2), Y^3/(Y^2 + 4p^2))$ (*). Using $Y = -2py/x$, we get $x = -2py^2/(x^2 + y^2)$, or $x(x^2 + y^2) + 2p y^2 = 0$ (**). We have shown that the locus is given by (*) and that all points on (*) are on (**). However, we must check that (**) does not include additional points. Writing (**) as $y^2 = -x^3/(x + 2p)$, shows that for each value of x in the range $(-2p, 0)$ there are exactly two possible values of y , and the only other point on (**) is the origin. Inspection shows that the same is true for (*), so the two expressions are equivalent.

Problem A5

Prove that the set of points satisfying $x^4 - x^2 = y^4 - y^2 = z^4 - z^2$ is the union of 4 straight lines and 6 ellipses.

Solution

$x^4 - x^2 = (x^2 - 1/2)^2 - 1/4$, so $x^4 - x^2 = y^4 - y^2$ is equivalent to $(x^2 - 1/2)^2 = (y^2 - 1/2)^2$ and hence to $x^2 - 1/2 = \pm(y^2 - 1/2)$, which is equivalent to $x = y$, or $x = -y$, or $x^2 + y^2 = 1$. Similarly, for $y^4 - y^2 = z^4 - z^2$. So the equation given is equivalent to: (1) $x = y$ and $y = z$, or (2) $x = y$ and $y = -z$, or (3) $x = -y$ and $y = z$, or (4) $x = -y$ and $y = -z$, or (5) $x = y$ and $y^2 + z^2 = 1$, or (6) $x = -y$ and $y^2 + z^2 = 1$, or (7) $x^2 + y^2 = 1$ and $y = z$, or (8) $x^2 + y^2 = 1$ and $y = -z$, or (9) $x^2 + y^2 = 1$ and $y^2 + z^2 = 1$.

Clearly (1) - (4) are straight lines. (5) is the intersection of a plane and a cylinder, which is an ellipse. Similarly (6), (7) and (8). (9) is slightly harder to see. If one's visualization is good, then one can see that the intersection of two cylinders with the same radius and axes intersecting at right angles is two perpendicular ellipses with a common minor axis. Otherwise, subtracting the two equations we see that $x = z$ or $-z$ and the intersection is also given by the intersection of a cylinder with two planes.

Problem A6

$p(x)$ is a polynomial with real coefficients and derivative $r(x) = p'(x)$. For some positive integers a, b , $r^a(x)$ divides $p^b(x)$. Prove that for some real numbers A and α and for some integer n , we have $p(x) = A(x - \alpha)^n$.

Solution

Write $p(x) = A \prod (x - \alpha_i)^{n_i}$.

Then $r(x) = p(x) \sum n_i/(x - \alpha_i) = (A \prod (x - \alpha_i)^{n_i - 1}) q(x)$, where no α_i is a root of $q(x)$. This is easily seen, because $q(x)$

is a sum of terms, all but one of which has a factor $(x - \alpha_i)$. But $q(x)$ divides $p(x)^b$ which has no roots except the α_i . Hence $q(x)$ must be a constant. But now the degree of $r(x)$ is wrong unless there is just one α_i .

Problem A7

a_i and b_i are real, and $\sum_{i=1}^{\infty} a_i^2$ and $\sum_{i=1}^{\infty} b_i^2$ converge. Prove that $\sum_{i=1}^{\infty} (a_i - b_i)^p$ converges for $p \geq 2$.

Solution

Notice first that it is sufficient to prove the result for $p = 2$. For that is equivalent to the statement that $\sum |a_i - b_i|^2$ converges. Hence for sufficiently large i , $|a_i - b_i| < 1$, and hence $|a_i - b_i|^p \leq |a_i - b_i|^2$. So $\sum (a_i - b_i)^p$ is absolutely convergent and hence convergent.

$(a_i - b_i)^2 = a_i^2 - 2a_i b_i + b_i^2$. The only tricky part is the middle term. It may be positive, so we cannot simply argue that $0 \leq (a_i - b_i)^2 \leq a_i^2 + b_i^2$. However, it is true that $0 \leq (a_i - b_i)^2 = 2a_i^2 + 2b_i^2 - (a_i + b_i)^2 \leq 2a_i^2 + 2b_i^2$. That suffices, since $\sum a_i^2$ and $\sum b_i^2$ are absolutely convergent, hence also $\sum (2a_i^2 + 2b_i^2)$.

Problem A8

Show that the area of the triangle bounded by the lines $a_i x + b_i y + c_i = 0$ ($i = 1, 2, 3$) is $\Delta^2 / [2(a_2 b_3 - a_3 b_2)(a_3 b_1 - a_1 b_3)(a_1 b_2 - a_2 b_1)]$, where Δ is the 3×3 determinant with columns a_i, b_i, c_i .

Solution

Fairly easy if you remember some formulae for determinants (which people did in those days). Of course, you can just slog through the expression (*) below in terms of a_i, b_i, c_i . That is doable, but completely mindless, the only required skill is doing elementary algebra fast without mistakes. I suppose I am fairly out of sympathy with the rather common Putnam style of problem where the basic idea is obvious, but you have to be skilful at evaluating integrals, determinants etc, often using tricks.

Take A_i be the cofactor of a_i in Δ . Similarly, B_i and C_i . [So, for example, $A_1 = b_2 c_3 - b_3 c_2$, $A_2 = b_3 c_1 - b_1 c_3$, $A_3 = b_1 c_2 - b_2 c_1$.]

The lines $a_2 x + b_2 y + c_2 = 0$ and $a_3 x + b_3 y + c_3 = 0$ intersect at $(A_1/C_1, B_1/C_1)$. Similarly, the other two points of intersection are $(A_2/C_2, B_2/C_2)$ and $(A_3/C_3, B_3/C_3)$.

The area of the triangle is therefore (the absolute value of) the determinant K with rows $A_1/C_1, B_1/C_1, 1; A_2/C_2, B_2/C_2, 1; A_3/C_3, B_3/C_3, 1$. (*) [For example, take a z -coordinate perpendicular to the plane, and take the cross product of the vectors along two sides.] But $K [C_1 C_2 C_3]$ is the determinant whose elements are the cofactors of the original determinant. This has value equal to Δ^2 . For example, on multiplying it by Δ , we get Δ down the diagonal and zeros elsewhere, and hence Δ^3 .

Problem B1

A stone is thrown from the ground with speed v at an angle θ to the horizontal. There is no friction and the ground is flat. Find the total distance it travels before hitting the ground. Show that the distance is greatest when $\sin \theta \ln(\sec \theta + \tan \theta) = 1$.

Solution

Take coordinates (time and distance) zeroed on the peak of the trajectory. After time t , the stone travels a distance $x = v t \cos \theta$ horizontally and a distance $1/2 g t^2$ vertically. So the trajectory is a parabola $2y = k x^2$, where $k = g/(v \cos \theta)^2$. The stone is on the ground at $t = \pm v/g \sin \theta$, at a horizontal distance $\pm a$ from the peak, where $a = v^2/g \sin \theta \cos \theta$.

The length of the parabola $y = k/2 x^2$ between $x = -a$ and $x = a$ is $2 \int_0^a (1 + k^2 x^2)^{1/2} dx$.

To do the integral it helps to remember that $1 + \sinh^2 z = \cosh^2 z$. So substituting $kx = \sinh z$, will essentially give us the integral of $\cosh^2 z$. That is doable, using the analog of the double angle formulae. So setting $I = \int (1 + k^2 x^2)^{1/2} dx$, and substituting $x = \sinh z$, we have $I = 1/k \int \cosh^2 z dz = 1/(2k) \int (\cosh 2z + 1) dz = 1/(4k) \sinh 2z + 1/(2k) z = 1/(2k) \sinh z \cosh z + 1/2 z = x/2 (1 + k^2 x^2)^{1/2} + 1/(2k) \sinh^{-1}(kx)$. We have $k = g/(v \cos \theta)^2$, and $a = v^2/g \sin \theta \cos \theta$, so $ka = \tan \theta$. So the required path length is $p(\theta) = a(1 + k^2 a^2)^{1/2} + 1/k \sinh^{-1} a = v^2/g (\sin \theta + \cos^2 \theta \sinh^{-1} \tan \theta)$.

To find the maximum, we differentiate, getting $p'(\theta) = \cos \theta - 2 \cos \theta \sin \theta \sinh^{-1} \tan \theta + \cos^2 \theta (1 + \tan^2 \theta)^{-1/2} \sec^2 \theta = 2 \cos \theta (1 - \sin \theta \sinh^{-1} \tan \theta)$. In the range $[0, \pi/2]$, $\tan \theta$ is monotone increasing. $\sinh^{-1} z$ is strictly monotone increasing for positive z , so $\sinh^{-1} \tan \theta$ is strictly monotone increasing on $(0, \pi/2)$. Indeed it evidently tends to ∞ as θ tends to $\pi/2$. Hence $(1 - \sin \theta \sinh^{-1} \tan \theta)$ is strictly monotone decreasing on $(0, \pi/2)$ and crosses zero once. $\cos \theta$ is monotone decreasing and positive, so $p'(\theta)$ is strictly monotone decreasing and crosses zero once. Hence $p(\theta)$ has a single maximum on $[0, \pi/2]$, which is achieved for the value φ in $(0, \pi/2)$ for which $(1 - \sin \varphi \sinh^{-1} \tan \varphi) = 0$. Rearranging, $\sinh(1/\sin \varphi) = \tan \varphi$. Squaring, adding 1, and taking the square root: $\cosh(1/\sin \varphi) = \sec \varphi$. Adding the last two equations: $e^{1/\sin \varphi} = \tan \varphi + \sec \varphi$, or $1/\sin \varphi = \ln(\tan \varphi + \sec \varphi)$, or $\sin \varphi \ln(\tan \varphi + \sec \varphi) = 1$.

Problem B2

C_1, C_2 are cylindrical surfaces with radii r_1, r_2 respectively. The axes of the two surfaces intersect at right angles and $r_1 > r_2$. Let S be the area of C_1 which is enclosed within C_2 . Prove that $S = 8r_2^2 A = 8r_1^2 C - 8(r_1^2 - r_2^2)B$, where $A = \int_0^1 (1 - x^2)^{1/2} (1 - k^2 x^2)^{-1/2} dx$, $B = \int_0^1 (1 - x^2)^{-1/2} (1 - k^2 x^2)^{-1/2} dx$, and $C = \int_0^1 (1 - x^2)^{-1/2} (1 - k^2 x^2)^{1/2} dx$, and $k = r_2/r_1$.

Solution

It is hard to see the point of the second half $[8r_2^2 A = 8r_1^2 C - 8(r_1^2 - r_2^2)B]$, which is trivial. Multiply top and bottom of the integrand in A by $(1 - x^2)^{1/2}$, so that we get $(1 - x^2)$ on the top. Then note that $k^2(1 - x^2) = 1 - k^2 x^2 + (k^2 - 1)$.

For the first half, the part of C_1 enclosed inside C_2 comprises two bent ovals, one at each end. It is tempting to think that when rolled flat, each piece is an ellipse. If that were true, then the problem would be trivial - the semi-axes are r_2 and $r_1 \sin^{-1} k$, so the total area would be $2\pi r_1 r_2 \sin^{-1} k$. But conics only remain conics when projected onto flat surfaces and here we are projecting onto a curved surface.

So we have to use integration. Take C_1 to be $x^2 + y^2 = r_1^2$, and C_2 to be $y^2 + z^2 = r_2^2$. We calculate the area of the quarter of the piece with $x > 0$ having $y, z > 0$. We may divide it into strips parallel to the z -axis. Take the angle θ to be the axial angle - the angle by which the strip has to be rotated about the z -axis from the $y = 0$ position (keeping in the surface C_1). [Points on the strip have the same x and y coordinates, but varying z -coordinates.] The strip has y -coordinate $r_1 \sin \theta$ and hence length $(r_2^2 - r_1^2 \sin^2 \theta)^{1/2}$ and width $r_1 d\theta$. Thus the area of the quarter-oval is $\int_0^{\sin^{-1} k} (r_2^2 - r_1^2 \sin^2 \theta)^{1/2} r_1 d\theta$. Put $t = 1/k \sin \theta$, and the integral becomes $r_2^2 \int_0^1 (1 - t^2)^{1/2} (1 - k^2 t^2)^{-1/2} dt$. Hence the complete oval is 4 times this, and the total area 8 times.

Comments. The original question was badly worded, because at first sight it appeared to be about volumes, not areas, which may have confused some. It is important to get a clear picture of the geometry - the easiest way to do this is to roll a piece of paper into a cylinder.

Problem B3

Let p be a positive real, let S be the parabola $y^2 = 4px$, and let P be a point with coordinates (a, b) . Show that there are 1, 2 or 3 normals from P to S according as $4(2p - a)^2 + 27pb^2 >, =$ or < 0 .

Solution

The general point on the parabola is $(pt^2, 2pt)$. The slope of the tangent is $2p/y = 1/t$, and so the slope of the normal is $-t$. Hence the equation of the normal is $(y - 2pt) = -t(x - pt^2)$. This passes through (a, b) iff $pt^3 + (2p - a)t - b = 0$ (*).

This is a cubic, so it has 1, 2 or 3 real roots. We have to decide which. If $(2p - a) > 0$, then $pt^3 + (2p - a)t$ is strictly increasing and takes all values in $(-\infty, \infty)$, so (*) has just one real root. The same is true if $(2p - a) = 0$, unless $b = 0$, in which case there are three coincident roots.

If $(2p - a) < 0$, then $pt^3 + (2p - a)t$ has a maximum and a minimum. Differentiating, we find that these are at $t = \pm \sqrt{-(2p - a)/(3p)}$, with values $\pm 2(2p - a)/3 \sqrt{-(2p - a)/(3p)}$. So there are 1, 2 or 3 real roots according as $b^2 >, =$ or $< 4(2p - a)^2/9 - (2p - a)/3p$, which is the condition in the question. Note that if $(2p - a) \geq 0$, then this expression is certainly > 0 , so the same rule applies.

Problem B4

Let S be the surface $ax^2 + by^2 + cz^2 = 1$ (a, b, c all non-zero), and let K be the sphere $x^2 + y^2 + z^2 = 1/a + 1/b + 1/c$ (known as the *director* sphere). Prove that if a point P lies on 3 mutually perpendicular planes, each of which is tangent to S , then P lies on K .

Solution

Let $f(x, y, z) = ax^2 + by^2 + cz^2$. Then the normal vector is the vector $\text{grad } f$, so the tangent plane at (u, v, w) is $a.u.x + b.v.y + c.w.z = 1$.

Note that this does not pass through the origin. The general plane not through the origin has equation $\mathbf{p} \cdot (\mathbf{x} - \lambda \mathbf{p}) = 0$ or $\mathbf{p} \cdot \mathbf{x} = \lambda$, where \mathbf{x} is the vector (x, y, z) representing a general point on the plane, $\mathbf{p} = (p, q, r)$ is a unit vector normal to the plane, and $\lambda > 0$ is the distance of the plane from the origin. If this is a tangent plane at some point of the quadric, then $a(p/\lambda a)^2 + b(q/\lambda b)^2 + c(r/\lambda c)^2 = 1$, or $p^2/a + q^2/b + r^2/c = \lambda^2$.

So suppose P is the point of intersection of three perpendicular tangent planes $p_i x + q_i y + r_i z = \lambda_i$, $i = 1, 2, 3$, where \mathbf{p}_i are orthonormal vectors. The squared distance of P from the origin is $\lambda_1^2 + \lambda_2^2 + \lambda_3^2 = (p_1^2 + p_2^2 + p_3^2)/a + (q_1^2 + q_2^2 + q_3^2)/b + (r_1^2 + r_2^2 + r_3^2)/c = 1/a + 1/b + 1/c$, since \mathbf{p}_i are orthonormal. [This is the key trick: if the rows of a matrix are orthonormal vectors, then the matrix is orthogonal and hence its *columns* are also orthonormal vectors.].

Problem B5

Find all rational triples (a, b, c) for which a, b, c are the roots of $x^3 + ax^2 + bx + c = 0$.

Solution

Answer: $(0, 0, 0); (1, -1, -1), (1, -2, 0)$.

We require (1) $a + b + c = -a$, (2) $ab + bc + ca = b$, and (3) $abc = -c$.

From (3), either $c = 0$, or $ab = -1$. If $c = 0$, then (1) becomes $b = -2a$, and (2) becomes $b(a - 1) = 0$. Hence either $a = b = 0$, or $a = 1, b = -2$.

So assume $c \neq 0$, and $ab = -1$. (1) becomes $c = -b - 2a$. Substituting in (2), we get: $-1 - (2a + b)(a + b) = b$, so $-a^2 - 2a^4 + 3a^2 - 1 = -a$, or $2a^4 - 2a^2 - a + 1 = 0$. So $a = 1$, or $2a^3 + 2a^2 - 1 = 0$ (*). The first possibility gives $a = 1, b = -1, c = -1$. Suppose $a = m/n$ is a root of (*) with m, n relatively prime integers. Then $2m^3 + 2m^2n - n^3 = 0$. So any prime factor of n must divide 2 and any prime factor of m must divide 1. Hence the only possibilities are $a = 1, -1, 1/2, -1/2$, and we easily check that these are not solutions. So (*) has no rational roots.

Problem B6

The $n \times n$ matrix (m_{ij}) is defined as $m_{ij} = a_i a_j$ for $i \neq j$, and $a_i^2 + k$ for $i = j$. Show that $\det(m_{ij})$ is divisible by k^{n-1} and find its other factor.

Solution

Answer: $\det(m_{ij}) = k^{n-1}(k + \sum a_i^2)$.

Induction on n . Clearly true for $n = 1$.

Expanding by the first row we get $k \cdot k^{n-2}(k + \sum_{i>1} a_i^2) + \det(m'_{ij})$, where m'_{ij} is the same as m_{ij} except that $m'_{11} = a_1^2$. Subtracting appropriate multiples of the first row from the others we zero all the elements outside the first row except those on the diagonal, which become k . Hence $\det(m'_{ij}) = k^{n-2} a_1^2$.

Problem B7

Given $n > 8$, let $a = \sqrt[n]{n}$ and $b = \sqrt[n+1]{n+1}$. Which is greater a^b or b^a ?

Solution

Answer: a^b is greater.

$a^b = e^{b \ln a}$ and $b^a = e^{a \ln b}$. So we have to decide which of $b \ln a$ and $a \ln b$ is greater, or, equivalently, which of $(\ln a)/a$ and $(\ln b)/b$ is greater. The latter is clearly more promising. So set $f(x) = (\ln x)/x$. Then $f'(x) = 1/x^2 - (\ln x)/x^2$ which is negative for $x > e$. Obviously $b > a$, so provided $a > e$, $(\ln a)/a > (\ln b)/b$ and hence $b \ln a > a \ln b$ and $a^b > b^a$. But $e^2 < 9$, so the result is certainly true for $n \geq 9$.

4th Putnam 1941

Problem A1

Prove that $(a-x)^6 - 3a(a-x)^5 + 5/2 a^2(a-x)^4 - 1/2 a^4(a-x)^2 < 0$ for $0 < x < a$.

Solution

Change variables to $t = 1 - x/a$ and the polynomial becomes $a^6(t^6 - 3t^5 + 5/2 t^4 - 1/2 t^2)$. This obviously has a factor t^2 , and almost obviously $(t-1)$. Dividing these out, we see that the resulting cubic has another factor $(t-1)$. So we can write the original as $a^6 t^2(t-1)^2(t(t-1) - 1/2)$, which evidently has the same sign as $t(t-1) - 1/2$. But that is clearly negative for t between 0 and 1.

Problem A2

Define $f(x) = \int_0^x \sum_{i=0}^{n-1} (x-t)^i / i! dt$. Find the n th derivative $f^{(n)}(x)$.

Solution

Note that x appears both in the integrand and in the limits, so a little care is needed. Write $g_r(x, t) = \sum_{i=0}^{r-1} (x-t)^i / i!$ so that $f(x) = \int_0^x g_n(x, t) e^{nt} dt$. By definition $f'(x) = \lim_{\delta x \rightarrow 0} (\int_0^{x+\delta x} g_n(x+\delta x, t) e^{nt} dt - \int_0^x g_n(x, t) e^{nt} dt) / \delta x = e^{nx} g_n(x, x) + \int_0^x g_n'(x, t) e^{nt} dt$, where the ' denotes the partial derivative wrt the first variable. But $g_n(x, x) = 1$, and $g_r'(x, t) = g_{r-1}(x, t)$, so $f'(x) = e^{nx} + \int_0^x g_{n-1}(x, t) e^{nt} dt$. Hence by an easy induction $f^{(n)}(x) = e^{nx} (1 + n + n^2 + \dots + n^{n-1})$.

Problem A3

A circle radius a rolls in the plane along the x -axis the envelope of a diameter is the curve C . Show that we can find a point on the circumference of a circle radius $a/2$, also rolling along the x -axis, which traces out the curve C .

Solution

Consider a circle radius $a/2$ with center initially at $(0, a/2)$ rolling along the x -axis. After rolling through an angle θ , the point initially at $(0, a)$ is at $a/2 \sin \theta, a/2 \cos \theta$ relative to the center and hence at $P(a\theta/2 + (a/2) \sin \theta, a/2 + (a/2) \cos \theta)$. The tangent at P is $(y - (a/2)(1 + \cos \theta)) / (x - (a/2)(\theta + \sin \theta)) = (-\sin \theta) / (1 + \cos \theta)$, or $x \sin \theta + y(1 - \cos \theta) = (a/2)(\theta \sin \theta + 2 \cos \theta + 2)$.

If we put $\theta = 2\phi$, then $\sin \theta = 2 \sin \phi \cos \phi$, $1 + \cos \theta = 2 \cos^2 \phi$, so the tangent at P has equation, $x \sin \phi + y \cos \phi = a(\phi \sin \phi + \cos \phi)$.

Now consider the circle radius a with center initially at $(0, a)$. When it has rolled through an angle ϕ , its center is at $(a\phi, a)$, so the diameter which is initially horizontal lies on the line $(y-a)/(x-a\phi) = -\tan \phi$, or $x \sin \phi + y \cos \phi = a(\phi \sin \phi + \cos \phi)$. In other words, the diameter is tangent to the point P of the curve traced out by the point on the circumference of the circle radius $a/2$. Hence the envelope of the diameter is that curve.

Problem A4

The real polynomial $x^3 + px^2 + qx + r$ has real roots $a \leq b \leq c$. Prove that f' has a root in the interval $[b/2 + c/2, b/3 + 2c/3]$. What can we say about f if the root is at one of the endpoints?

Solution

$p(x) = (x-a)(x-b)(x-c)$, so $p'(x) = (x-a)(x-b) + (x-b)(x-c) + (x-a)(x-c)$.

We can write $p'(x) = (x-b)(x-c) + (x-a)(2x-b-c)$, so $p'(b/2 + c/2) = -1/4 (c-b)^2 \leq 0$, with equality iff $b = c$.

$p'(b/3 + 2c/3) = -2/9 (c-b)^2 + (x-a) 1/3 (c-b) = 1/3 (c-b) (b-a) \geq 0$ with equality iff $a = b$ or $b = c$.

If $b = c$, then b is a repeated root and $p'(b) = 0$. If $a = b$, then $p'(b/3 + 2c/3) = 0$. Otherwise, $p'(x)$ is negative at $b/2 + c/2$ and positive at $b/3 + 2c/3$, so it has a zero in the interior of the interval.

Problem A6

f is defined for the non-negative reals and takes positive real values. The centroid of the area lying under the curve $y = f(x)$ between $x = 0$ and $x = a$ has x -coordinate $g(a)$. Prove that for some positive constant k , $f(x) = k g'(x)/(x - g(x))^2 e^{\int 1/(t - g(t)) dt}$.

Problem A7

Do either (1) or (2):

(1) Do either (1) or (2):

(1) Let A be the 3 x 3 matrix

$$\begin{matrix} 1+x^2-y^2-z^2 & 2(xy+z) & 2(zx-y) \\ 2(xy-z) & 1+y^2-z^2-x^2 & 2(yz+x) \\ 2(zx+y) & 2(yz-x) & 1+z^2-x^2-y^2 \end{matrix}$$

Show that $\det A = (1 + x^2 + y^2 + z^2)^3$.

(2) A solid is formed by rotating about the x-axis the first quadrant of the ellipse $x^2/a^2 + y^2/b^2 = 1$. Prove that this solid can rest in stable equilibrium on its vertex (corresponding to $x = a, y = 0$ on the ellipse) iff $a/b \leq \sqrt{8/5}$.

Solution

(1) subtract z times row 2 from row 1 and add y times row 3 to row 1. After taking out the common factor $1+x^2+y^2+z^2$ from row 1 we get:

$$\begin{matrix} 1 & z & -y \\ 2(xy-z) & 1+y^2-z^2-x^2 & 2(yz+x) \\ 2(zx+y) & 2(yz-x) & 1+z^2-x^2-y^2 \end{matrix}$$

Subtract z times col 1 from col 2 and add y times col 1 to col 3. We get:

$$\begin{matrix} 1 & 0 & 0 \\ 2(xy-z) & 1+y^2+z^2-x^2-2xyz & 2x(1+y^2) \\ 2(zx+y) & -2x(1+z^2) & 1+z^2-x^2+y^2+2xyz \end{matrix}$$

Multiplying this out, we get $(1-x^2+y^2+z^2)^2 - 4x^2y^2z^2 + 4x^2(1+y^2+z^2+y^2z^2) = (1+x^2+y^2+z^2)^2$. Hence with the additional factor we took out, we get the result.

(2) We first have to find the position of the centre of mass on the axis. The moment about the y-axis of the solid is $\int_0^a \pi y^2 x dx = \pi b^2 \int_0^a (x - x^3/a^2) dx = \pi b^2 a^2/4$. The volume is $\int_0^a \pi y^2 dx = \pi b^2 \int_0^a (1 - x^2/a^2) dx = 2/3 \pi b^2 a$. Hence the centre of mass is a distance $3a/8$ from the flat surface or $5a/8$ from the point of contact.

Now suppose the point of contact is at $(a \cos t, b \sin t)$. The tangent has gradient $-b/a \cot t$, so the normal has gradient $a/b \tan t$. So the equation of the normal is $y - b \sin t = a/b \tan t (x - a \cos t)$. This meets the x-axis at $a(1 - b^2/a^2) \cos t$. For stability we want this to be closer to the origin than the centre of mass, in other words we want $(1 - b^2/a^2) \cos t < 3/8$. The point of contact is at $\cos t = 1$, so we require $(1 - b^2/a^2) < 3/8$ or $b/a > \sqrt{5/8}$.

Problem B1

A particle moves in the plane so that its angular velocity about the point $(1, 0)$ equals minus its angular velocity about the point $(-1, 0)$. Show that its trajectory satisfies the differential equation $y' x(x^2 + y^2 - 1) = y(x^2 + y^2 + 1)$. Verify that this has as solutions the rectangular hyperbolae with center at the origin and passing through $(\pm 1, 0)$.

Solution

The angular momentum (for unit mass) about $(1, 0)$ is $(x - 1) dy/dt - y dx/dt$. Hence the angular velocity is $((x - 1) dy/dt - y dx/dt) / ((x - 1)^2 + y^2)$. Similarly, the angular velocity about $(-1, 0)$ is $((x + 1) dy/dt - y dx/dt) / ((x + 1)^2 + y^2)$. Hence the trajectory satisfies: $((x - 1) y' - y)(x^2 + y^2 + 1 + 2x) + ((x + 1) y' - y)(x^2 + y^2 + 1 - 2x) = 0$ or $2x y'(x^2 + y^2 + 1) - 4x y' - 2y(x^2 + y^2 + 1) = 0$, or $x y'(x^2 + y^2 - 1) = y(x^2 + y^2 + 1)$.

We see immediately that $xy = 0$ is a solution and almost immediately that $x^2 - y^2 = 1$ is a solution. Hence also all linear combinations $x^2 + kxy - y^2 = 1$. These are rectangular hyperbolae because the sum of the coefficients of x^2 and y^2 are zero.

Problem B2

Find:

- (1) $\lim_{n \rightarrow \infty} \sum_{1 \leq i \leq n} 1/\sqrt{(n^2 + i^2)}$;
- (2) $\lim_{n \rightarrow \infty} \sum_{1 \leq i \leq n} 1/\sqrt{(n^2 + i)}$;
- (3) $\lim_{n \rightarrow \infty} \sum_{1 \leq i \leq n^2} 1/\sqrt{(n^2 + i)}$;

Solution

(1) $1/\sqrt{(n^2 + i^2)} = (1/n) / \sqrt{(1 + (i/n)^2)}$. So the sum is just a Riemann sum for the integral $\int_0^1 dx/\sqrt{(1 + x^2)} = \sinh^{-1}1 = \ln(1 + \sqrt{2}) = 0.8814$.

(2) $1/\sqrt{(n^2 + i)} = (1/n) / \sqrt{(1 + i/n^2)}$. Each term is less than $1/n$, so the (finite) sum is less than 1. But each term is at least $(1/n) / \sqrt{(1 + 1/n)}$. So the sum is at least $1/\sqrt{(1 + 1/n)}$, which tends to 1. Hence the limit of the sum is 1.

(3) $1/\sqrt{(n^2 + i)} = n(1/n^2) / \sqrt{(1 + i/n^2)}$. Now $\sum (1/n^2) / \sqrt{(1 + i/n^2)}$ is just a Riemann sum for $\int_0^1 dx/\sqrt{(1 + x)} = 2\sqrt{(1 + x)}|_0^1 = 2(\sqrt{2} - 1)$. So the sum given tends to $2(\sqrt{2} - 1)n$, which diverges to infinity. [Or simpler, there are n^2 terms, each at least $1/\sqrt{(2n^2)} = 1/(n\sqrt{2})$, so the sum is at least $n/\sqrt{2}$ which diverges.]

Problem B3

Let y and z be any two linearly independent solutions of the differential equation $y'' + p(x)y' + q(x)y = 0$. Let $w = yz$. Find the differential equation satisfied by w .

Solution

We have $w = yz$, $w' = y'z + yz'$, $w'' = y''z + 2y'z' + yz''$. Hence $w'' + pw' + 2qw = 2y'z'$ (1). Now $(y'z)' = y''z + y'z'$, $2py'z' = py'z' + py'z'$, $qw' = qyz' + qy'z$, so $(y'z)' + 2py'z' + qw' = 0$ (2).

Differentiating (1) we get: $w''' + p'w' + pw'' + 2q'w + 2qw' = 2(y'z)'' = -4py'z' - 2qw'$ (using (2)), $= -2p(w'' + pw' + 2qw) - 2qw'$. Rearranging, this gives: $w''' + 3p'w' + (p' + 2p^2 + 4q)w' + (4pq + 2q')w = 0$.

Problem B4

Given an ellipse center O , take two perpendicular diameters AOB and COD . Take the diameter $A'O'B'$ parallel to the tangents to the ellipse at A and B (this is said to be *conjugate* to the diameter AOB). Similarly, take $C'O'D'$ conjugate to COD . Prove that the rectangular hyperbola through $A'B'C'D'$ passes through the foci of the ellipse.

Solution

Take the ellipse as $x^2/a^2 + y^2/b^2 = 1$ and the point A as $(a \cos t, b \sin t)$. Then AB has slope $b/a \tan t$, so CD has slope $-a/b \cot t$. The tangent at A has slope $-b/a \cot t$. Suppose C is $(a \cos u, b \sin u)$, then $b/a \tan u = -a/b \cot t$ and the tangent at C has slope $-b/a \cot u = b^3/a^3 \tan t$. Hence the line pair $A'B', C'D'$ has equation $(y + x b/a \cot t)(y - x b^3/a^3 \tan t)$. Now we have the equations for two distinct conics through $A'B'C'D'$: the original ellipse and the line pair $A'B', C'D'$. The equation of any other conic through these four points must be a linear combination of the equations of these two, in other words, $(x^2/a^2 + y^2/b^2 - 1) + k(y + x b/a \cot t)(y - x b^3/a^3 \tan t) = 0$ for some k .

The criterion for a rectangular hyperbola is that the coefficients of x^2 and y^2 should have sum zero, or that $1/a^2 + 1/b^2 + k - k b^4/a^4 = 0$. Hence $k = a^2/(b^4 - b^2 a^2)$ and the equation of the rectangular hyperbola is $x^2 - y^2 + (b/a \tan t - a/b \cot t)xy = a^2 - b^2$. But the foci are at $(\pm(a^2 - b^2)^{1/2}, 0)$, so they lie on the rectangular hyperbola.

Problem B5

A wheel radius r is traveling along a road without slipping with angular velocity $\omega > \sqrt{(g/r)}$. A particle is thrown off the rim of the wheel. Show that it can reach a maximum height above the road of $(r\omega + g/\omega)^2/(2g)$. [Ignore air resistance.]

Solution

Suppose the pebble leaves the wheel from a point on the rim which is at an angle θ to the vertical. Its point of departure is a distance $r + r \cos \theta$ above the road. Its upward velocity is $r\omega \sin \theta$, so it ascends a further $(r\omega \sin \theta)^2/2g$. Thus the total height is $r + r^2\omega^2/2g + r \cos \theta - r^2\omega^2/2g \cos^2 \theta = r + g/2\omega^2 + r^2\omega^2/2g - r^2\omega^2/2g (\cos \theta - g/r\omega^2)$

(*). We are given that $g/r\omega^2 < 1$, so (*) has a maximum when $\cos \theta = g/r\omega^2$ and the maximum value is $1/2g (r^2\omega^2 + 2gr + g^2/\omega^2) = (r\omega + g/\omega)^2/2g$.

Problem B6

f is a real valued function on $[0, 1]$, continuous on $(0, 1)$. Prove that $\int_{x=0}^{x=1} \int_{y=x}^{y=1} \int_{z=x}^{z=y} f(x) f(y) f(z) dz dy dx = 1/6 (\int_{x=0}^{x=1} f(x) dx)^3$.

Solution

Let S_{xyz} be the points in the cube for which $x \leq y \leq z$, let S_{yxz} be the points for which $y \leq x \leq z$ and so on. Then the union of the six sets is the cube and the intersection of any two has measure zero. Also by changing the variables of integration we see that the integral of $f(x) f(y) f(z)$ over each set is the same. Hence the integral over S_{xzy} is $1/6$ of the integral over the cube. But the integral over S_{xzy} is just $\int_{x=0}^{x=1} \int_{y=x}^{y=1} \int_{z=x}^{z=y} f(x) f(y) f(z) dz dy dx$ and the integral over the cube is $(\int_{x=0}^{x=1} f(x) dx)^3$.

Problem B7

Do either (1) or (2):

(1) f is a real-valued function defined on the reals with a continuous second derivative and satisfies $f(x + y) f(x - y) = f(x)^2 + f(y)^2 - 1$ for all x, y . Show that for some constant k we have $f''(x) = \pm k^2 f(x)$. Deduce that $f(x)$ is one of $\pm \cos kx, \pm \cosh kx$.

(2) a_i and b_i are constants. Let A be the $(n+1) \times (n+1)$ matrix A_{ij} , defined as follows: $A_{i1} = 1; A_{1j} = x^{j-1}$ for $j \leq n; A_{1(n+1)} = p(x); A_{ij} = a_{i-1}^{j-1}$ for $i > 1, j \leq n; A_{i(n+1)} = b_{i-1}$ for $i > 1$. We use the identity $\det A = 0$ to define the polynomial $p(x)$. Now given any polynomial $f(x)$, replace b_i by $f(b_i)$ and $p(x)$ by $q(x)$, so that $\det A = 0$ now defines a polynomial $q(x)$. Prove that $f(p(x))$ is a multiple of $\prod (x - a_i)$ plus $q(x)$.

Solution

(1) Putting $y = 0$ gives $f(0)^2 = 1$, so $f(0) = \pm 1$. Differentiating wrt y gives $f'(x+y) f(x-y) - f(x+y) f'(x-y) = 2 f(y) f'(y)$. Putting $y = 0$ gives $f'(0) = 0$. Differentiating wrt x gives $f''(x+y) f(x-y) = f(x+y) f''(x-y)$. Putting $x = y = z/2$ gives $f''(z) = h f(z)$, where $h = \pm f''(0)$. If h is positive, we may put $h = k^2$ and integrate, using $f(0) = \pm 1, f'(0) = 0$ to get $f(x) = \pm \cos kx$. If h is negative, we may put $h = -k^2$ and integrate to get $f(x) = \pm \cosh kx$.

(2)

5th Putnam 1942

Problem A1

ABCD is a square side $2a$ with vertices in that order. It rotates in the first quadrant with A remaining on the positive x -axis and B on the positive y -axis. Find the locus of its center.

Solution

Answer: the segment (a, a) to $(a\sqrt{2}, a\sqrt{2})$.

Let AB make an angle θ with the x -axis. Then we find that the coordinates of the center to be $x = y = a \cos \theta + a \sin \theta$. But $a \cos \theta + a \sin \theta = a \sqrt{2} \sin(\theta + \pi/4)$.

Problem A2

a and b are unequal reals. What is the remainder when the polynomial $p(x)$ is divided $(x - a)^2(x - b)$.

Solution

Suppose the remainder is $cx^2 + dx + e$. We have $p(a) = ca^2 + da + e$, $p(b) = cb^2 + db + e$. Also, differentiating, we get $p'(a) = 2ca + d$. Solving, $c = p'(a)/(a - b) - p(a)/(a - b)^2 + p(b)/(a - b)^2$, $d = (2a p(a) - 2a p(b) - (a^2 - b^2)p'(a))/(a - b)^2$, $e = p(a) - a^2(p(a) - p(b))/(a - b)^2 + ab p'(a)/(a - b)$.

Problem A3

Does $\sum_{n \geq 0} n! k^n / (n + 1)^n$ converge or diverge for $k = 19/7$?

Solution

The n th term divided by the $(n-1)$ th term is $k n^{n-1} / (n+1)^n = k / (1 + 1/n)^n$ which tends to k/e . But $k/e < 1$, so the series converges by the ratio test.

Problem A4

Let C be the family of conics $(2y + x)^2 = a(y + x)$. Find C' , the family of conics which are orthogonal to C . At what angle do the curves of the two families meet at the origin?

Solution

For most points P in the plane we can find a unique conic in the family passing through the point. Thus we should be able to find the gradient of members of the family at (x, y) in a formula which is independent of a . We then use this to get a formula for the gradient of the orthogonal family and solve the resulting first-order differential equation to get the orthogonal family.

Thus we have $8y y' + 4x y' + 4y + 2x = ay' + a = (y' + 1)(2y + x)^2 / (y + x)$. So $y'(2y+x)(4(x+y) - (2y+x)) = (2y+x)^2 - (x+2y)(x+y)$, or $y'(2y+x)(2y+3x) = -2x(2y+x)$, so $y' = -x/(2y+3x)$. Hence the orthogonal family satisfies $y' = (2y+3x)/x$. So $y'/x^2 - 2y/x^3 = 3/x^2$. Integrating $y = bx^2 - 3x$. These are all parabolas.

All members of both families pass through the origin. Changing coordinates to $X = x + 2y$, $Y = y - 2x$, the equation of a member of the first family becomes $X^2 = a(3X - Y)/5$ or $Y = -5/a(X - 3a/10)^2 + 9a/20$. This has gradient 3 (in the new system) at the origin. In the old system the tangent is $y = -x$. The orthogonal set obviously has gradient -3 at the origin. If the angle between them is k , then $\tan k = (-1 + 3)/(1 + 3) = 1/2$. So $k = \tan^{-1} 1/2$.

Problem A5


C is a circle radius a whose center lies a distance b from the coplanar line L . C is rotated through π about L to form a solid whose center of gravity lies on its surface. Find b/a .

Answer

$(\pi + \sqrt{(\pi^2 + 2\pi - 4)}) / (2\pi - 4) = \text{about } 2.9028$

Solution

The solid is half a torus. We can divide it into a large number of thin disks. Each disk has variable thickness, with thickness proportional to the distance from L . So we must integrate to find the distance of the centroid of the disk from L . Take the density to be kd , where d is the distance from L .



Take x to be distance along the line perpendicular to x , and θ to be the angle between the radius vector and the x -axis. We have $x = a \cos \theta$, so $dx = -a \sin \theta d\theta$. The mass is $\int_0^\pi 2a \sin \theta (a \sin \theta d\theta) k(b + a \cos \theta) = 2a^2bk \int_0^\pi \sin^2 \theta d\theta + 2a^3k \int_0^\pi \sin^2 \theta \cos \theta d\theta = a^2bk\pi + 0$. So the mass times the centroid distance is $\int_0^\pi 2a^2k \sin^2 \theta (a \cos \theta + b)^2 d\theta = 2a^4k \int_0^\pi \sin^2 \theta \cos^2 \theta d\theta + 4a^3bk \int_0^\pi \sin^2 \theta \cos \theta d\theta + 2a^2b^2k \int_0^\pi \sin^2 \theta d\theta = \frac{1}{2}a^4k \int_0^\pi \sin^2 2\theta d\theta + 0 + a^2b^2k\pi = ka^2\pi(a^2/4 + b^2)$. So the centroid distance is $b + a^2/4b$. Thus we can regard the mass as uniformly spread over a semicircle radius $b + a^2/4b$.

We need another integration to find the distance of the mass of a semicircle radius r from its center. It is $(1/\pi r) \int_0^\pi r^2 \sin \theta d\theta = 2r/\pi$. Thus the cm of the half-torus is a distance $(2/\pi)(b + a^2/4b)$ from L . We want it to be a distance $b-a$ from L so that it lies on the surface. Thus $(2/\pi)(b + a^2/4b) = b - a$, so $(2\pi-4)b^2 - 2\pi ab - a^2 = 0$. Hence $b/a = (\pi + \sqrt{\pi^2 + 2\pi - 4})/(2\pi - 4) = \text{about } 2.9028$.

Problem A6

P is a plane and H is the half-space on one side of P . K is a fixed circle in P . C is a circle in P which cuts K at an angle α . Let C have center O and radius r . $f(C)$ is the point in H on the normal to P through O and a distance r from O . Show that the locus of $f(C)$ is a one-sheet hyperboloid and that it has two families of rulings in it.

Problem B1

S is a solid square side $2a$. It lies in the quadrant $x \geq 0, y \geq 0$, and it is free to move around provided a vertex remains on the x -axis and an adjacent vertex on the y -axis. P is a point of S . Show that the locus of P is part of a conic. For what P does the locus degenerate?

Solution

Let A be the vertex that moves along the x -axis and B the vertex that moves along the y -axis. Suppose that when AB is horizontal P has coordinates b, c . In the general configuration let be the angle BAO be θ . Then P has coordinates $x = (2a - b) \cos \theta + c \sin \theta, y = b \sin \theta + c \cos \theta$. Hence $cx - (2a - b)y = (b^2 + c^2 - 2ab) \sin \theta, bx - cy = (2ab - b^2 - c^2) \cos \theta$. Squaring and adding we eliminate θ to get: $(b^2 + c^2) x^2 - 4ac xy + (4a^2 + b^2 + c^2 - 4ab) y^2 = (b^2 + c^2 - 2ab)^2$, which is the equation of a conic. So the locus of P must form part of this conic.


The conic degenerates if $b^2 + c^2 = 2ab$. In this case, the equation becomes $2ab x^2 - 4ac xy + (4a^2 - 2ab)y^2 = 0$, or $bx^2 - 2c xy + (2a - b)y^2 = 0$, or $b^2x^2 - 2bc xy + c^2 y^2 = 0$, or $bx = cy$. So in this case the locus lies on a straight line. We may write the condition $b^2 + c^2 = 2ab$ as $(a - b)^2 + c^2 = a^2$, which shows that such P lie on the semicircle diameter AB .

Problem B2

Let P_a be the parabola $y = a^3x^2/3 + a^2x/2 - 2a$. Find the locus of the vertices of P_a , and the envelope of P_a . Sketch the envelope and two P_a .

Solution

We can write the equation of P_a as $(y + 35a/16) = (a^3/3)(x + 3/4a)^2$, so the vertex is $x = -3/4a, y = -35a/16$. The locus of the vertex is $xy = 105/64$.



The graph shows $P_3, P_2, P_1, P_{1/2}, P_{1/3}$ and the two hyperbolae $yx = -7/6, yx = 10/3$. It shows that for positive a , the parabolas touch the lower branches of the hyperbolae. For negative a they touch the upper branches.

That is not hard to verify. We claim that P_a and $xy = 10/3$ touch at $x = -2/a, y = -5a/3$. The point obviously lies on $xy = 10/3$. We have $(a^3/3)(x + 3/4a)^2 = (1/3)(-2 + 3/4)^2 a = 25a/48 = (-5a/3 + 35a/16)$, so the point also lies on P_a . The gradient of $xy = 10/3$ at the point is $-10/(3x^2) = -5a^2/6$. The gradient of P_a at the point is $2a^3x/3 + a^2/2 = -5a^2/6$.

Similarly, we claim that P_a and $xy = -7/6$ touch at $x = 1/a, y = -7a/6$. The point obviously lies on $xy = -7/6$. We have $(a^3/3)(x + 3/4a)^2 = (a/3)(1 + 3/4)^2 = 49a/48 = (-7a/6 + 35a/16)$, so the point also lies on P_a . The gradient of $xy = -7/6$ at the point is $7/(6x^2) = 7a^2/6$. The gradient of P_a at the point is $2a^3x/3 + a^2/2 = a^2(2/3 + 1/2) = 7a^2/6$.

It is less clear how you get the hyperbolae. One standard approach is to look for the singular points of the mapping $f(a,t) = (t, a^3t^2/3 + a^2t/2 - 2a)$. The matrix for the derivative is:

$$\begin{pmatrix} 1 & 0 \\ 2a^3t/3 + a^2/2 & a^2t^2 + at - 2 \end{pmatrix}$$

which has zero determinant when $at = 1$ or -2 , so $xy = 7/6$ or $-10/3$.

Problem B3

$f(x, y)$ and $g(x, y)$ satisfy the differential equation $f_1(x, y) g_2(x, y) - f_2(x, y) g_1(x, y) = 1$ (*). Taking $r = f(x, y)$ and y as independent variables, and $x = h(r, y), g(x, y) = k(r, y)$, show that $k_2(r, y) = h_1(r, y)$. Integrate and hence obtain a solution to (*). What other solutions does (*) have?

Problem B4

A particle moves in a circle through the origin under the influence of a force a/r^k towards the origin (where r is its distance from the origin). Find k .

Solution


The equations of motion are $r(\theta')^2 - r'' = a/r^k, r^2\theta'' = A$ (conservation of angular momentum).

If the particle moves in a circle as described, then we can write its orbit as $r = B \cos \theta$. Differentiating, $r' = -B \theta' \sin \theta = -AB/r^2 \sin \theta$. Differentiating again, $r'' = -AB \cos \theta A/r^2 + 2AB/r^3 \sin \theta r' = -A^2/r^3 - 2A^2B^2/r^5 \sin^2 \theta = -A^2/r^3 - 2A^2B^2/r^5 (1 - r^2/B^2) = A^2/r^3 - 2A^2B^2/r^5$. So substituting back in the equation of motion we get: $2A^2B^2/r^5 = a/r^k$. Hence $k = 5$.

Note that this is unphysical, since we require infinite velocity as we reach the origin.

Problem B5

Let $f(x) = x/(1 + x^6 \sin^2 x)$. Sketch the curve $y = f(x)$ and show that $\int_0^\infty f(x) dx$ exists.



Solution

Obviously $f(x)$ is positive for positive x . But it has an infinite number of spikes at $x = n\pi$. The spike at $n\pi$ has height $n\pi$, so we have to show that the integral is bounded above.

We have $\sin x > 1/2x$ near $x = 0$ (certainly for $x < \pi/3$). So $|\sin x| > 1/(n\pi)^k$ except possibly for $|x| < 2/(n\pi)^k$. Let I_n be the interval centered on $n\pi$ width $4/(n\pi)^k$. For $x \in I_n$ we have $f(x) < 2n\pi$, so the integral of $f(x)$ over the interval is less than $8/(n\pi)^{k-1}$. The total integral over all such intervals is bounded provided that $k > 2$. Outside such

intervals, $x^6 \sin^2 x > 1/2 x^6/x^{2k}$, so $f(x) < 2/x^{2k-5}$. Hence the interval of $f(x)$ over 0 to ∞ excluding the intervals I_n is bounded provided $k > 5/2$. By taking $k = 2 1/4$, for example, we get that the whole integral is bounded.

6th Putnam 1946

Problem A1

$p(x)$ is a real polynomial of degree less than 3 and satisfies $|p(x)| \leq 1$ for $x \in [-1, 1]$. Show that $|p'(x)| \leq 4$ for $x \in [-1, 1]$.

Solution

Let $p(x) = ax^2 + bx + c$, so $p'(x) = 2ax + b$. It is evidently sufficient to show that $|2a + b|$ and $|2a - b| \leq 4$. $p(0) = c$, $p(1) = a + b + c$, $p(-1) = a - b + c$, so $2a + b = 3/2 p(1) + 1/2 p(-1) + 2 p(0)$. But $|p(1)|, |p(-1)|, |p(0)| \leq 1$, so $|2a + b| \leq 4$. Similarly, $2a - b = 1/2 p(1) + 3/2 p(-1) + 2 p(0)$.

Problem A2

\mathbb{R} is the reals. For functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$ and $x \in \mathbb{R}$ define $I(fg) = \int_1^x f(t) g(t) dt$. If $a(x), b(x), c(x), d(x)$ are real polynomials, show that $I(ac) I(bd) - I(ad) I(bc)$ is divisible by $(x - 1)^4$.

Solution

Let $F(x) = I(ac) I(bd) - I(ad) I(bc)$. Then clearly $F(1) = 0$ (since all the integrals are over an empty range). Differentiating, we get, $F' = acI(bd) + bdI(ac) - adI(bc) - bcI(ad)$. So $F'(1) = 0$. Differentiating again: $F'' = a'cI(bd) + ac'I(bd) + abcd + b'dI(ac) + bd'I(ac) + abcd - a'dI(bc) - ad'I(bc) - abcd - b'cI(ad) - bc'I(ad) - abcd = a'cI(bd) + ac'I(bd) + b'dI(ac) + bd'I(ac) - a'dI(bc) - ad'I(bc) - b'cI(ad) - bc'I(ad)$. So $F''(1) = 1$.

Differentiating again: $F''' = a''cI(bd) + 2a'c'I(bd) + ac''I(bd) + a'bcd + abc'd + b''dI(ac) + 2b'd'I(ac) + bd''I(ac) + ab'cd + abcd' - a''dI(bc) - 2a'd'I(bc) - ad''I(bc) - a'b'cd - abcd' - b''cI(ad) - 2b'c'I(ad) - bc''I(ad) - ab'cd - abc'd = a''cI(bd) + 2a'c'I(bd) + ac''I(bd) + b''dI(ac) + 2b'd'I(ac) + bd''I(ac) - a''dI(bc) - 2a'd'I(bc) - ad''I(bc) - b''cI(ad) - 2b'c'I(ad) - bc''I(ad)$. So $F'''(1) = 1$.

That is sufficient to prove the result. But notice that if we differentiate again, then just collecting the terms that do not involve $I(fg)$ we get (after some cancellation) $2a'b'c'd + 2ab'c'd' - 2a'b'c'd' - 2ab'c'd$, which is not, in general, zero for $x = 1$. So in general we do not have $F(x)$ divisible by $(x - 1)^5$.

Problem A3

ABCD are the vertices of a square with A opposite C and side $AB = s$. The distances of a point P in space from A, B, C, D are a, b, c, d respectively. Show that $a^2 + c^2 = b^2 + d^2$, and that the perpendicular distance k of P from the plane ABCD is given by $8k^2 = 2(a^2 + b^2 + c^2 + d^2) - 4s^2 - (a^4 + b^4 + c^4 + d^4 - 2a^2c^2 - 2b^2d^2)/s^2$.

Solution

Let Q be the point of the plane ABCD closest to P. Let O be the center of the square ABCD. Let QO make an angle θ with AC. Then using the cosine rule we have: $AQ^2 = AO^2 + OQ^2 - 2AO \cdot OQ \cos \theta$, $CQ^2 = CO^2 + OQ^2 + 2CO \cdot OQ \cos \theta$ (*). Adding: $AQ^2 + CQ^2 = 2AO^2 + 2OQ^2$. But $a^2 = AQ^2 + k^2$ etc, so $a^2 + c^2 = 2k^2 + s^2 + 2QO^2$. Similarly, $b^2 + d^2 = 2k^2 + s^2 + 2QO^2$. We have established that $a^2 + c^2 = b^2 + d^2$.

We have also established that $8k^2 = 2(a^2 + b^2 + c^2 + d^2) - 4s^2 - 8QO^2$ (**). The angle ϕ between QO and BD is $\pi/2 - \theta$. So $\cos \phi = \sin \theta$. Hence, going back to (*), $AQ^2 - CQ^2 = -4 AO \cdot OQ \cos \theta$, and $BQ^2 - DQ^2 = \pm 4 AO \cdot OQ \sin \theta$. But $AQ^2 - CQ^2 = a^2 - c^2$ etc. So $(a^2 - c^2)^2 + (b^2 - d^2)^2 = 16 AO^2 OQ^2 = 8s^2 OQ^2$. Substituting in (**) gives the required result.

Problem A4

\mathbb{R} is the reals. $f : \mathbb{R} \rightarrow \mathbb{R}$ has a continuous derivative, $f(0) = 0$, and $|f'(x)| \leq |f(x)|$ for all x . Show that f is constant.

Solution

Suppose f is not constant. Then take an interval $[a, b]$ of length $< 1/2$ such that $f(a) = 0$, $f(b) \neq 0$, and $|f(b)| \geq |f(x)|$ for $x \in [a, b]$. Now applying the mean value theorem to the interval gives an immediate contradiction.

Problem A5

Let T be a tangent plane to the ellipsoid $x^2/a^2 + y^2/b^2 + z^2/c^2 = 1$. What is the smallest possible volume for the tetrahedral volume bounded by T and the planes $x = 0, y = 0, z = 0$?

Solution

Answer: $\sqrt{3} abc/2$.

The normal at (x_0, y_0, z_0) is $(xx_0/a^2, yy_0/b^2, zz_0/c^2)$. So the tangent plane is $x x_0/a^2 + y y_0/b^2 + z z_0/c^2 = 1$. This cuts the three axes at $x = a^2/x_0, y = b^2/y_0, z = c^2/z_0$. We can regard two of these lengths as defining the base of the tetrahedron, and the third as forming its height. Hence its volume is $a^2 b^2 c^2 / (6x_0 y_0 z_0)$.

We wish to maximize $x_0 y_0 z_0$. That is equivalent to maximising $x_0^2/a^2 y_0^2/b^2 z_0^2/c^2$. But we know that the sum of these three numbers is 1, so their maximum product is $1/27$ (achieved when they are all equal - the arithmetic/geometric mean result). Hence $x_0 y_0 z_0$ has maximum value $abc/(3\sqrt{3})$.

Problem A6

A particle moves in one dimension. Its distance x from the origin at time t is $a + bt^2 + ct^3$. Find an expression for the particle's acceleration in terms of a, b, c and its speed v .

Solution

Differentiating, $v = 3ct^2 + 2bt + a$ (*), and hence the acceleration $f = 6ct + 2b$. So $t = (f - 2b)/6c$. Substituting in (*) gives $v = (f - 2b)^2/12c + b(f - 2b)/3c + a$. Hence $12cv = f^2 - 4b^2 + 12ac$. So $f = 2\sqrt{(b^2 + 3cv - 3ac)}$.

Problem B1

Two circles C_1 and C_2 intersect at A and B . C_1 has radius 1. L denotes the arc AB of C_2 which lies inside C_1 . L divides C_1 into two parts of equal area. Show L has length > 2 .

Solution

Let O_1 be the center of C_1 , and O_2 the center of C_2 . Let the line $O_1 O_2$ meet the arc AB of C_2 at P . If P lies between O_1 and O_2 , then the tangent to C_2 at P divides C_1 into two unequal parts and the area $C_1 \cap C_2$ lies inside the smaller part. Contradiction. So O_1 must lie between P and O_2 . But now the arc AP is greater than the segment AP , which is greater than $AO_1 = 1$. Hence $L > 2$.

Problem B2

P_0 is the parabola $y^2 = mx$, vertex $K(0, 0)$. If A and B points on P_0 whose tangents are at right angles, let C be the centroid of the triangle ABK . Show that the locus of C is a parabola P_1 . Repeat the process to define P_n . Find the equation of P_n .

Solution

The gradient at the point (x_1, y_1) is $1/2 m/y_1$. So the tangents at $(x_1, y_1), (x_2, y_2)$ are perpendicular iff $y_1 y_2 = -m^2/4$. So we may take one point as $(x, y) = (t^2/m, t)$ and the other as $(1/16 m^3/t^2, -1/4 m^2/t)$. Hence the centroid is (x, y) , where $x = 1/3 (t^2/m + 1/16 m^3/t^2), y = 1/3 (t - 1/4 m^2/t)$. But this lies on the parabola $y^2 = m/3 (x - m/6)$. But since any value of t was possible and hence any value of y , any point of this parabola is a centroid from some pair of points A, B .

Repeating, we get that the equation for P_n is $y^2 = m/3^n (x - m/6(1 + 1/3 + \dots + 1/3^{n-1})) = m/3^n (x - m/4 (1 - 1/3^n))$.

Problem B3

The density of a solid sphere depends solely on the radial distance. The gravitational force at any point inside the sphere, a distance r from the center, is kr^2 (where k is a constant). Find the density (in terms of G, k and r), and the gravitational force at a point outside the sphere. [You may assume the usual results about the gravitational attraction of a spherical shell.]

Solution

A shell exerts no net attraction on a point inside and acts on a point outside as if all its mass was concentrated at its center. So let the density at a radial distance r be $\rho(r)$. Then $G/r^2 \int_0^r 4\pi t^2 \rho(t) dt = kr^2$. Hence $\int_0^r t^2 \rho(t) dt = k/(4G\pi) r^4$. Differentiating: $r^2 \rho(r) = k/(G\pi) r^3$. So $\rho(r) = k/(G\pi) r$.

The force at the surface of the sphere (R) is kR^2 . Hence the force at a distance $r > R$ from the center of the sphere is kR^4/r^2 . [We know the force is a simple inverse square law outside the sphere and it must be continuous at the surface.]

Problem B4

Define $a_n = 2(1 + 1/n)^{2n+1} / ((1 + 1/n)^n + (1 + 1/n)^{n+1})$. Prove that a_n is strictly monotonic increasing.

Solution

$a_n = 2(n+1)^{n+1} / (n^n(2n+1))$. Let $f(n) = \ln a_n / 2 = (n+1) \ln(n+1) - n \ln n - \ln(2n+1)$. Regard f as a function of a real variable and differentiate. $f'(x) = \ln(x+1) - \ln x - 2/(2x+1)$. Differentiate again: $f''(x) = 1/(x+1) - 1/x + 4/(2x+1)^2 = -1/(x(x+1)(2x+1)^2)$.

So $f''(x) < 0$ for all $x > 0$. Hence $f'(x)$ is decreasing. But we can write $f'(x) = \ln(1 + 1/x) - 2/(2x + 1)$ which tends to 0 as x tends to infinity. So $f'(x) > 0$ for all $x > 0$. Hence $f(x)$ is strictly increasing for all positive x . Hence $f(n+1) > f(n)$ and so $\exp f(n+1) > \exp f(n)$, which is the result we want.

Problem B5

Let m be the smallest integer greater than $(\sqrt{3} + 1)^{2n}$. Show that m is divisible by 2^{n+1} .

Solution

This is a fairly well-known problem. $(\sqrt{3} - 1) < 1$ and (we will show) $(\sqrt{3} + 1)^{2n} + (\sqrt{3} - 1)^{2n}$ (*) is an integer. Hence it must be the required integer.

But it is easy to check that (*) is the solution of a recurrence relation, in fact the relation $u_n = 8u_{n-1} - 4u_{n-2}$, $u_0 = 2$, $u_1 = 8$. That establishes that (*) is an integer, and a trivial induction shows that (*) is divisible by 2^{n+1} .

Problem B6

The particle P moves in the plane. At $t = 0$ it starts from the point A with velocity zero. It is next at rest at $t = T$, when its position is the point B . Its path from A to B is the arc of a circle center O . Prove that its acceleration at each point in the time interval $[0, T]$ is non-zero, and that at some point in the interval its acceleration is directly towards the center O .

Solution

This is almost trivial. Take polar coordinates with origin O . Let the radius of the circular arc be k . The particle's radial acceleration is $k(d\theta/dt)^2$ towards O . Its tangential acceleration is $k d^2\theta/dt^2$. Since the particle is not at rest between A and B , $(d\theta/dt)$ is non-zero, so the radial acceleration is non-zero (and hence its total acceleration).

$(d\theta/dt)$ is zero at A and at B , so its derivative (and hence the tangential acceleration) must be zero at some point between.

7th Putnam 1947

Problem A1

The sequence a_n of real numbers satisfies $a_{n+1} = 1/(2 - a_n)$. Show that $\lim_{n \rightarrow \infty} a_n = 1$.

Solution

This is slightly messy. First, since $k = 1/(2 - k)$ implies $k = 1$, it is obvious that if the sequence tends to a limit, then the limit is 1.

Next, if $0 < a_n < 1$, then $1 < 2 - a_n < 2$, so $1/2 < a_{n+1} < 1$. So once the sequence gets into the interval $(0, 1)$ it stays there. But $a_{n+1} - a_n = (a_n - 1)^2 / (2 - a_n) > 0$ for $a_n < 2$ (*). So once the sequence gets into the interval $(0, 1)$ it is monotonic increasing and bounded above by 1, and hence tends to a limit (which must be 1).

If $a_n < 0$, then $0 < a_{n+1} < 1$, so we are also home if a member of the sequence is negative. Similarly, $a_n > 2$ implies $a_{n+1} < 0$. If $a_n = 1$, then all following terms are 1 and so the limit is 1. So the only issue is if $1 < a_1 < 2$.

But then (*) shows that whilst the sequence remains in the interval $(1, 2)$ it is monotonic increasing. It cannot tend to a limit, because that limit would have to be 1, which is impossible. So it cannot stay in the interval. We cannot have $a_m = 2$, because then we would not have $a_{m+1} = 1/(2 - a_m)$, so we must have $a_m > 2$ for some m and then we are home.

Problem A2

\mathbb{R} is the reals. $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous and satisfies $f(r) = f(x) f(y)$ for all x, y , where $r = \sqrt{x^2 + y^2}$. Show that $f(x) = f(1)$ to the power of x^2 .

Solution

Induction on n shows that $f(x \sqrt{n}) = f(x)^n$, and hence $f(n x) = f(x)$ to the power of n^2 . In particular, taking $x = 1/n$, $f(1) = f(1/n)$ to the power of n^2 . Hence, provided $f(1)$ is non-zero, $f(1/n) = f(1)$ to the power of $1/n^2$. Hence $f(m/n) = f(1)$ to the power of $(m/n)^2$. So we have established that $f(x) = f(1)$ to the power of x^2 for all rational x . But f is continuous, so the relation holds for all x .

If $f(1) = 0$, then the same reasoning establishes that $f(x) = 0$ for all x .

Problem A3

ABC is a triangle and P an interior point. Show that we cannot find a piecewise linear path $K = K_1 K_2 \dots K_n$ (where each $K_i K_{i+1}$ is a straight line segment) such that: (1) none of the K_i do not lie on any of the lines AB, BC, CA, AP, BP, CP ; (2) none of the points A, B, C, P lie on K ; (3) K crosses each of AB, BC, CA, AP, BP, CP just once; (4) K does not cross itself.

Solution

Each time K crosses the boundary of a triangle it moves from the outside to the inside or vice versa. K has two endpoints, so we can find one of the three triangles ABP, BCP, CAP in which it does not start or finish. But that is impossible - on the first crossing it must go from outside to inside, on the second from inside to outside and on the third from outside to inside.

Problem A4

Take the x -axis as horizontal and the y -axis as vertical. A gun at the origin can fire at any angle into the first quadrant ($x, y \geq 0$) with a fixed muzzle velocity v . Assuming the only force on the pellet after firing is gravity (acceleration g), which points in the first quadrant can the gun hit?

Solution

Let the angle of the gun to the x -axis be θ . Then the equations of motion are: $x = t v \cos \theta$, $y = t v \sin \theta - 1/2 g t^2$. So the pellet moves along the parabola $y = x \tan \theta - x^2 g / 2v^2 \sec^2 \theta$ (*).

We can view (*) as an equation for θ given x, y . Put $k = \tan \theta$, then the equation becomes $g/2v^2 x^2 k^2 - x k + y + gx^2/2v^2$. This has real roots iff $x^2 \geq 2g/v^2 x^2 y - g^2/v^4 x^4$ and hence iff $y \leq 2g^2/v^2 - x^2/v^4 x^2$ (**). Since $x \geq 0$, we see directly from the quadratic that the sum and the product of the roots are both non-negative, so (**) is the condition for the equation to have at least one non-negative root in k and hence at least one root for θ in the range 0 to $\pi/2$. Thus the gun can hit points in the first quadrant under (or on) the parabola given by (**).

Problem A5

The sequences a_n, b_n, c_n of positive reals satisfy: (1) $a_1 + b_1 + c_1 = 1$; (2) $a_{n+1} = a_n^2 + 2b_n c_n, b_{n+1} = b_n^2 + 2c_n a_n, c_{n+1} = c_n^2 + 2a_n b_n$. Show that each of the sequences converges and find their limits.

Solution

$a_{n+1} + b_{n+1} + c_{n+1} = (a_n + b_n + c_n)^2$, so by a trivial induction $a_n + b_n + c_n = 1$. There appears to be symmetry between the three sequences, so we conjecture that each converges to $1/3$.

Suppose $a_n \leq b_n \leq c_n$. We have $a_{n+1} = a_n^2 + 2b_n c_n \leq a_n c_n + b_n c_n + c_n c_n = c_n(a_n + b_n + c_n) = c_n$. Similarly, $b_{n+1} = b_n^2 + 2a_n c_n \leq b_n c_n + a_n c_n + c_n^2 = c_n$, and $c_{n+1} = c_n^2 + 2a_n b_n \leq c_n^2 + a_n c_n + b_n c_n = c_n$. Hence the largest of $a_{n+1}, b_{n+1}, c_{n+1}$ is no bigger than the largest of a_n, b_n, c_n . An exactly similar argument works for the smallest. Hence the largest forms a monotonic decreasing sequence which is bounded below and the smallest forms a monotonic increasing sequence which is bounded above.

Let $b_n - a_n = h \geq 0, c_n - b_n = k \geq 0$. Then $a_{n+1} - b_{n+1} = (a_n - b_n)(a_n + b_n - 2c_n)$, so $|a_{n+1} - b_{n+1}| = h(h + 2k) \leq (h + k)^2$. Similarly, $|b_{n+1} - c_{n+1}| = |b_n - c_n| |b_n + c_n - 2a_n| = k(2h + k) \leq (h + k)^2$, and $|c_{n+1} - a_{n+1}| = |c_n - a_n| |c_n + a_n - 2b_n| = (h + k)(k - h) \leq (h + k)^2$. So for $n+1$ the difference between the biggest and the smallest is the square of the difference for n . But a_1, b_1, c_1 are all positive and hence, by a trivial induction, a_n, b_n, c_n are positive. Their sum is 1 so the difference between the biggest and smallest must be less than 1. Hence the difference tends to zero. Hence a_n, b_n, c_n all tend to $1/3$.

Problem A6

A is the matrix

a b c

d e f

g h i

$\det A = 0$ and the cofactor of each element is its square (for example the cofactor of b is $fg - di = b^2$). Show that all elements of A are zero.

Solution

$a^2 e^2 - b^2 d^2 = (ei - fh)(ai - cg) - (fg - di)(ch - bi) = (ae - bd) i^2 + (cd - af) hi + (bf - ce) gi = (g^3 + h^3 + i^3) i = 0$, since $0 = \det A = g^3 + h^3 + i^3$. Hence $ae = \pm bd$. Similarly $cd = \pm af, bf = \pm ce$. Multiplying the three equations together we get $abcdef = -abcdef$ unless at least one of the equations has a plus sign. In the first case, at least one of a, b, c, d, e, f is zero. In the second case, the element corresponding to the cofactor is zero - for example $ae = bd$ implies $i^2 = 0$ and hence $i = 0$. So either a member of the first two rows is zero, or a member of the last row is zero.

wlog we may assume $a = 0$. That implies b or $d = 0$ also. [Note that if, for example, i was the zero element, then we would have $ei = \pm fh$, by an argument similar to that above and hence f or $h = 0$]. If $b = 0$, then since $a^3 + b^3 + c^3 = 0$, we have also $c = 0$. Similarly, if $d = 0$, then $g = 0$. So we now have a complete row or column zero. But now the square of any other element is a linear combination of elements in the that row or column and hence zero. Suppose, for example, $a = b = c = 0$. Then $g^2 = bf - ce = 0$, and similarly for the other five elements.

Problem B1

Let \mathbb{R} be the reals. $f : [1, \infty) \rightarrow \mathbb{R}$ is differentiable and satisfies $f'(x) = 1/(x^2 + f(x)^2)$ and $f(1) = 1$. Show that as $x \rightarrow \infty, f(x)$ tends to a limit which is less than $1 + \pi/4$.

Solution

Clearly $f'(x)$ is always positive. But $f(1) = 1$, so $f(x) > 1$ for all x . Hence $f'(x) < 1/(x^2 + 1)$ for all x . Hence $f(x) = 1 + \int_1^x f'(t) dt < 1 + 1 + \int_1^x 1/(1 + t^2) dt = 1 + (\tan^{-1} t)|_1^x = 1 + \tan^{-1} x - \pi/4 < 1 + \pi/2 - \pi/4 = 1 + \pi/4$. Since $f'(x)$ is positive, $f(x)$ is monotone increasing. It is bounded above by $1 + \pi/4$, so it must tend to a limit less than $1 + \pi/4$.

Problem B2

\mathbb{R} is the reals. $f : (0, 1) \rightarrow \mathbb{R}$ is differentiable and has a bounded derivative: $|f'(x)| \leq k$. Prove that: $|\int_0^1 f(x) dx - \sum_{i=1}^n f(i/n)/n| \leq k/n$.

Solution

The worst case for the difference between $1/n \sum_{i=1}^n f(i/n)$ and $\int_{i/n-1/n}^{i/n} f(x) dx$ is if $f'(x) = k$ (or $-k$) for the entire range, in which case the difference is the area of a triangle base $1/n$ and height k/n . Hence the difference for the complete Riemann sum is at worst $k/(2n)$.

Problem B3

Let O be the origin $(0, 0)$ and C the line segment $\{(x, y) : x \in [1, 3], y = 1\}$. Let K be the curve $\{P : \text{for some } Q \in C, P \text{ lies on } OQ \text{ and } PQ = 0.01\}$. Let k be the length of the curve K . Is k greater or less than 2?

Solution

Answer: less.

If we use polar coordinates, then $r = \operatorname{cosec} \theta - .01$, so the length is $\int_{\pi/4}^{\arctan(1/3)} \sqrt{(\operatorname{cosec} \theta - .01)^2 + \operatorname{cosec}^2 \theta \cot^2 \theta} d\theta$. This is obviously horrendous.

The trick is that if we just remove the $.01$, then the integral gives the curve length for the line segment C , which is 2. But the presence of the $-.01$ obviously *reduces* the integrand at every point of the range, so the integral above must have value *less* than 2.

Problem B4

$p(z) \equiv z^2 + az + b$ has complex coefficients. $|p(z)| = 1$ on the unit circle $|z| = 1$. Show that $a = b = 0$.

Solution

In particular, $|p(1)| = |p(-1)| = 1$, so $1 + a + b$ and $1 - a + b$ lie on the unit circle. Hence their midpoint $1 + b$ lies in the unit disk. Similarly, $|p(i)| = |p(-i)| = 1$, so $-1 + ia + b$ and $-1 - ia + b$ lie on the unit circle and hence their midpoint $-1 + b$ lies in the unit disk. But $1 + b$ and $-1 + b$ are a distance 2 apart, so they must lie at either end of a diameter of the unit circle and hence $b = 0$. Now $1 + a$ and $1 - a$ lie on the unit circle, as does their midpoint 1. Hence they must coincide and so $a = 0$.

Problem B5

Let $p(x)$ be the polynomial $(x - a)(x - b)(x - c)(x - d)$. Assume $p(x) = 0$ has four distinct integral roots and that $p(x) = 4$ has an integral root k . Show that k is the mean of a, b, c, d .

Solution

a, b, c, d must be integers. $(k - a)(k - b)(k - c)(k - d) = 4$ and all of $(k - a), (k - b), (k - c), (k - d)$ are integers. They all divide 4, so they must belong to $\{-4, -2, -1, 1, 2, 4\}$. They are all distinct, so at most two of them have absolute value 1. Hence none of them can have absolute value 4 - or their product would be at least 8. Hence they must be $-2, -1, 1, 2$. Hence their sum is 0, so $4k = a + b + c + d$.

Problem B6

P is a variable point in space. Q is a fixed point on the z -axis. The plane normal to PQ through P cuts the x -axis at R and the y -axis at S . Find the locus of P such that PR and PS are at right angles.

Solution

Answer: sphere centre Q , radius QO , where O is the origin, excluding the two circles formed by the intersection of the sphere with the y - z plane and the x - z plane.

Let Q be $(0, 0, r)$. Let P be (a, b, c) . If R is $(x, 0, 0)$, then PR and PQ are perpendicular, so their dot product is 0, so $a(a - x) + b^2 + c(c - r) = 0$, hence $ax = (a^2 + b^2 + c^2 - cr)$. Similarly, if S is $(0, y, 0)$, then $by = (a^2 + b^2 + c^2 - cr)$. We require PR and PS perpendicular so $a(a - x) + (b - y)b + c^2 = 0$, hence $ax + by = a^2 + b^2 + c^2$. So $a^2 + b^2 + c^2 - 2cr = 0$ and hence $a^2 + b^2 + (c - r)^2 = r^2$, which shows that P lies on the sphere centre Q radius QO .

Conversely, suppose P lies on the sphere. Then SP and SO are tangents to the sphere and hence equal. Similarly, $RP = RO$, so PRS and ORS are similar. Hence $\angle RPS = \angle ROS = 90^\circ$. However, if P lies in the z - x plane, then S is not a finite point and if P lies in the y - z plane, then R is not a finite point. So we must exclude points lying on these two circles of the sphere.

8th Putnam 1948

Problem A1

C is the complex numbers. $f: C \rightarrow R$ is defined by $f(z) = |z^3 - z + 2|$. What is the maximum value of f on the unit circle $|z| = 1$?

Solution

Answer: $\sqrt{13}$.

Put $z = e^{i\theta}$, and use $\cos 2\theta = 2c^2 - 1$, $\cos 3\theta = 4c^3 - 3c$, where $\cos \theta = c$, to get: $|f(z)|^2 = 6 - 2\cos(3\theta - \theta) + 4\cos^3\theta - 4\cos\theta = 4(4c^3 - c^2 - 4c + 2)$. The cubic has stationary points where $12c^2 - 2c - 4 = 0$ or $c = 2/3$ or $-1/2$. So the maximum value is at $c = 1$ or -1 (the endpoints), or $2/3$ or $-1/2$ (the stationary points). Substituting in, we find that the maximum is actually at $c = -1/2$ with value $\sqrt{13}$.

Problem A2

K is a cone. s is a sphere radius r , and S is a sphere radius R . s is inside K touches it along all points of a circle. S is also inside K and touches it along all points of a circle. s and S also touch each other. What is the volume of the finite region between the two spheres and inside K ?

Solution

A straight slog is fairly long and tiresome.

Slice off the top of a sphere radius r by a cut a distance d from the surface. A simple integration shows that the volume removed is $\pi d^2(3r - d)/3$.

Let h be the distance from the vertex of the cone along the axis to the nearest sphere. Similar triangles gives $(h+r)/r = (h+2r+R)/R$. Hence $h = 2r^2/(R-r)$. Let the plane through the circle of contact between the small sphere and the cone cut the cone's axis at a distance $h+d$ from its vertex. Similarly, let the plane through the circle of contact of the other sphere cut the cone's axis a distance D from the point of contact between the two spheres. Let t be the distance from the cone's vertex to the circle of contact with the small sphere. Then $t^2 = h(h+2r)$. By similar triangles $(h+d)/t = t/(h+r)$, so $d = 2r^2/(R+r)$. Hence by similar triangles $D = 2rR/(R+r)$.

Let v be the volume in the cone between the vertex and the small sphere. We find this as the volume of a cone with circular base less the volume of a slice of sphere. The cone has height $h+d$. The square of the radius of the base is $(2rd-d^2)$. Hence $v = \pi/3 ((h+d)(2rd-d^2) - d^2(3r-d))$. This simplifies to $4r^5\pi/(3(R^2-r^2))$.

Hence V the volume of the corresponding region between the vertex and the large sphere (assuming the small sphere is temporarily removed) is $4r^2R^3\pi/(3(R^2-r^2))$. Hence the required volume is $V - v - 4\pi r^3/3 = 4\pi r^2R^2/3(R+r)$.

Problem A3

a_n is a sequence of positive reals decreasing monotonically to zero. b_n is defined by $b_n = a_n - 2a_{n+1} + a_{n+2}$ and all b_n are non-negative. Prove that $b_1 + 2b_2 + 3b_3 + \dots = a_1$.

Solution

We have that $b_1 + 2b_2 + 3b_3 + \dots + n b_n = a_1 - (n+1)a_{n+1} + na_{n+2} = a_1 - (n+1)(a_{n+1} - a_{n+2}) - a_{n+2}$. Also we are given that all b_i are non-negative. So $\sum b_n$ is monotonic increasing and bounded above by a_1 . So it converges to a limit $L \leq a_1$.

$b_n = (a_n - a_{n+1}) - (a_{n+1} - a_{n+2})$. Hence $b_n + b_{n+1} + b_{n+2} + \dots + b_{n+m} = (a_n - a_{n+1}) - (a_{n+m+1} - a_{n+m+2})$. But a_m tends to zero, so $(a_{n+m+1} - a_{n+m+2})$ tends to zero as m tends to infinity. Hence $b_n + b_{n+1} + b_{n+2} + \dots$ converges to $a_n - a_{n+1}$. Hence $(n+1)(b_{n+1} + b_{n+2} + b_{n+3} + \dots)$ converges to $(n+1)(a_{n+1} - a_{n+2})$. Hence $L \geq b_1 + 2b_2 + 3b_3 + \dots + n b_n + (n+1)(b_{n+1} + b_{n+2} + b_{n+3} + \dots) = a_1 - (n+1)a_{n+1} + n a_{n+2} + (n+1)(a_{n+1} - a_{n+2}) = a_1 - a_{n+2}$. But a_{n+2} tends to zero. Hence $L = a_1$.

Problem A4

Let D be a disk radius r . Given $(x, y) \in D$, and $R > 0$, let $a(x, y, R)$ be the length of the arc of the circle center (x, y) , radius R , which is outside D . Evaluate $\lim_{R \rightarrow 0} R^{-2} \int_D a(x, y, R) dx dy$.

Solution

Answer: $4\pi r$.

Let P be a point in the disk a distance x from its centre. Suppose that the circle centre P radius R cuts the disk

perimeter at A and B. Let angle APB be 2θ . We have $r^2 = x^2 + R^2 + 2xR \cos \theta$. Hence the length of arc outside the disk is $2R \cos^{-1}((r^2 - x^2 - R^2)/2xR)$. This applies for $r - R \leq x \leq r$. For smaller x the small circle lies entirely in the disk. Thus we have to evaluate $1/R^2 \int 2\pi x 2R \cos^{-1}((r^2 - x^2 - R^2)/2xR) dx$. Put $x = r - yR$, and we get $4\pi r \int_0^1 (1 - yR/r) \cos^{-1}((2Rry - (y^2 + 1)R^2)/2R(r - yR)) dy$. As R tends to 0, $(1 - yR/r)$ tends to 1 and $(2Rry - (y^2 + 1)R^2)/2R(r - yR)$ tends to y , so we get $4\pi r \int_0^1 \cos^{-1} y dy = 4\pi r (y \cos^{-1} y - \sqrt{1 - y^2})|_0^1 = 4\pi r$.

Problem A5

Let $\alpha_1, \alpha_2, \dots, \alpha_n$ be the n th roots of unity. Find $\prod_{i < j} (\alpha_i - \alpha_j)^2$.

Solution

Let $\alpha_k = \exp(2\pi i(k-1)/n)$. So $\alpha_1 = 1$, and the other α_k are roots of $x^{n-1} + x^{n-2} + \dots + x + 1 = 0$. Hence $(1 - \alpha_2), \dots, (1 - \alpha_n)$ are roots of $(1 - x)^{n-1} + \dots + (1 - x) + 1 = 0$. The coefficient of x^0 is n and the coefficient of x^{n-1} is $(-1)^{n-1}$. Hence $\prod_{i=2}^n (1 - \alpha_i) = n$.

Hence $n/\alpha_k = (\alpha_k - \alpha_1)(\alpha_k - \alpha_2) \dots (\alpha_k - \alpha_n)$, where the product excludes $(\alpha_k - \alpha_k)$. Hence $n^n/(\alpha_2 \dots \alpha_n) = \prod_{i \text{ not equ } j} (\alpha_i - \alpha_j) = (-1)^{n(n-1)/2} \prod_{i < j} (\alpha_i - \alpha_j)^2 = (-1)^{(n-1)(n-2)/2} n^n$.

Problem A6

Do either (1) or (2):

(1) On each element ds of a closed plane curve there is a force $1/R ds$, where R is the radius of curvature. The force is towards the center of curvature at each point. Show that the curve is in equilibrium.

(2) Prove that $x + 2/3 x^3 + 2 \cdot 4/3 \cdot 5 x^5 + \dots + 2 \cdot 4 \cdot \dots \cdot 2n/(3 \cdot 5 \cdot \dots \cdot 2n+1) x^{2n+1} + \dots = (1 - x^2)^{-1/2} \sin^{-1} x$.

Solution

(2) Let $f(x) = x + 2/3 x^3 + 8/15 x^5 + \dots + 1/2 n! n!/2n+1! (2x)^{2n+1} + \dots$.
Hence $x f(x) = x^2 + 2/3 x^4 + 8/15 x^6 + \dots + 1/4 n! n!/2n+1! (2x)^{2n+2} + \dots$.
Also $f'(x) = 1 + 2x^2 + 8/3 x^4 + \dots + n! n!/2n! (2x)^{2n} + \dots$.
and $(1 - x^2) f'(x) = 1 + x^2 + 2/3 x^4 + \dots + n-1! n-1!/2n! n/2 (2x)^{2n} + \dots$
 $= 1 + x^2 + 2/3 x^4 + \dots + 1/4 n-1! n-1!/2n-1! (2x)^{2n} + \dots$.

Hence the derivative of $\sqrt{1 - x^2} f(x)$ is $1/\sqrt{1 - x^2} (-x f(x) + (1 - x^2) f'(x)) = 1/\sqrt{1 - x^2}$. So $\sqrt{1 - x^2} f(x) = \text{const} + \sin^{-1}(x)$. Putting $x = 0$, we find that the constant is 0.

Problem B1

$p(x)$ is a cubic polynomial with roots α, β, γ and $p'(x)$ divides $p(2x)$. Find the ratios $\alpha : \beta : \gamma$.

Solution

$p(x) = (x - \alpha)(x - \beta)(x - \gamma)$. Hence $p'(x) = 3x^2 - 2(\alpha + \beta + \gamma)x + \beta\gamma + \gamma\alpha + \alpha\beta$. This must have roots $\alpha/2, \beta/2$. Hence $2(\alpha + \beta + \gamma)/3 = (\alpha + \beta)/2$ and $\alpha\beta/4 = (\beta\gamma + \gamma\alpha + \alpha\beta)/3$. This leads to $\alpha = (-2 + 2i/\sqrt{3})\gamma, \beta = (-2 - 2i/\sqrt{3})\gamma$.

Problem B2

A circle radius r is tangent to the three coordinate planes ($x=0, y=0, z=0$) in space. Find the locus of its center.

Problem B3

Show that $[\sqrt{n} + \sqrt{n+1}] = [\sqrt{4n+2}]$ for positive integers n .

Solution

We have $(\sqrt{n} + \sqrt{n+1})^2 = 2n + 1 + 2\sqrt{n^2 + n}$. Now $n^2 < n^2 + n < n^2 + n + 1/4 = (n + 1/2)^2$, so $\sqrt{4n+1} < \sqrt{n} + \sqrt{n+1} < \sqrt{4n+2}$. Hence $[\sqrt{4n+1}] \leq [\sqrt{n} + \sqrt{n+1}] \leq [\sqrt{4n+2}]$. But a square must be congruent to 0 or 1 mod 4, so $[\sqrt{4n+2}] = [\sqrt{4n+1}]$. Hence result.

Problem B4

R is the reals. For what λ can we find a continuous function $f : (0, 1) \rightarrow R$, not identically zero, such that $\int_0^1 \min(x, y) f(y) dy = \lambda f(x)$ for all $x \in (0, 1)$?

Problem B5

Find the area of the region $\{ (x, y) : |x + yt + t^2| \leq 1 \text{ for all } t \in [0, 1] \}$.

Solution

We cannot have $|x| > 1$, for then the inequality is not satisfied at $t = 0$. We cannot have $y > 1$ for then $yt + t^2 > 2$ at $t = 1$ and hence $x + yt + t^2 > 1$. Similarly, we cannot have $y < -3$ for then $yt + t^2 < -2$ at $t = 1$ and hence $x + yt + t^2 < -1$.

We can write $x + yt + t^2 = x - y^2/4 + (t + y/2)^2$. So for $-3 \leq y \leq -2$, the maximum of $(t + y/2)^2$ occurs at $t = 0$ with value $y^2/4$, and its minimum is $(1 + y/2)^2$ at $t = 1$. Hence the minimum of $x + yt + t^2$ is $1 + x + y$ and the maximum is x . So the region is bounded by $x \leq 1$ and $x + y \geq -2$.

For $-2 \leq y \leq -1$, the maximum of $(t + y/2)^2$ is $y^2/4$ at $t = 0$, but the minimum is 0 at $t = -y/2$. Hence the maximum of $x + yt + t^2$ is x and the minimum is $x - y^2/4$. So the region is bounded by $x \leq 1$ and $x - y^2/4 \geq -1$.

For $-1 \leq y \leq 0$, the maximum of $(t + y/2)^2$ is $(1 + y/2)^2$ and the minimum is 0 at $t = -y/2$. Hence the maximum of $x + yt + t^2$ is $1 + x + y$ and the minimum is $x - y^2/4$. So the region is bounded by $x + y \leq 0$ and $x - y^2/4 \geq -1$.

For $0 \leq y \leq 1$, the maximum of $(t + y/2)^2$ is $(1 + y/2)^2$ and the minimum is $y^2/4$. Hence the maximum of $x + yt + t^2$ is $1 + x + y$ and the minimum is x . So the region is bounded by $x + y \leq 0$ and $x \geq -1$.

Thus the region is the rhombus with vertices at $(1, -1)$, $(1, -3)$, $(-1, -1)$, $(-1, 1)$, except that the vertex $(-1, -1)$ is cut off by the curve $x = y^2/4 - 1$, which forms the boundary between $(0, -2)$ and $(-1, 0)$. The rhombus has area 4 (base 2 and height 2). The area between the parabola and the lines $x = -1$ and $y = -2$ is $\int_0^{-2} y^2/4 dy = 8/12$. The area between the line $y = -2 - x$ and the lines $x = -1$ and $y = -2$ is $1/2$, so the area cut off the rhombus by the parabola is $2/3 - 1/2 = 1/6$. Hence the area required is $4 - 1/6 = 3 \frac{5}{6}$.

Problem B6

Do either (1) or (2):

(1) Take the origin O of the complex plane to be the vertex of a cube, so that OA, OB, OC are edges of the cube. Let the feet of the perpendiculars from A, B, C to the complex plane be the complex numbers u, v, w . Show that $u^2 + v^2 + w^2 = 0$.

(2) Let (a_{ij}) be an $n \times n$ matrix. Suppose that for each i , $2|a_{ii}| > \sum_{j \neq i} |a_{ij}|$. By considering the corresponding system of linear equations or otherwise, show that $\det a_{ij} \neq 0$.

Solution

(2) If the det is non-zero, then we can find x_i not all zero so that $\sum x_i a_{ij} = 0$ for each j . Take k so that $|x_k| \geq |x_i|$ for all i . Then $|\sum_{i \neq k} x_i a_{ik}| \leq \sum_{i \neq k} |x_i a_{ik}| \leq |x_k| \sum_{i \neq k} |a_{ik}| < |x_k| |a_{kk}|$, so we cannot have $\sum x_i a_{ik} = 0$. Contradiction.

9th Putnam 1949

Problem A1

Do either (1) or (2)

(1) Let L be the line through $(0, -a, a)$ parallel to the x -axis, M the line through $(a, 0, -a)$ parallel to the y -axis, and N the line through $(-a, a, 0)$ parallel to the z -axis. Find the equation of S , the surface formed from the union of all lines K which intersect each of L , M and N .

(2) Let S be the surface $xy + yz + zx = 0$. Which planes cut S in circles? In parabolas?

Problem A2

Take points O, P, Q, R in space. Let the volume of the parallelepiped with edges OP, OQ, OR be V . Let V' be the volume of the parallelepiped which has O as one vertex and which has OP, OQ, OR as altitudes to three faces. Show that $V V' = OP^2 OQ^2 OR^2$. Generalize to n dimensions.

Problem A3

All the complex numbers z_n are non-zero and $|z_m - z_n| > 1$ (for any $m \neq n$). Show that $\sum 1/z_n^3$ converges.

Solution

We show that it is absolutely convergent. Take a disk radius $1/2$ centred on each z_n . The disks must be disjoint since $|z_m - z_n| > 1$ for distinct m, n . Now consider how many z_n can lie in the annulus $N \leq |z| < N+1$. If z_n lies in the annulus, then at least half the corresponding disk must also, so if M is the number of z_n , then $M \pi/8 < \pi ((N+1)^2 - N^2) = (2N+1) \pi$, and the sum of $1/|z_n|^3$ for these z_n is at most $M/N^3 = 8(2N+1)/N^3$. But $\sum 8(2N+1)/N^3$ converges, since $\sum 1/N^2$ converges.

Problem A4

Take P inside the tetrahedron $ABCD$ to minimize $PA + PB + PC + PD$. Show that $\square APB = \square CPD$ and that the bisector of APB also bisects CPD .

Problem A5

Let $p(z) \equiv z^6 + 6z + 10$. How many roots lie in each quadrant of the complex plane?

Solution

There are no real roots, since $p(z)$ has a minimum (for real value of z) at $z = -1$ of 5 . Similarly there are no purely imaginary roots, since $p(iy)$ has imaginary part $6iy$ (and $z = 0$ is not a root). Also $a + ib$ is a root iff $a - ib$ is a root, so either there is one root in each of the 1st and 4th quadrants and two in each of the 2nd and 3rd, or vice versa.

The argument principle states that the change in $\arg p(z)$ as we move (anti-clockwise) round a closed contour which passes through no zeros and contains k zeros is $2\pi k$. Take a contour in the first quadrant from $z = 0$ to $z = X$, then around the arc $|z| = X$ to iX , then back along the imaginary axis to $z = 0$. Moving along the real axis, $\arg p(z)$ does not change. Moving along the arc the change will be the same as that in $\arg z^6$ for X sufficiently large and hence will be 3π . As we move back down the imaginary axis, the imaginary part remains positive, so $p(z)$ is confined to the upper half plane, and we end up at $p(z) = 10$, so the change must be $-\pi$. Thus the change for a complete circuit is 2π , and so there is just one root in the first quadrant.

Problem A6

Show that $\prod_{n=1}^{\infty} (1 + 2 \cos(2z/3^n))/3 = (\sin z)/z$ for all complex z .

Solution

For $z = 0$, we may take $(\sin z)/z = 1$ by continuity. In this case, each term on the lhs is 1, so the relation holds. Assume now that z is non-zero.

We have $\sin z = \sin(2z/3 + z/3) = \sin 2z/3 \cos z/3 + \cos 2z/3 \sin z/3$, and $\sin z/3 = \sin(2z/3 - z/3) = \sin 2z/3 \cos z/3 - \cos 2z/3 \sin z/3$. Hence $\sin z = (1 + \cos 2z/3) \sin z/3$. Hence $(\sin z)/z = (\sin z/3)/(z/3) (1 + \cos 2z/3)/3$. Iterating: $(\sin z)/z = (\sin z/3^N)/(z/3^N) \prod_{n=1}^N (1 + 2 \cos(2z/3^n))/3$. But as N tends to infinity $z/3^N$ tends to zero and hence $(\sin z/3^N)/(z/3^N)$ tends to 1. Hence the result.

Problem B1

Show that for any rational $a/b \in (0, 1)$, we have $|a/b - 1/\sqrt{2}| > 1/(4b^2)$.

Solution

If $|a/b - 1/\sqrt{2}| \leq 1/(4b^2)$, then $|a^2/b^2 - 1/2| = |a/b - 1/\sqrt{2}| |a/b + 1/\sqrt{2}| < |a/b - 1/\sqrt{2}| (1 + 1) < 1/2b^2$. So $|2a^2 - b^2| < 1$. Hence $2a^2 = b^2$. But that is impossible since $\sqrt{2}$ is irrational. Note that the restriction to the interval $(0, 1)$ is unnecessary.

Problem B2

Do either (1) or (2)

(1) Prove that $\sum_2^\infty \cos(\ln \ln n) / \ln n$ diverges.

(2) Let k, a, b, c be real numbers such that $a, k > 0$ and $b^2 < ac$. Show that $\int_U (k + ax^2 + 2bxy + cy^2)^{-2} dx dy = \pi / (k \sqrt{ac - b^2})$, where U is the entire plane.

Solution

(1) We have $\ln n^2 = 2 \ln n$, so for any m in the range $n, n+1, n+2, \dots, n^2$ we have $1/\ln m > 1/2 \cdot 1/\ln n$. Also $\ln \ln n^2 = \ln \ln n + \ln 2 < \ln \ln n + \pi/3$. So given any sufficiently large integer N , take $x > N$ so that $\ln \ln x = a$ a multiple of 2π , then we can find at least $N^2/2$ consecutive integers n in the range x to x^2 such that (A) $1/\ln n > 1/(2 \ln N)$ and (B) $\ln \ln n$ lies between $2k\pi$ and $2k\pi + \pi/3$ (for some integer k) and hence so that $\cos \ln \ln n > 1/2$. Hence for each such integer $(\cos \ln \ln n)/\ln n > 1/(4 \ln N)$. So $\sum (\cos \ln \ln n)/\ln n$ over all these terms is at least $N^2/(8 \ln N)$ which can be made arbitrarily large. Hence the sequence does not converge.

(2) $ax^2 + 2bxy + cy^2 = h$ is the equation of an ellipse. If we rotate the axes to line up with its axes we get new coordinates X, Y so that it becomes $AX^2 + BY^2 = h$ with $AB = ac - b^2$. Thus the integral is transformed to $\int_U (k + AX^2 + BY^2)^{-2} dX dY$. Take $u = X\sqrt{A}, v = Y\sqrt{B}$ and the integral becomes $1/\sqrt{ac - b^2} \int_U (k + u^2 + v^2)^{-2} du dv$. Now take $u = r \cos \theta, v = r \sin \theta$ and we get $1/\sqrt{ac - b^2} \int_U (k + u^2 + v^2)^{-2} du dv = 1/\sqrt{ac - b^2} \int_U (k + r^2)^{-2} r dr d\theta = 2\pi / \sqrt{ac - b^2} \int (k + r^2)^{-2} r dr = 2\pi / (k \sqrt{ac - b^2}) \int (1 + s^2)^{-2} s ds = \pi / (k \sqrt{ac - b^2}) (-1/(1 + s^2))_0^\infty = 2\pi / (k \sqrt{ac - b^2})$.

Problem B3

C is a closed plane curve. If $P, Q \in C$, then $|PQ| < 1$. Show that we can find a disk radius $1/\sqrt{3}$ which contains C .

Solution

C can be any set of points in the plane. The usual proof uses Helly's theorem. Given any three points of the set we can find a circle radius $1/\sqrt{3}$ which contains them. This is fairly obvious. Take the points to be P, Q, R with PQ the longest side of the triangle. Take S on the same side of PQ as R so that PQS is equilateral. Then R must lie inside the bounding circle of PQS (in fact it must lie inside the region bounded PQ , the arc of the circle centre P from S to Q , and the arc of the circle centre Q from S to P , and this region lies entirely inside the bounding circle of PQS).

Another way of expressing this is that the three circles radius $1/\sqrt{3}$ and centres P, Q, R have a point in common. We can now apply Helly's theorem to the set of circles radius $1/\sqrt{3}$ and with centres at all the points of the set C . Every three of these circles have a point in common (proved above) and hence by Helly's theorem there is a point common to all of them. But this point can be taken as the centre of the required circle.

Problem B4

Let $(1 + x - \sqrt{x^2 - 6x + 1})/4 = \sum_1^\infty a_n x^n$. Show that all a_n are positive integers.

Solution

We have $1 + x - \sqrt{x^2 - 6x + 1} = 4 \sum a_n x^n$, and hence $1 + x + \sqrt{x^2 - 6x + 1} = 2x + 2 - 4 \sum a_n x^n$. Multiplying, $1 + 2x + x^2 - x^2 + 6x - 1 = 8x = 8(a_1x + a_2x^2 + \dots) (1 + (1 - 2a_1)x - 2a_2x^2 - 2a_3x^3 - \dots)$. Hence:

$$a_1 = 1$$

$$a_2 + a_1(1 - 2a_1) = 0$$

$$a_3 + a_2(1 - 2a_1) - 2a_1a_2 = 0$$

$$a_4 + a_3(1 - 2a_1) - 2a_2a_2 - 2a_1a_3 = 0 \dots$$

The n th equation has the form $a_n +$ combination of a_1, \dots, a_{n-1} with integer coefficients, so by a simple induction, a_n is integral.

Problem B5

a_n is a sequence of positive reals. Show that $\limsup_{n \rightarrow \infty} (a_1 + a_{n+1})/a_n \geq e$.

Solution

It is sufficient to show that $(a_1 + a_{n+1})/a_n \geq 1 + 1/n$ for infinitely many n . For these n then form a sequence for which $((a_1 + a_{n+1})/a_n)^n$ has a limit point (possibly infinity) not less than $\lim (1 + 1/n)^n = e$.

If that were false, then we would have $(a_1 + a_{n+1})/a_n < 1 + 1/n$ for all n not less than some N . In particular, taking $n = N, N+1, N+2, \dots$ we get $a_1/N+1 + a_{N+1}/N+1 < a_N/N$, $a_1/N+2 + a_{N+2}/N+2 < a_{N+1}/N+1$, $a_1/N+3 + a_{N+3}/N+3 < a_{N+2}/N+2$, \dots . Hence $a_n/N > a_1(1/N+1 + 1/N+2 + \dots + 1/N+M) + a_{N+M}/N+M > a_1(1/N+1 + 1/N+2 + \dots + 1/N+M)$. But this holds for any M and $(1/N+1 + 1/N+2 + \dots)$ diverges. Contradiction.

Problem B6

C is a closed convex curve. If P lies on C and T_P is the tangent at P , then T_P varies continuously with P . Let O be a point inside C . Given a point P on C , define $f(P)$ to be the point where the perpendicular from O to T_P intersects C . Given P_1 , define the sequence P_n by $P_{n+1} = f(P_n)$. Assume that f is continuous and that, for each P , C lies entirely on one side of T_P . Show that P_n converges. Find $S = \{ P : P = \lim_{n \rightarrow \infty} P_n \text{ for some } P_1 \}$.

10th Putnam 1950

Problem A1

a and b are positive reals and $a > b$. Let C be the plane curve $r = a - b \cos \theta$. For what values of b/a is C convex?

Solution

The curvature is $(x'y'' - x''y')/(x'^2 + y'^2)^{3/2}$. So for convexity we require $x'y'' \geq x''y'$. Putting $k = b/a$, $c = \cos \theta$, $s = \sin \theta$, we have $x/a = c - kc^2$, $x'/a = -s + 2kcs$, $x''/a = -c - 2ks^2 + 2kc^2$ and $y/a = s - ksc$, $y'/a = c - kc^2 + ks^2$, $y''/a = -s + 4ksc$. Thus the condition becomes after a little cancellation, $1 + 2k^2 - 3kc \geq 0$. c takes values in the range -1 to 1 , so for the condition to be true for all points of the curve we require $1 + 2k^2 - 3k \geq 0$. But $1 + 2k^2 - 3k = (2k - 1)(k - 1)$. We are given that $k < 1$, so we must have $k < 1/2$.

For small k , the curve is approximately a circle centred on the origin. As k increases it develops a flattening near $x = a$, $y = 0$. For $k > 1/2$, this becomes a dimple in the surface, so that convexity is broken. For $k = 1$, the depression in the surface extends as far as the centre (the origin). For $k > 0$, the curve intersects itself at the origin so that it comprises two ovals one inside the other.

Problem A2

Does the series $\sum_{n=2}^{\infty} 1/\ln n!$ converge? Does the series $1/3 + 1/(3 \cdot 3^{1/2}) + 1/(3 \cdot 3^{1/2} \cdot 3^{1/3}) + \dots + 1/(3 \cdot 3^{1/2} \cdot 3^{1/3} \dots 3^{1/n}) + \dots$ converge?

Solution

(1) $\ln n! < n \ln n$, so $\sum_{n=2}^{\infty} 1/\ln n! > \sum_{n=2}^{\infty} 1/(n \ln n)$, which diverges. For $\int 1/(x \ln x) = \ln \ln x$.

(2) We have $1 + 1/2 + 1/3 + \dots + 1/n$ is approx $\ln n$, and $1/3^{\ln n}$ is approx $1/n^{\ln 3}$. But $\ln 3 > 1$, so we expect this to converge. In fact $1 + 1/2 + 1/3 + \dots + 1/n > \int_1^{n+1} dx/x = \ln(n+1) > \ln n$. So $1/3 + 1/(3 \cdot 3^{1/2}) + 1/(3 \cdot 3^{1/2} \cdot 3^{1/3}) + \dots + 1/(3 \cdot 3^{1/2} \cdot 3^{1/3} \dots 3^{1/n}) + \dots < 1/3^{\ln 2} + 1/3^{\ln 3} + 1/3^{\ln 4} + \dots = 1/1 + 1/2^k + 1/3^k + \dots$, where $k = \ln 3$, which converges.

Problem A3

The sequence a_n is defined by $a_0 = \alpha$, $a_1 = \beta$, $a_{n+1} = a_n + (a_{n-1} - a_n)/(2n)$. Find $\lim a_n$.

Solution

$a_{n+1} - a_n = -1/2n (a_n - a_{n-1})$, hence $a_{n+1} - a_n = (-1)^n/(2 \cdot 4 \cdot 6 \dots \cdot 2n) (a_1 - a_0)$. Then $a_n = (a_n - a_{n-1}) + (a_{n-1} - a_{n-2}) + \dots + (a_1 - a_0) + a_0$.

Hence the limit is $\beta + (\beta - \alpha)/2 - (\beta - \alpha)/2 \cdot 4 + (\beta - \alpha)/2 \cdot 4 \cdot 6 - \dots$. But $1/2 - 1/2 \cdot 4 + 1/2 \cdot 4 \cdot 6 - \dots = 1 - 1/\sqrt{e}$, so the limit is $\alpha + (\beta - \alpha)/\sqrt{e}$.

Problem A4

Do either (1) or (2)

(1) P is a prism with triangular base. A is a vertex. The total area of the three faces containing A is $3k$. Show that if the volume of P is maximized, then each of the three faces has area k and the two lateral faces are perpendicular to each other.

(2) Let $f(x) = x + x^3/(1 \cdot 3) + x^5/(1 \cdot 3 \cdot 5) + x^7/(1 \cdot 3 \cdot 5 \cdot 7) + \dots$, and $g(x) = 1 + x^2/2 + x^4/(2 \cdot 4) + x^6/(2 \cdot 4 \cdot 6) + \dots$. Show that $\int_0^x \exp(-t^2/2) dt = f(x)/g(x)$.

Solution

(1) Let the sides of the base at A be a , b with angle between them θ . Let the height be h . Then the volume is $V = 1/2 abh \sin \theta$ and $3k = ah + bh + 1/2 ab \sin \theta$. Now $V^2 = (1/2 ab \sin \theta) (ah) (bh) 1/2 \sin \theta$. By the arithmetic geometric mean theorem we have that $(1/2 ab \sin \theta) (ah) (bh) \leq k^3$, with equality iff $ah = bh = 1/2 ab \sin \theta$. So $V^2 \leq k^3/2 \sin \theta$. Hence $V \leq \sqrt{k^3/2}$ with equality iff $\theta = \pi/2$ and $ah = bh = 1/2 ab$.

(2) $\exp(-t^2/2) = 1 - t^2/2 + t^4/2 \cdot 4 - t^6/2 \cdot 4 \cdot 6 + \dots$. We notice that this is similar to $g(t)$. In fact $g(t) = \exp(t^2/2)$. So the required relation is $f(x) = \exp(x^2/2) \int_0^x \exp(-t^2/2) dt$. Differentiating gives $f'(x) = x f(x) + 1$. All that is just motivation.

We start from the series for $f(x)$. By the ratio test it is absolutely convergent, so we may differentiate term by term to get $f'(x) = 1 + x^2/1 + x^4/1 \cdot 3 + x^6/1 \cdot 3 \cdot 5 + \dots = x f(x) + 1$. Multiplying this by $\exp(-x^2/2)$ we get $\exp(-x^2/2) f'(x) - x \exp(-x^2/2) f(x) = \exp(-x^2/2)$. Integrating and using $f(0) = 0$, we get $\exp(-x^2/2) f(x) = \int_0^x \exp(-t^2/2) dt$ or $f(x)/g(x) = \int_0^x \exp(-t^2/2) dt$.

Problem A5

Let N be the set of natural numbers $\{1, 2, 3, \dots\}$. Let Z be the integers. Define $d : N \rightarrow Z$ by $d(1) = 0$, $d(p) = 1$ for p prime, and $d(mn) = m d(n) + n d(m)$ for any integers m, n . Determine $d(n)$ in terms of the prime factors of n . Find all n such that $d(n) = n$. Define $d_1(m) = d(m)$ and $d_{n+1}(m) = d(d_n(m))$. Find $\lim_{n \rightarrow \infty} d_n(63)$.

Solution

$d(p^a) = a p^{a-1}$ by a trivial induction on a . Hence for $n = p^a q^b \dots$ we have $d(n) = n (a/p + b/q + \dots)$ by a trivial induction on the number of primes.

Hence $d(n) = n$ iff $a/p + b/q + \dots = 1$. Multiplying through by all the prime denominators gives integral terms. All but the first are clearly divisible by p , so the first must be also. Hence $a/p = 1$ and b/q etc are zero. In other words, $n = p^p$ for some prime p .

We find $d_1(63) = 51$, $d_2(63) = 20$, $d_3(63) = 24$, $d_4(63) = 44$, $d_5(63) = 48$. Now suppose n is divisible by 16. Then $n = 2^k p^a q^b \dots$, where p, q, \dots are all odd and $k \geq 4$. Hence $d(n) = n(4/2 + a/p + b/q + \dots)$. Now all of $2n, na/p, nb/q, \dots$ are integral and multiples of 16. So $d(n)$ is at least twice n and a multiple of 16. So if we have $d_k(m)$ divisible by 16, then $d_{n+k}(m) \geq 2^h$ which diverges. Hence $d_k(63)$ tends to infinity.

Problem A6

Let $f(x) = \sum_{n=0}^{\infty} a_n x^n$ and suppose that each $a_n = 0$ or 1. Do either (1) or (2):

(1) Show that if $f(1/2)$ is rational, then $f(x)$ has the form $p(x)/q(x)$ for some integer polynomials $p(x)$ and $q(x)$.

(2) Show that if $f(1/2)$ is not rational, then $f(x)$ does not have the form $p(x)/q(x)$ for any integer polynomials $p(x)$ and $q(x)$.

Solution

(1) $f(1/2)$ is the binary expansion of some real in the closed interval $[0, 1]$. So if it is rational, then the binary expansion is periodic after some point. In other words, there are positive integers N and k such that $a_n = a_{n+k}$ for $n > N$. Hence $f(x) = a_0 + a_1 x + \dots + a_N x^N + (a_{N+1} x^{N+1} + \dots + a_{N+k-1} x^{N+k-1})(1 + x^k + x^{2k} + \dots) = a_0 + a_1 x + \dots + a_N x^N + (a_{N+1} x^{N+1} + \dots + a_{N+k-1} x^{N+k-1})/(1 - x^k)$. Hence we can take $q(x) = 1 - x^k$ and $p(x) = (a_0 + a_1 x + \dots + a_N x^N)(1 - x^k) + (a_{N+1} x^{N+1} + \dots + a_{N+k-1} x^{N+k-1})$.

(2) is trivial. If $f(x) = p(x)/q(x)$, then $f(1/2) = p(1/2)/q(1/2)$ which is rational. The only slight complication is if $q(1/2) = 0$. But we can assume that $p(x)/q(x)$ is in lowest terms, so if $q(1/2) = 0$, then $p(x)$ is non-zero, but then $p(x) = f(x) q(x)$. We know that $f(1/2) \leq 1/2 + 1/4 + 1/8 + \dots = 1$, so $f(1/2) q(1/2) = 0$. Hence $p(1/2) = 0$. Contradiction.

Problem B1

Given n , not necessarily distinct, points P_1, P_2, \dots, P_n on a line. Find the point P on the line to minimize $\sum |PP_i|$.

Solution

Assume that the points are in the order given. For two points A and B , $|PA| + |PB| = AB$ for P on the segment AB and $|PA| + |PB| > AB$ for P outside it. Thus we minimise $|PP_1| + |PP_n|$ by placing P between P_1 and P_n . Similarly, we minimise $|PP_2| + |PP_{n-1}|$ by placing P between P_2 and P_{n-1} . And so on. But note that if we minimise $|PP_2| + |PP_{n-1}|$ then we also minimise $|PP_1| + |PP_n|$. Thus for n odd we must take P to be the central point, for n even we can take P to be any point on the segment between the two central points.

Problem B2

An ellipse with semi-axes a and b has perimeter length $p(a, b)$. For b/a near 1, is $\pi(a + b)$ or $2\pi \sqrt{ab}$ the better approximation to $p(a, b)$?

Solution

Answer: $\pi(a + b)$.

Put $b^2 = a^2(1 - \epsilon)$. The perimeter is $4 \int_0^{\pi/2} (a^2 \sin^2 t + b^2 \cos^2 t)^{1/2} dt = 4a \int (1 - \epsilon \cos^2 t)^{1/2} dt = 4a \int (1 - 1/2 \epsilon \cos^2 t - 1/8 \epsilon^2 \cos^4 t + O(\epsilon^4)) dt = 2\pi a (1 - \epsilon/4 - 3\epsilon^2/64 + O(\epsilon^4))$.

$b = a(1 - \epsilon)^{1/2} = a(1 - \epsilon/2 - \epsilon^2/8 + O(\epsilon^3))$. Hence $\pi(a + b) = 2\pi a (1 - \epsilon/4 - 4\epsilon^2/64 + O(\epsilon^3))$. Whilst $2\pi\sqrt{ab} = 2\pi a (1 - \epsilon/2 - \epsilon^2/8 + O(\epsilon^3))^{1/2} = 2\pi a (1 - \epsilon/4 - 6\epsilon^2/64 + O(\epsilon^3))$.

Hence the error in $\pi(a + b)$ is $\pi a \epsilon^2/32 + O(\epsilon^3)$, and the error in $2\pi\sqrt{ab}$ is $3\pi a \epsilon^2/32 + O(\epsilon^3)$, which is roughly three times as large for small ϵ .

Problem B3

Leap years have 366 days; other years have 365 days. Year $n > 0$ is a leap year iff (1) 4 divides n , but 100 does not divide n , or (2) 400 divides n . n is chosen at random from the natural numbers. Show that the probability that December 25 in year n is a Wednesday is not $1/7$.

Solution

$365 = 52 \cdot 7 + 1$, so in non-leap years Christmas day is a day later in the week than the previous year. In leap years it is two days later. Now consider a period of 400 years. There are 100 multiples of 4, 3 of which are multiples of 100 but not 400, so there are 97 leap years. Thus after 400 years the day has advanced $303 \cdot 1 + 97 \cdot 2 = 497$ days or exactly 41 weeks. In other words after 400 years Christmas is on the same day of the week. This cycle then repeats indefinitely. But 400 is not a multiple of 7, so each day cannot occur with exactly probability $1/7$.

If we look in more detail at the 400 years we find that Su, Tu, Fr each occur 58 times, We and Th each occur 57 times, and Mo and Sa each occur 56 times (compared with the expected $57 \frac{1}{7}$).

Problem B4

A long, light cylinder has elliptical cross-section with semi-axes $a > b$. It lies on the ground with its main axis horizontal and the major axes horizontal. A thin heavy wire of the same length as the cylinder is attached to the line along the top of the cylinder. [We could take the cylinder to be the surface $|x| \leq L, y^2/a^2 + z^2/b^2 = 1$. Contact with the ground is along $|x| \leq L, y = 0, z = -b$. The wire is along $|x| \leq L, y = 0, z = b$.] For what values of b/a is the cylinder in stable equilibrium?

Solution

Take the ellipse to be $x^2/a^2 + y^2/b^2 = 1$. We need the normal to the ellipse at a point near $(0, -b)$ to meet the y -axis above $(0, b)$ (because then the heavy wire will be the correct side of the vertical through the point of contact to right the cylinder).

The gradient at $(a \cos t, b \sin t)$ is $-b/a \cot t$, so the normal has slope $a/b \tan t$, so its equation is $(y - b \sin t) = a/b \tan t (x - a \cos t)$. This meets the y -axis at $y = b \sin t - a^2/b \sin t$. For points near $(0, -b)$, $\sin t$ is $-(1 - \epsilon)$ so we need $a^2/b - b > b$ or $a > b\sqrt{2}$.

Problem B5

Do either (1) or (2):

(1) Show that if $\sum(a_n + 2a_{n+1})$ converges, then so does $\sum a_n$.

(2) Let S be the surface $2xy = z^2$. The surface S and the variable plane P enclose a cone with volume $\pi a^3/3$, where a is a positive real constant. Find the equation of the envelope of P . What is the envelope in the case of a general cone?

Solution

(1) Suppose $a_n + 2a_{n+1}$ converges to $3k$. We show that a_n converges to k .

Given any $\epsilon > 0$, take N so that $a_n + 2a_{n+1}$ is within ϵ of $3k$ for all $n \geq N$. Take a positive integer M such that a_N is within $(2^M + 1)\epsilon$ of k .

Then a_{N+1} is within $((2^M + 1)\epsilon + \epsilon)/2 = (2^{M-1} + 1)\epsilon$ of $(3k - k)/2 = k$. By a trivial induction a^{N+M} is within 2ϵ of k .

Then a^{N+M+1} is within $(2\epsilon + \epsilon)/2$, and hence within ϵ , of k . So by a trivial induction, a_n is within 2ϵ of k for all $n > N + M$.

(2) Answer: the 2-sheet hyperboloid $u^2 = v^2 + z^2 + a^2$.

The nature of the surface S becomes much clearer if we change coordinates to u, v, z with z unchanged and the x, y

axes rotated through $\pi/4$ to $u = (x + y)/\sqrt{2}$, $v = (-x + y)/\sqrt{2}$ (or inversely, $x = (u - v)/\sqrt{2}$, $y = (u + v)/\sqrt{2}$). Now S becomes $u^2 - v^2 = z^2$ or $v^2 + z^2 = u^2$, which is evidently a right circular cone with vertex the origin and axis the u -axis.

Evidently the plane $u = a$ forms a cone volume $1/3 \pi a^3$ with the surface (the base is a circle radius a and the height is a). If the plane moves round to be almost tangent to the sides of the cone it should still cut off a cone with the same volume, so we might suspect that the envelope is a two sheet hyperboloid asymptotic to the cone and passing through the points $u = \pm a$, $v = z = 0$. This has equation $u^2 = v^2 + z^2 + a^2$. This is a surface of revolution formed by rotating the hyperbola $u^2 = v^2 + a^2$ about the u -axis, so it is sufficient to look at tangent planes that are parallel to (ie do not intersect) the z -axis. We need to show that these cut off cones volume $1/3 \pi a^3$.

The tangent to the hyperbola at $u = a \cosh t$, $v = a \sinh t$ is $(v - a \sinh t) = \coth t (u - a \cosh t)$ or $v \sinh t - u \cosh t + a = 0$ (*). It cuts $v = u$ at $u = v = a/(\cosh t - \sinh t)$ and it cuts $v = -u$ at $u = -v = a/(\cosh t + \sinh t)$. So the corresponding plane cuts the hyperboloid in an ellipse with major axis length $2A$, where $2A$ is the distance between these two points of intersection, which is $2a \cosh 2t$. At the centre of the ellipse (which we find as midway between the two points of intersection) we have $u = a \cosh t$, $v = a \sinh t$, so the extremities of minor axis have $z^2 = u^2 - v^2 = a^2$. Thus the minor axis has length $2a$. Hence the area of the ellipse is $\pi a^2 \sqrt{2 \cosh 2t}$. The distance of the origin from the plane is its distance from the line (*) which is $a/\sqrt{2 \cosh 2t}$. Hence the volume of the cone is $1/3$ area of base \times height = $1/3 \pi a^3$, as we had guessed.

Finally, if we have a general cone (which does not have circular cross-section or right-angle), we can transform it into a right circular cone by an affine transformation. Such transformations preserve the ratios of distances and hence the ratios of volumes. They also preserve tangency, so the envelope in this more general case will still be a hyperboloid asymptotic to the cone.

Problem B6

- (1) The convex polygon C' lies inside the polygon C . Is it true that the perimeter of C' is no longer than the perimeter of C ?
- (2) C is the convex polygon with shortest perimeter enclosing the polygon C' . Is it true that the perimeter of C is no longer than the perimeter of C' ?
- (3) The closed convex surface S' lies inside the closed surface S . Is it true that $\text{area } S' \leq \text{area } S$?
- (4) S is the smallest convex surface containing the closed surface S' . Is it true that $\text{area } S \leq \text{area } S'$?

Solution

(1) True. On each side of C' take a semi-infinite strip perpendicular to the side and extending outwards from C' . Since C' is convex, these strips are all disjoint. But the intersection of C with each strip is at least as long as the corresponding side of C' , hence the total length of C that intersects the strips is at least as long as the perimeter of C' .

(2) True. Take C as the convex hull of C' . It is sufficient to prove that the perimeter of C is no longer than the perimeter of C' . Evidently the vertices of C form a subset of the vertices of C' . If C' includes a vertex P which is not a vertex of C , then removing it does not increase the perimeter of C' (by the triangle inequality) and leaves C unaffected. So we can assume that C and C' have the same vertices. If C' just has the vertices of C permuted cyclically (or in reverse order), then its perimeter is the same. If not, then there are vertices A, B which are adjacent in C' but not in C . We show that in this case we can permute the vertices of C' so as to shorten the perimeter. We can repeat this process, shortening the perimeter each time. But there are only finitely many permutations so we must eventually reach a perimeter with the same length as C .

Suppose C' has vertices in the order $X_1 X_2 \dots X_n$. wlog we may assume that X_1 and X_2 are not adjacent in C , so we can find X_i on one side of $X_1 X_2$ and X_j on the other. Then one of the pairs $X_i X_{i+1}$, $X_{i+1} X_{i+2}$, ..., $X_{j-1} X_j$ must straddle $X_1 X_2$. Suppose it is $X_k X_{k+1}$. Consider the quadrilateral $X_1 X_k X_2 X_{k+1}$. Suppose its diagonals $X_1 X_2$ and $X_k X_{k+1}$ meet at O . Then $X_1 X_2 + X_k X_{k+1} = X_1 O + O X_2 + X_k O + O X_{k+1} = (X_1 O + O X_k) + (X_2 O + O X_{k+1}) > X_1 X_k + X_2 X_{k+1}$. So if we take C'' to have vertices in the order $X_1 X_k X_{k-1} \dots X_3 X_2 X_{k+1} X_{k+2} \dots X_n$, then C'' has shorter perimeter.

(3) True. The same argument as (1) works. Take semi-infinite columns on each face of the inner polyhedron.

(4) False. Take the 4 vertices of a regular tetrahedron and its centre. We can find a surface S' with arbitrarily small area which contains these 5 points, but the convex hull S is the tetrahedron.

11th Putnam 1951

Problem A1

A is a skew-symmetric real 4×4 matrix. Show that $\det A \geq 0$.

Solution

Let $A =$

$$\begin{pmatrix} 0 & a & b & c \\ -a & 0 & d & e \\ -b & -d & 0 & f \\ -c & -e & -f & 0 \end{pmatrix}$$

Then $\det A = a^2f^2 + 2acdf - 2abef + b^2e^2 - 2bcde + c^2d^2$.

At this point it is helpful to notice that a only appears with f , b with e , and c with d . So putting $X = af$, $Y = cd$, $Z = be$, we have that $\det A = X^2 + Y^2 + Z^2 + 2XY - 2XZ - 2YZ$. This easily factorizes as $(X + Y - Z)^2$.

Problem A2

k is a positive real and P_1, P_2, \dots, P_n are points in the plane. What is the locus of P such that $\sum PP_i^2 = k$? State in geometric terms the conditions on k for such points P to exist.

Solution

Let P_i have coordinates (a_i, b_i) . Take the origin at the centroid of the points, so that $\sum a_i = \sum b_i = 0$. Then if P has coordinates (x, y) , we have $\sum PP_i^2 = \sum((x - a_i)^2 + (y - b_i)^2) = n(x^2 + y^2) + \sum(a_i^2 + b_i^2)$. Let the origin be O , so that $\sum(a_i^2 + b_i^2) = \sum OP_i^2$. Then if $\sum OP_i^2 > k$, the locus is empty. Otherwise it is the circle with centre at the centroid of the n points and radius $\sqrt{(k - \sum OP_i^2)/n}$.

Problem A3

Find $\sum_{0}^{\infty} (-1)^n / (3n + 1)$.

Solution

We have $\ln(1 + x) = x - x^2/2 + x^3/3 - x^4/4 + \dots$ (1). Put $\omega = e^{2\pi i/3}$, so that $\omega^3 = 1$ and $1 + \omega + \omega^2 = 0$. Then $\omega^2 \ln(1 + \omega x) = x - \omega x^2/2 + \omega^2 x^3/3 - x^4/4 + \dots$ (2) and $\omega \ln(1 + \omega^2 x) = x - \omega^2 x^2/2 + \omega x^3/3 - x^4/4 + \dots$ (3).

Adding (1), (2), (3) we get $3(x - x^4/4 + x^7/7 - \dots) = \ln(1 + x) + \omega^2 \ln(1 + \omega x) + \omega \ln(1 + \omega^2 x)$. Hence the required series sums to $1/3 (\ln 2 + \omega^2 \ln(1 + \omega) + \omega \ln(1 + \omega^2))$.

If $k = \ln(1 + \omega)$, then $e^k = 1 + \omega = -\omega^2 = e^{i\pi/3}$, so $k = i\pi/3$. Hence $\omega^2 \ln(1 + \omega) = -(1 + i\sqrt{3})/2 \cdot i\pi/3 = \pi/2\sqrt{3} - i\pi/6$. Similarly, $\omega \ln(1 + \omega^2) = -i\pi/3 (-1/2 + i\sqrt{3}/2) = \pi/2\sqrt{3} + i\pi/6$. Hence the series sums to $1/3 \ln 2 + \pi/3\sqrt{3}$.

Problem A4

Sketch the curve $y^4 - x^4 - 96y^2 + 100x^2 = 0$.

Solution

The graph is in three parts. It is symmetrical on reflection in the x -axis and on reflection in the y -axis. The central part is a figure of 8 with crossing point at the origin and touching the lines $y/x = \pm\sqrt{(100/96)}$. The top and bottom of the figure of 8 are at $(0, \pm\sqrt{96})$. The right hand part is a sort of dimpled hyperbola with asymptotes $y/x = \pm 1$. It comes in along each asymptote and touches the line $x = 8$ at $y = \pm\sqrt{48}$, then it dimples back away from the origin to cut the x -axis at $x = 10$. The left hand part of the curve is just the reflection of this in the y -axis.

Problem A5

Show that a line in the plane with rational slope contains either no lattice points or an infinite number. Show that given any line L of rational slope we can find $\delta > 0$, such that no lattice point is a distance k from L where $0 < k < \delta$.

Solution

The first part is obvious. If the line passes through (m, n) and has slope a/b (where m, n, a, b are all integers), then

it also passes through $(m + kb, n + ka)$ for any integer a . Although not required, we note that there are lines with rational slope passing through no lattice points, for example, $y = 1/2$ and $y = x + 1/2$.

Suppose L is $bx - ay + c = 0$. The distance of L from (x, y) is $|bx - ay + c|/\sqrt{(a^2 + b^2)}$. If (x, y) is a lattice point then $bx - ay$ is an integer. Hence $|bx - ay + c|$ is either zero or at least 1 if c is an integer, or at least the distance of c from the nearest integer if c is not an integer. So we can find $k > 0$, so that $|bx - ay + c| > k$ for all lattice points not on L . Hence the distance $|bx - ay + c|/\sqrt{(a^2 + b^2)}$ is at least $k/\sqrt{(a^2 + b^2)}$ for all lattice points not on L . So we can take δ to be $k/\sqrt{(a^2 + b^2)}$.

Problem A6

Let C be a parabola. Take points P, Q on C such that (1) PQ is perpendicular to the tangent at P , (2) the area enclosed by the parabola and PQ is as small as possible. What is the position of the chord PQ ?

Solution

We may take the parabola as $y = x^2$ and P as (a, a^2) . The tangent at P has slope $2a$, so the normal has slope $-1/2a$ and equation $(y - a^2) + 1/2a(x - a) = 0$. This meets the parabola at $x^2 + x/2a - a^2 - 1/2 = 0$, so the other root is $x = -1/2a - a$ and Q is $(-1/2a - a, (a + 1/2a)^2)$. The area under the curve between P and Q is $1/3(a^3 + (a + 1/2a)^3)$. The area under the chord PQ is $(2a + 1/2a)(a^2 + (a + 1/2a)^2)/2 = 2a^3 + 3a/2 + 1/2a + 1/16a^3$. Hence the area enclosed is $4/3 a^3 + a + 1/4a + 1/48a^3$. We wish to minimise this. The gradient is $4a^2 + 1 - 1/4a^2 - 1/16a^4 = 1/16a^4(4a^2 + 1)^2(2a - 1)(2a + 1)$. Hence there is a minimum at $a = 1/2$ (and another at $a = -1/2$). In the minimum position P is $(1/2, 1/4)$ or $(-1/2, 1/4)$, so the minimum positions are the intersection of the parabola with the perpendicular to the axis through the focus $(0, 1/4)$.

Problem A7

Show that if $\sum a_n$ converges, then so does $\sum a_n/n$.

Solution

Put $s_n = a_1 + a_2 + \dots + a_n$. Then $s_n - s_{n-1} = a_n$, and so $a_1/1 + a_2/2 + a_3/3 + \dots + a_n/n = s_1(1/1 - 1/2) + s_2(1/2 - 1/3) + \dots + s_n(1/n - 1/n+1) + s_n/n+1$. Now $s_1(1/1 - 1/2) + s_2(1/2 - 1/3) + \dots + s_n(1/n - 1/n+1) + \dots$ is absolutely convergent, because the sequence $|s_n|$ is bounded, say by B . Hence $|s_1(1/1 - 1/2)| + |s_2(1/2 - 1/3)| + \dots + |s_n(1/n - 1/n+1)| \leq B(1/1 - 1/2) + (1/2 - 1/3) + \dots + (1/n - 1/n+1) = B(1 - 1/n+1) < B$. Obviously $s_n/n+1$ tends to zero, so $\sum a_n/n$ is absolutely convergent and hence convergent.

Problem B1

R is the reals. $f, g : R^2 \rightarrow R$ have continuous partial derivatives of all orders. What conditions must they satisfy for the differential equation $f(x, y) dx + g(x, y) dy = 0$ to have an integrating factor $h(xy)$?

Solution

If $h(xy)$ is an integrating factor, then $f(x, y) h(xy) dx + g(x, y) h(xy) dy$ is an exact form. In other words, $\partial/\partial y (f(x, y) h(xy)) = \partial/\partial x (g(x, y) h(xy))$. So $h \partial f/\partial y + x f h' = h \partial g/\partial x + y g h'$. Hence $h'/h = (\partial f/\partial y - \partial g/\partial x) / (yg - xf)$.

The lhs is a function of xy , so the rhs must be also. Equally if the rhs is a function of xy , then we can immediately integrate to get $h(xy)$. Thus the required necessary and sufficient condition for the existence of an integrating factor is that $(\partial f/\partial y - \partial g/\partial x) / (yg - xf)$ should be a function of xy .

Problem B2

R is the reals. Find an example of functions $f, g : R \rightarrow R$, which are differentiable, not identically zero, and satisfy $(f/g)' = f'/g'$.

Solution

Suppose we take $f = gh$. Then we require $h' = h + g/g' h'$. If we take $g/g' = 2$, then $h' = -h$. So this suggests $f(x) = e^{-x^2}$, $g(x) = e^{x^2}$. Checking, we see that that works.

Problem B3

Show that $\ln(1 + 1/x) > 1/(1 + x)$ for $x > 0$.

Solution

We have the well-known $e^y > 1 + y$ for all y not equal to 1. [Prove, for example, by differentiating.] So $\ln(1 + y) < y$. Put $y = -1/(1 + x)$, so $1 + y = x/(1 + x)$. Hence $1/(1 + x) < -\ln(x/(1+x)) = \ln(1 + 1/x)$.

Problem B4

Can we find four distinct concentric circles all touching an ellipse?

Solution

Answer: yes, we can find four such circles for any ellipse except the circle.

We show first that we can find four such circles for any non-circular ellipse. Let the ellipse have centre O , major axis AB (length $2a$) and minor axis CD (length $2b$). Take a point P with $OP < (a - b)/2$. Then using the triangle inequality we have $PA > OA - OP = (a + b)/2$, $PB > OB - OP = (a + b)/2$. $PC < OC + OP = (a + b)/2$, $PD < OD + OP = (a + b)/2$. Put $k = (a + b)/2$. Then, if Q is a point on the ellipse, $(PQ - k)$ is positive at A and B and negative at C and D . Hence there must be at least four points Q on the ellipse at which it is stationary and hence at which PQ is normal to the ellipse. There is a point A' near A which is a maximum and hence has $PA' > k$. Similarly, there is a point B' near B with $PB' > k$. There is a point C' near C with $PC' < k$ and a point D' near D with $PD' < k$. Let us assume that P is in the quadrant AOC and not on either axis. The reflection D'' of D' in the line AB lies at or near C' and hence has $PD'' \geq PC'$ (which is known to be a minimum). Put P lies closer to D'' than to D' . Hence $PC' < PD'$. A similar argument shows that $PB' > PA'$. So we have $PB' > PA' > k > PD' > PC'$. Thus the circles centre P through A' , B' , C' and D' are all distinct and all touch the ellipse.

Note that we cannot find 4 circles in the case when the ellipse is circular. For given a point P and a circle E centre O , a circle centre P can only touch E on the line OP , but that implies there can only be two such circles.

Problem B5

T is a torus, center O . The plane P contains O and touches T . Prove that $P \cup T$ is two circles.

Solution

We may take the torus to be $(\sqrt{(x^2 + y^2)} - a)^2 + z^2 = b^2$, where $a > b$. Viewed in the x - z plane the torus appears as two circles each radius b and centred at $x = \pm a, z = 0$. The plane P appears as the line through the origin tangent to both circles and hence having equation $cz = bx$, where $c = \sqrt{a^2 - b^2}$. P also contains the y -axis which is perpendicular to the line $z = kx$. Hence the equation of the plane P is $z = kx$.

It is far from obvious that P meets T in two circles. If we draw the plane P we see that T contains the four points on the y -axis, $y = \pm a \pm b$, and also the points a distance c either side of the origin. The obvious way to put these onto two circles is to take circles radius c , centred at $(x, y, z) = (0, \pm b, 0)$. These can be written as $(x, y, z) = (c \cos t, \pm b + a \sin t, b \cos t)$. Hence $\sqrt{(x^2 + y^2)} = a \pm b \sin t$. So $(\sqrt{(x^2 + y^2)} - a)^2 + z^2 = b^2 \sin^2 t + b^2 \cos^2 t = b^2$. Thus these circles do indeed lie on the torus.

We may write the equation of T as $2a\sqrt{(x^2 + y^2)} = x^2 + y^2 + z^2 + c^2$. Hence $4a^2(x^2 + y^2) = (x^2 + y^2 + z^2 + c^2)^2$. So $(x^2 + y^2 + z^2 - c^2)^2 = 4a^2x^2 + 4a^2y^2 - 4c^2(x^2 + y^2 + z^2) = 4b^2x^2 + 4b^2y^2 - 4c^2z^2$. Hence $(x^2 + y^2 + z^2 - c^2)^2 - (2by)^2 = 4b^2x^2 - 4b^2c^2$. So $(x^2 + (y + b)^2 + z^2 - a^2)(x^2 + (y - b)^2 + z^2 - a^2) = 4(bx - cz)(bx + cz)$.

So if a point lies on the plane P , which has equation $bx - cz = 0$ and on the torus T , then it also lies on one of the two spheres $(x^2 + (y + b)^2 + z^2 - a^2) = 0$ and $(x^2 + (y - b)^2 + z^2 - a^2) = 0$. But a plane always intersects a sphere in a circle (or not at all) so the intersection of P and T is at most two circles. We have already established that it contains two circles, so it cannot contain any other points.

Problem B6

The real polynomial $p(x) \equiv x^3 + ax^2 + bx + c$ has three real roots $\alpha < \beta < \gamma$. Show that $\sqrt{(a^2 - 3b)} < (\gamma - \alpha) \leq 2\sqrt{(a^2/3 - b)}$.

Solution

$a^2 - 3b = (\alpha + \beta + \gamma)^2 - (\alpha\beta + \beta\gamma + \gamma\alpha) = (\gamma - \alpha)^2 - (\gamma - \beta)(\beta - \alpha) < (\gamma - \alpha)^2$. Hence $\sqrt{(a^2 - 3b)} < (\gamma - \alpha)$.

Also $4(a^2 - 3b) - 3(\gamma - \alpha)^2 = (\gamma - \alpha)^2 - 4(\gamma - \beta)(\beta - \alpha) = ((\gamma - \beta) + (\beta - \alpha))^2 - 4(\gamma - \beta)(\beta - \alpha) = ((\gamma - \beta) - (\beta - \alpha))^2 \geq 0$. Hence $(\gamma - \alpha) \leq 2\sqrt{(a^2/3 - b)}$.

Problem B7

In 4-space let S be the 3-sphere radius r: $w^2 + x^2 + y^2 + z^2 = r^2$. What is the 3-dimensional volume of S? What is the 4-dimensional volume of its interior?

Solution

Answer: $2\pi^2 r^3$, $1/2 \pi^2 r^4$.

A hyperplane perpendicular to the x-axis will cut S in a sphere. If the hyperplane is at a distance k from the origin, then the radius of the sphere will be $\sqrt{(r^2 - k^2)}$. So the 4-dimensional volume $= \int_{-r}^r 4\pi/3 (r^2 - x^2)^{3/2} dx = 4\pi r^4/3 \int_{-1}^1 (1 - t^2)^{3/2} dt$.

Let $K = \int_{-1}^1 (1 - t^2)^{3/2} dt$. We find that $A = t(1 - t^2)^{3/2}$ differentiates to $K - 3t^2(1 - t^2)^{1/2}$. $B = t(1 - t^2)^{1/2}$ differentiates to $(1 - t^2)^{1/2} - t^2(1 - t^2)^{-1/2}$. $C = \sin^{-1}t$ differentiates to $(1 - t^2)^{-1/2}$. Hence $B + C$ differentiates to $2(1 - t^2)^{1/2}$, so $2A + 3(B + C)$ differentiates to $8K$. So $\int_{-1}^1 (1 - t^2)^{3/2} dt = (\text{terms involving powers of } (1 - t^2)^{1/2} + 3/8 \sin^{-1}t)|_{-1}^1 = 3\pi/8$. Hence the 4-dimensional volume is $\pi^2 r^4/2$.

The volume of the 3-dimensional surface is the derivative wrt r of the 4-dimensional volume (because the 4-dimensional volume for $r + \delta r$ is the 4-dimensional volume for r + the 3-dimensional volume of the bounding sphere times δr). Hence the 3-dimensional volume is $2\pi^2 r^3$.

12th Putnam 1952

Problem A1

$p(x)$ is a polynomial with integral coefficients. The leading coefficient, the constant term, and $p(1)$ are all odd. Show that $p(x)$ has no rational roots.

Solution

Suppose there is a rational root p/q . Let $p(x) = \sum_0^n a_r x^r$. Then $\sum a_r p^r q^{n-r} = 0$. So p divides a_0 and hence is odd, and q divides a_n and must also be odd. But now $\sum a_r p^r q^{n-r} = a_r = 1 \pmod{2}$. Contradiction.

Problem A2

Show that the solutions of the differential equation $(9 - x^2)(y')^2 = 9 - y^2$ are conics touching the sides of a square.

Solution

We have $dy/dx = \pm \sqrt{(1 - (y/3)^2)} / \sqrt{(1 - (x/3)^2)}$. Integrating, $\sin^{-1}y/3 = \pm \sin^{-1}x/3 + A$, where A is a constant. Hence $y/3 = \sin A \cos(\pm \sin^{-1}x/3) \pm x/3 \cos A = \sin A \sqrt{(1 - (x/3)^2)} \pm x/3 \cos A$. So $y = k(9 - x^2) \pm hx$, where $h^2 + k^2 = 1$. So $y^2 \pm 2hxy + x^2 = 9(1 - h^2)$. This is evidently a family of conics. Note that h is not restricted to being less than 1. By taking a suitable complex constant A we can get any value for h . But since negative values are possible, there is no benefit in retaining the \pm sign. So we can write the family of conics simply as $y^2 + 2hxy + x^2 = 9(1 - h^2)$.

We might guess that the square is $x = \pm 3, y = \pm 3$. If we substitute $x = 3$ in the equation for the conic, we get $y^2 + 6hy + 9 = 9 - 9h^2$, or $(y - 3h)^2 = 0$. This has a repeated root $y = 3h$, which shows that the line $x = 3$ touches the conic. Similarly for the other four sides of the square.

Problem A3

Let the roots of the cubic $p(x) \equiv x^3 + ax^2 + bx + c$ be α, β, γ . Find all (a, b, c) so that $p(\alpha^2) = p(\beta^2) = p(\gamma^2) = 0$.

Solution

Answer: $(0, 0, 0)$ roots $0, 0, 0$

$(-1, 0, 0)$ roots $0, 0, 1$

$(-2, 1, 0)$ roots $0, 1, 1$

$(0, -1, 0)$ roots $0, 1, -1$

$(1, 1, 0)$ roots $0, \omega, \omega^2$, where $\omega = e^{2\pi i/3}$

$(-3, 3, -1)$ roots $1, 1, 1$

$(-1, -1, 1)$ roots $1, 1, -1$

$(1, -1, -1)$ roots $1, -1, -1$

$(0, 0, -1)$, roots $1, \omega, \omega^2$

$(i, 1, -i)$, roots $1, -1, -i$

$(-i, 1, i)$, roots $1, -1, i$

$(1-\omega, 1-\omega, -\omega)$, roots ω, ω, ω^2

$(1-\omega^2, 1-\omega^2, -\omega^2)$, roots $\omega, \omega^2, \omega^2$

$(1+\omega, 1+\omega, \omega)$, roots $\omega, -\omega, \omega^2$

$(1+\omega^2, 1+\omega^2, \omega^2)$, roots $\omega, \omega^2, -\omega^2$

$((1-i\sqrt{7})/2, (-1-i\sqrt{7})/2, -1)$, roots k, k^2, k^4 , where $k = e^{2i\pi/7}$

$((1+i\sqrt{7})/2, (-1+i\sqrt{7})/2, -1)$, roots k^3, k^5, k^6 .

$p(x) = 0$ has only three roots, so the squares must all be found in $\{\alpha, \beta, \gamma\}$. Let α be the root with the largest modulus, then $|\alpha|^2 > |\alpha|$ unless $|\alpha| \leq 1$. Let γ be the root with the smallest non-zero modulus, then $|\gamma|^2 < |\beta|$ unless $|\gamma| \geq 1$. Hence all non-zero roots have modulus 1.

All roots zero is possible and gives $(a, b, c) = (0, 0, 0)$.

Suppose $\beta = \gamma = 0$ and α is non-zero. Then $\alpha^2 = \alpha$, so $\alpha = 1$. This gives $(a, b, c) = (-1, 0, 0)$.

Suppose $\gamma = 0$, and α, β are non-zero. If $\alpha^2 = \alpha$, then $\alpha = 1$. Then $\beta^2 = 1$ or β , so $\beta = 1$ or -1 . This gives $(a, b, c) = (-2, 1, 0)$ or $(0, -1, 0)$. If α^2 is not α (and β^2 not β), then we must have $\alpha^4 = \alpha$, so α and β are ω and ω^2 , where $\omega = e^{2i\pi/3}$. This gives $(a, b, c) = (1, 1, 1)$.

The remaining possibility is that all roots are non-zero (and have modulus 1). There might be 3, 2, 1 or 0 roots equal to 1. The first case gives $(-3, 3, -1)$. In the second case the third root must satisfy $\gamma^2 = 1$ or γ and hence must be -1 , giving $(a, b, c) = (-1, -1, 1)$.

We consider the third case (just one root equal to 1). If both the roots not equal to 1 have square 1, then they must both be -1 and $(a, b, c) = (1, -1, -1)$. If one has square 1 and one not. Then they must be -1 and $\pm i$. This gives $(a, b, c) = (i, 1, i)$ or $(-i, 1, -i)$. If neither have square 1, then we must have $\beta^2 = \gamma, \gamma^2 = \beta$, hence $\beta^3 = 1$ and β and γ are ω and ω^2 .

The final case is that no roots are 1. If we have $\alpha^2 = \beta$ and $\beta^2 = \alpha$, then α and β are ω and ω^2 and $\gamma^2 = \omega$ or ω^2 , so $\gamma = \pm\omega$ or $\pm\omega^2$. So the roots are ω, ω, ω^2 or $\omega, \omega^2, \omega^2$ or $\omega, -\omega, \omega^2$ or $\omega, \omega^2, -\omega^2$. This gives $(a, b, c) = (1-\omega, 1-\omega, \omega)$ or $(1-\omega^2, 1-\omega^2, \omega^2)$ or $(1+\omega, 1+\omega, \omega)$ or $(1+\omega^2, 1+\omega^2, \omega^2)$.

The final possibility is $\alpha^2 = \beta, \beta^2 = \gamma, \gamma^2 = \alpha$. Hence $\alpha^7 = 1$. If we put $k = e^{i2\pi/7}$, then the roots could be k, k^2, k^4 or k^6, k^5, k^3 . This corresponds to $(a, b, c) = ((1-i\sqrt{7})/2, (-1-i\sqrt{7})/2, -1), ((1+i\sqrt{7})/2, (-1+i\sqrt{7})/2, -1)$.

Problem A4

A map represents the polar cap from latitudes -45° to 90° . The pole (latitude 90°) is at the center of the map and lines of latitude on the globe are represented as concentric circles with radii proportional to $(90^\circ - \text{latitude})$. How are east-west distances exaggerated compared to north-south distances on the map at a latitude of -30° ?

Solution

Answer: too high by a factor $4\pi/\sqrt{27} = 2.42$.

Let the globe have radius R . Then the distance from the pole to the circle of latitude at -30 is $2\pi R/3$. The circumference of the circle is $2\pi R (\sqrt{3})/2$. The ratio of circumference to distance is $(\sqrt{27})/2$. On the map the corresponding ratio is 2π . Thus the map overstates by a factor $4\pi/\sqrt{27}$.

Problem A5

a_i are reals $\neq 1$. Let $b_n = 1 - a_n$. Show that $a_1 + a_2b_1 + a_3b_1b_2 + a_4b_1b_2b_3 + \dots + a_nb_1b_2 \dots b_{n-1} = 1 - b_1b_2 \dots b_n$.

Solution

Induction on n . Obvious for $n = 1$. Suppose true for n . Then $a_1 + a_2b_1 + a_3b_1b_2 + a_4b_1b_2b_3 + \dots + a_{n+1}b_1b_2 \dots b_n = 1 - b_1b_2 \dots b_n + a_{n+1}b_1b_2 \dots b_n = 1 - b_1b_2 \dots b_n(1 - a_{n+1}) = 1 - b_1 \dots b_{n+1}$, which is the result for $n+1$.

Problem A6

Prove that there are only finitely many cuboidal blocks with integer sides $a \times b \times c$, such that if the block is painted on the outside and then cut into unit cubes, exactly half the cubes have no face painted.

Solution

It is sufficient to show that $abc = 2(a-2)(b-2)(c-2)$ has only finitely many solutions in integers $c \leq b \leq a$. If $c \leq 4$, then $c \geq 2(c-2)$, but $ab > (a-2)(b-2)$, so there are no solutions. If $c \geq 10$, then $c/(c-2) \leq 10/8$, so $D \leq 125/64 < 2$, where for convenience we have written $abc/((a-2)(b-2)(c-2))$ as D , so there are no solutions. Hence $c = 5, 6, 7, 8, \text{ or } 9$.

If $c = 5$ and $b \geq 24$, then $D \leq 5/3 (24/22)^2 < 2$, so there are no solutions. Now $a/(a-2)$ is strictly monotonic decreasing, so there is at most one solution for given b, c . Hence there are only finitely many solutions for $c = 5$.

Similarly, for $c = 6$, we find that are only solutions for $b < 16$; for $c = 7$, we find $b < 14$; for $c = 8, b < 12$ and for $c = 9, b < 12$. Hence there are only finitely many solutions in each case.

Problem A7

Let O be the center of a circle C and P_0 a point on the circle. Take points P_n on the circle such that angle $P_nOP_{n-1} = +1$ for all integers n . Given that π is irrational, show that given any two distinct points P, Q on C , the (shorter) arc PQ contains a point P_n .

Solution

All the points must be distinct, for if we had $P_n = P_m$ for $n < m$, then the integer $m - n$ would be a multiple of 2π and hence π would be rational.

Choose N so that the arc length $PQ < 2\pi R/N$, where R is the radius of the circle. Divide the circle into N equal arcs length $2\pi R/N$. Then at least two of the points P_1, P_2, \dots, P_{N+1} must lie in the same arc. Suppose the points are P_n and P_{n+m} (with m, n positive), so that $P_n P_{n+m}$ is an arc of length less than $2\pi R/N$. Hence the arcs $P_{n+m} P_{n+2m}, P_{n+2m} P_{n+3m} \dots$ have the same length. So for some k , P_{n+km} will lie in the arc PQ .

Problem B1

ABC is a triangle with, as usual, $AB = c$, $CA = b$. Find necessary and sufficient conditions for $b^2 c^2 / (2bc \cos A) = b^2 + c^2 - 2bc \cos A$.

Solution

Answer: $BC = AB$ or CA .

We have that $a^2 = b^2 + c^2 - 2bc \cos A$. So the condition implies that $(b^2 + c^2 - a^2) / (2bc) = \cos A = bc / (2a^2)$. Hence, $b^2 c^2 = a^2 (b^2 + c^2 - a^2)$, so $(a^2 - b^2)(a^2 - c^2) = 0$. Hence $a = b$ or c .

Conversely, if $a = b$, then the altitude from C meets AB at its midpoint and so $\cos A = \cos B = c / (2a) = bc / (2a^2)$. Hence $a^2 = bc / (2 \cos A)$, so $(b^2 + c^2 - 2bc \cos A) = b^2 c^2 / (2bc \cos A)$.

Problem B2

Find the surface comprising the curves which satisfy $dx/(yz) = dy/(zx) = dz/(xy)$ and which meet the circle $x = 0$, $y^2 + z^2 = 1$.

Solution

We have $x dx = y dy = z dz$. Integrating, $y^2 = x^2 + h$, $z^2 = x^2 + k$ (*). We are told that some point of the circle $x = 0$, $y^2 + z^2 = 1$ belongs to the curve, so h and k must be non-negative with sum 1. Hence the curve (*) lies in the surface $2x^2 + 1 = y^2 + z^2$, which is a one-sheet hyperboloid with the x -axis as an axis of symmetry. However, not all points of this surface lie on a curve (*). Clearly we require $|y| \geq |x|$ and $|z| \geq |x|$. Equally, it is clear that this is a sufficient condition for we can then find h, k to satisfy (*).

Problem B3

Let $A(x) =$ be the matrix

$$\begin{matrix} 0 & a-x & b-x \\ -a-x & 0 & c-x \\ -b-x & -c-x & 0 \end{matrix}$$

For which (a, b, c) does $\det A(x) = 0$ have a repeated root in x ?

Solution

Multiplying out, we find that $\det A(x) = -2x^3 + 2(bc - ac + ab)x$. This has roots 0 and $\pm\sqrt{(ab + bc - ac)}$. This has a repeated root iff $ac = ab + bc$.

Problem B4

The solid S consists of a circular cylinder radius r , height h , with a hemispherical cap at one end. S is placed with the center of the cap on the table and the axis of the cylinder vertical. For some k , equilibrium is stable if $r/h > k$, unstable if $r/h < k$ and neutral if $r/h = k$. Find k and show that if $r/h = k$, then the body is in equilibrium if any point of the cap is in contact with the table.

Solution

Let O be the centre of the base of the hemispherical cap. As the solid rolls, O is always a height r above the table. Consider a point P on the axis a distance d from O and inside the cap. If the solid is rolled to a position where the axis is at an angle θ to the vertical, then P ends up a vertical distance $d \cos \theta$ below O , so its height has increased. Similarly, a point on the axis a distance d above O would end up a vertical distance $d \cos \theta$ above O , so its height

would decrease. So equilibrium will be stable, neutral or unstable according as the centre of mass is below, at or above O.

We start by finding the position of the centre of mass of the cap. Take the x-axis along the axis with the origin at O. Assume the density is ρ . The moment about an axis perpendicular to the x-axis is $\int_0^r \pi(r^2 - x^2)\rho x dx = \pi \rho r^4/4$. The centre of mass of the cylindrical portion of the solid is obviously a distance $h/2$ from O, so we require $(\pi \rho r^2 h) h/2 = \pi \rho r^4/4$ or $r/h = \sqrt{2}$ get the centre of mass at O. Thus $k = \sqrt{2}$.

If the centre of mass is at O, then the centre of mass is always at the same height above the table, so the solid is in equilibrium however far it is tilted from the vertical.

Problem B5

The sequence a_n is monotonic and $\sum a_n$ converges. Show that $\sum n(a_n - a_{n+1})$ converges.

Solution

The sum of the first n terms is $(a_1 - a_2) + 2(a_2 - a_3) + \dots + n(a_n - a_{n+1}) = a_1 + a_2 + a_3 + \dots + a_n - n a_{n+1}$. We are given that $\sum a_n$ converges, so it is sufficient to show that the sequence $n a_{n+1}$ converges to zero.

But since $\sum a_n$ converges, $|a_{n+1} + a_{n+2} + \dots + a_{2n}|$ is arbitrarily small for n sufficiently large. Since a_n is monotonic, this implies that $n a_{n+1}$ is arbitrarily small for n sufficiently large.

Problem B6

A, B, C are points of a fixed ellipse E. Show that the area of ABC is a maximum iff the centroid of ABC is at the center of E.

Solution

Take the ellipse as $x^2/a^2 + y^2/b^2 = 1$. Take the three points as A ($a \cos u, b \sin u$), B ($a \cos v, b \sin v$), C ($a \cos w, b \sin w$). If the area is a maximum then the tangent at A must be parallel to BC (otherwise we could keep B and C fixed and move A to increase the altitude and hence the area). So we have $-b/a \cot u = (b \sin v - b \sin w)/(a \cos v - a \cos w)$ and hence $\cos(u - v) = \cos(u - w)$. Similarly, $\cos(v - u) = \cos(v - w)$.

Hence $2u = v + w \pmod{2\pi}$. Similarly $2v = u + w \pmod{2\pi}$, so $3u = 3v \pmod{2\pi}$. Hence u, v, w are (in some order and for some k) $k, k + 2\pi/3, k - 2\pi/3$. Hence the centroid is at $x = a \cos k + a \cos(k+2\pi/3) + a \cos(k-2\pi/3) = 0$, $y = b \sin k + b \sin(k+2\pi/3) + b \sin(k-2\pi/3) = 0$.

Conversely, if the centroid of A, B, C is at the centre, then $\cos u + \cos v + \cos w = 0$, $\sin u + \sin v + \sin w = 0$. Hence $\cos(u - v) = \cos u (-\cos u - \cos w) + \sin u (-\sin u - \sin w) = -1 - \cos(u - w)$. Hence $\cos(u - v) + \cos(u - w) = -1$. Similarly, $\cos(v - u) + \cos(v - w) = -1$, hence $\cos(u - w) = \cos(v - w)$. So $\cos w (\cos u - \cos v) = \sin w (\sin v - \sin u)$, so $-b/a \cot w = (b \sin v - b \sin u)/(a \cos v - a \cos u)$, and hence the tangent at C is parallel to AB. Similarly, the tangent at A is parallel to BC and the tangent at B is parallel to AC. Hence the area is a maximum.

Problem B7

Let R be the reals. Define a_n by $a_1 = \alpha \in R$, $a_{n+1} = \cos a_n$. Show that a_n converges to a limit independent of α .

Solution

$|\cos x| \leq 1$, so $-1 \leq a_2 \leq 1$. Hence $0 < a_3 < 1$ and so $0 < a_n < 1$ for $n \geq 3$.

$\cos x$ is strictly monotonic decreasing over the range 0 to 1, so if $a_n < a_{n+1}$ then $a_{n+2} < a_{n+1}$. Now the gradient of $\cos x$ is greater than -1 throughout the interval (0, 1), so if $a_n < a_{n+1}$ then $\cos a_n - \cos a_{n+1} < a_{n+1} - a_n$. Hence $a_n < a_{n+2} < a_{n+1}$. Similarly, if $a_n > a_{n+1}$, then $a_n > a_{n+2} > a_{n+1}$. Thus either a_{2n} is an increasing sequence bounded above and a_{2n+1} is a decreasing sequence bounded below or vice versa. Hence both a_{2n} and a_{2n+1} converge.

By the mean value theorem $|a_{n+1} - a_{n+2}| = |\cos a_n - \cos a_{n+1}| = \sin k |a_n - a_{n+1}|$ for some k in (0, 1). But $\sin k < \sin 1 < 0.9$. So a_{2n} and a_{2n+1} must converge to the same limit. Suppose this limit is h . Then $h = \cos h$. But $\cos x$ is strictly decreasing in (0, 1) and x is strictly increasing, so they have a single point of intersection in (0, 1). Thus h must be the unique point in (0, 1) at which $h = \cos h$.

13th Putnam 1953

Problem A1

Show that $(2/3)n^{3/2} < \sum_{r=1}^n \sqrt{r} < (2/3)n^{3/2} + (1/2)\sqrt{n}$.

Solution

The gradient of \sqrt{x} is falling from $x=0$ to $x=n$, so $(2/3)n^{3/2} = \int_0^n \sqrt{x} dx < \sqrt{1} + \sqrt{2} + \sqrt{3} + \dots + \sqrt{n}$. That gives the first inequality.

To get the second, we note that the chords joining (r, \sqrt{r}) to $(r+1, \sqrt{r+1})$ all lie under the curve, so if we subtract the area of the little triangles from $\sum_{r=1}^n \sqrt{r}$ then we get something less than the integral. The triangles have area $1/2\sqrt{1} + 1/2(\sqrt{2} - \sqrt{1}) + 1/2(\sqrt{3} - \sqrt{2}) + \dots + 1/2(\sqrt{n} - \sqrt{n-1}) = 1/2\sqrt{n}$. That gives the second inequality.

Problem A2

The complete graph with 6 points and 15 edges has each edge colored red or blue. Show that we can find 3 points such that the 3 edges joining them are the same color.

Solution

Take any point A. It has 5 edges, so at least 3 of them must be the same color. wlog it is red. So we have B, C, D with AB, AC, AD all red. Now if any of the three edges BC, CD, DB is red, then that gives us a red triangle. But if they are all blue, then BCD is a blue triangle.

Problem A3

a, b, c are real, and the sum of any two is greater than the third. Show that $2(a+b+c)(a^2+b^2+c^2)/3 > a^3+b^3+c^3+abc$.

Solution

wlog we may take $a \geq b \geq c$. c (and hence also a and b) must be positive. For if $c \leq 0$, then $a \geq b+c$. Contradiction.

Multiplying out the required relation and cancelling, we must prove that $2(a^2b+a^2c+b^2a+b^2c+c^2a+c^2b) > a^3+b^3+c^3+3abc$. But we have $a^2(b+c) > a^3$, $b^2(c+a) > b^3$, $c^2(a+b) > c^3$. So it is sufficient to prove that $a^2b+a^2c+b^2a+b^2c+c^2a+c^2b > 3abc$. But $a^2b \geq abc$ and $ab^2 \geq abc$. Also $b^2c+bc^2 = bc(b+c) > abc$. Similarly, $ac(a+c) > abc$. So we have in fact proved the slightly stronger result with abc replaced by $4abc/3$.

Problem A4

Using $\sin x = 2 \sin x/2 \cos x/2$ or otherwise, find $\int_0^{\pi/2} \ln \sin x dx$.

Solution

Let $I = \int_0^{\pi/2} \ln \sin x dx$. Making the suggested substitution, we get $I = \pi/2 \ln 2 + \int_0^{\pi/2} \ln \sin x/2 \cos x/2 dx$. Putting $y = x/2$ we get $\int_0^{\pi/2} \ln \sin x/2 \cos x/2 dx = 2 \int_0^{\pi/4} \ln \sin y + \ln \cos y dy$. But $\cos y = \sin(\pi/2 - y)$, so $\int_0^{\pi/4} \ln \cos y dy = \int_{\pi/4}^{\pi/2} \ln \sin y dy$. Hence $I = \pi/2 \ln 2 + 2I$, so $I = -\pi/2 \ln 2$.

Problem A5

S is a parabola with focus F and axis L. Three distinct normals to S pass through P. Show that the sum of the angles which these make with L less the angle which PF makes with L is a multiple of π .

Solution

We start by finding a formula for $\tan(A+B+C+D)$. Applying $\tan(A+B) = (\tan A + \tan B)/(1 - \tan A \tan B)$ twice and writing $a = \tan A$, $b = \tan B$, $c = \tan C$, $d = \tan D$, we get $\tan(A+B+C+D) = (\sum a - \sum abc)/(1 - \sum ab + abcd)$.

Take the parabola as $y = x^2$. The gradient at (k, k^2) is $2k$, so the normal has slope $-1/2k$. The normal has equation $2k^3 + (1-2y)k - x = 0$. For given x, y this is a cubic in k, so has 1 or real 3 roots. Their slopes satisfy the cubic $4xh^3 + 2(1-2y)h^2 + 1 = 0$ (*).

The focus is at $(0, 1/4)$, so the line joining (x, y) has slope $(y - 1/4)/x$ and negative is $(1/4 - y)/x$. The quartic with this root and the 3 roots of (*) is $(h + (y - 1/4)/x)(4xh^3 + 2(1 - 2y)h^2 + 1)$ (**). So the angles the normals make with L less the angle which PF makes with L is a multiple of π iff $(\sum a = \sum abc)$, where a, b, c, d are the slopes of the

four lines and hence iff the coefficients of h^3 and h in (***) are the same. But the coefficient of h^3 is $2(1 - 2y) + 4x$ ($y - 1/4)/x = 1$ and the coefficient of h is 1.

Problem A6

Show that $\sqrt{7}, \sqrt{7 - \sqrt{7}}, \sqrt{7 - \sqrt{7 + \sqrt{7}}}, \sqrt{7 - \sqrt{7 + \sqrt{7 - \sqrt{7}}}}, \dots$ converges and find its limit.

Solution

Answer: 2.

Let x_n be the n th number in the sequence. We have $x_{n+2} = \sqrt{7 - \sqrt{7 + x_n}}$. Hence if $x_n < 2$, then $x_{n+2} > 2$, and if $x_n > 2$, then $x_{n+2} < 2$. So if the sequence converges, its limit must be 2.

If $x_n = 2 + \varepsilon$, with $0 < \varepsilon < 1$, then $9 + \varepsilon < 9 + \varepsilon + \varepsilon^2/36 = (3 + \varepsilon/6)^2$, so $x_{n+2} > \sqrt{7 - (3 + \varepsilon/6)} = \sqrt{4 - \varepsilon/6}$. But certainly $\varepsilon/6 < \varepsilon/3 - \varepsilon^2/144$, so $\sqrt{4 - \varepsilon/6} > 2 - \varepsilon/12$. Thus x_{n+2} differs from 2 by less than $\varepsilon/12$. Similarly, if $x_n = 2 - \varepsilon$, with $0 < \varepsilon < 1$, then $9 - \varepsilon > 9 - \varepsilon/2 + \varepsilon^2/144 = (3 - \varepsilon/12)^2$, so $x_{n+2} < \sqrt{4 + \varepsilon/12} < 2 + \varepsilon/48$.

So we have established that if $|x_n - 2| < 1$, then $|x_{n+2} - 2| < 1/12 |x_n - 2|$. But we certainly have $|x_1 - 2| < 1$, and $|x_2 - 2| < 1$, so x_n converges to 2.

Problem A7

$p(x) \equiv x^3 + ax^2 + bx + c$ has three positive real roots. Find a necessary and sufficient condition on a, b, c for the roots to be $\cos A, \cos B, \cos C$ for some triangle ABC .

Solution

If $A + B + C = 180^\circ$, then $\cos A = -\cos(B + C) = \sin B \sin C - \cos B \cos C$. Squaring, we get $\cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cos B \cos C = 1$. So a necessary condition is $a^2 - 2b - 2c = 1$.

Conversely, suppose that this condition holds. Then if the roots are p, q, r , we have $p^2 + q^2 + r^2 + 2pqr = 1$ (*). We are given that the roots are all positive, so $2pqr > 0$, hence $p^2 < 1$ and so $p < 1$. Similarly for q and r . So we can find angles A, B, C greater than 0 and less than 90° such that $p = \cos A, q = \cos B, r = \cos C$. Now we can rewrite (*) as $(1 - \cos^2 B)(1 - \cos^2 C) = \cos^2 A + 2 \cos A \cos B \cos C + \cos^2 B \cos^2 C = (\cos A + \cos B \cos C)^2$. But $(1 - \cos^2 B) = \sin^2 B, (1 - \cos^2 C) = \sin^2 C$, so we have $\sin B \sin C = \pm(\cos A + \cos B \cos C)$. But we know that A, B, C are between 0 and 90° , so $\cos A, \cos B, \cos C, \sin B, \sin C$ are all positive. Hence we must use the $+$ sign and we have $\cos A = \sin B \sin C - \cos B \cos C$, so $A + B + C = 180^\circ$. Hence the condition is also sufficient.

Problem B1

Does $\sum_{n=1}^{\infty} 1/n^{1+1/n}$ converge?

Solution

Answer: no.

$x < e^x$ for all $x \geq 0$, so $x^{1/x} < e < 3$. Hence $1/n^{1+1/n} > 1/3n$. But $\sum 1/n$ diverges.

Problem B2

$p(x)$ is a real polynomial of degree n such that $p(m)$ is integral for all integers m . Show that if k is a coefficient of $p(x)$, then $n! k$ is an integer.

Solution

Note that $n!$ is best possible, because $1/n! x(x+1) \dots (x+n-1)$ is always integral for integral x (it is the binomial coefficient $\binom{x+n-1}{n}$).

We need a standard result from the calculus of differences. Let $\Delta f(x) = f(x+1) - f(x)$. Then $p(x) = p(0) + \Delta_1 x + 1/2! \Delta_2 x(x-1) + 1/3! \Delta_3 x(x-1)(x-2) + \dots + 1/n! \Delta_n x(x-1) \dots (x-n+1)$ (*), where $\Delta_m = \Delta^m p(0)$ (thus $\Delta_1 = p(1) - p(0), \Delta_2 = p(2) - 2p(1) + p(0)$ etc).

Assume this is true. Then if $p(x)$ is integral for all integral x , all the Δ_m must be integral. So the result above gives $n! p(x) =$ an integral combination of integral polynomials. Hence all the coefficients of $n! p(x)$ are integral.

To prove the result, notice first that if $f(x) = x(x-1)(x-2) \dots (x-m+1)$, then $\Delta^r f(0) = m!$ for $r = m$ and 0 otherwise. So if we call the rhs of (*) $q(x)$, then $\Delta^m q(0)$ is a sum of terms which are all zero except for $\Delta^m(1/m!)$

$\Delta^m x(x-1)\dots(x-m+1)(0) = \Delta_m$. Hence $\Delta^m p(0) = \Delta^m q(0)$ for $m = 1, 2, \dots, n$. Also $p(0) = q(0)$, so by a simple induction $p(m) = q(m)$ for $m = 0, 1, 2, \dots, n$. But $q(x)$ is a polynomial of degree at most n . If polynomials of degree at most n agree at $n+1$ points, then they must be identical. Hence $p(x) = q(x)$.

Problem B3

k is real. Solve the differential equations $y' = z(y+z)^k$, $z' = y(y+z)^k$ subject to $y(0) = 1$, $z(0) = 0$.

Solution

Adding, $(y+z)' = (y+z)^{k+1}$. Integrating $(y+z)^k = 1/(1-kx)$ (1).

Multiplying opposite sides of the two given equations together, we get $yy'(y+z)^k = zz'(y+z)^k$. Hence $yy' = zz'$. Integrating $y^2 - z^2 = 1$. Hence $(y+z)^k(y-z)^k = 1$, so $(y-z)^k = 1-kx$ (2).

For k non-zero, we can immediately solve (1) and (2) to get $y = 1/2((1-kx)^{1/k} + 1/(1-kx)^{1/k})$, $z = 1/2((1-kx)^{1/k} - 1/(1-kx)^{1/k})$.

For $k = 0$, the original equations simplify to $y' = z$, $z' = y$. So $y'' = y$. So $y = A \cosh x + B \sinh x$, $z = A \sinh x + B \cosh x$. Applying the initial conditions, $y = \cosh x$, $z = \sinh x$.

Problem B4

R is the reals. S is a surface in R^3 containing the point $(1, 1, 1)$ such that the tangent plane at any point $P \in S$ cuts the axes at three points whose orthocenter is P . Find the equation of S .

Solution

Consider a plane cutting the axes at $\mathbf{a} = (a, 0, 0)$, $\mathbf{b} = (0, b, 0)$, $\mathbf{c} = (0, 0, c)$. If the orthocentre is at $\mathbf{p} = (x, y, z)$, then we have $(\mathbf{p} - \mathbf{a}) \cdot (\mathbf{b} - \mathbf{c}) = (\mathbf{p} - \mathbf{b}) \cdot (\mathbf{a} - \mathbf{c}) = 0$. But $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{c} = \mathbf{c} \cdot \mathbf{a} = 0$, so we have $\mathbf{p} \cdot (\mathbf{b} - \mathbf{c}) = \mathbf{p} \cdot (\mathbf{a} - \mathbf{c}) = 0$. In other words the line from the origin $(0, 0, 0)$ to (x, y, z) is normal to the plane. So the surface satisfies the condition that all its normals pass through the origin and it passes through $(1, 1, 1)$. This implies that it is the sphere $x^2 + y^2 + z^2 = 3$.

Note, however, that for points with one coordinate zero, the tangent plane will meet one axis at infinity, so we should arguable exclude all such points. That divides the sphere into 8 disconnected pieces. The piece containing $(1, 1, 1)$ is that in the positive octant ($x > 0, y > 0, z > 0$).

Problem B5

The coefficients of the complex polynomial $z^4 + az^3 + bz^2 + cz + d$ satisfy $a^2d = c^2 \neq 0$. Show that the ratio of two of the roots equals the ratio of the other two.

Solution

We start with a straight slog. Let the roots be p, q, r, s . We have $a^2d - c^2 = (p+q+r+s)^2 pqr - (pqr + pqs + prs + qrs)^2$. The terms like $2p^2q^2rs$ all cancel, leaving $p^3qrs + pq^3rs + pqr^3s + pqr^3s - p^2q^2r^2 - p^2q^2s^2 - p^2r^2s^2 - q^2r^2s^2$.

The trick now is to factorise this. We might suspect that $p^2q - r^2s$ is a factor. But in that case $pr - qs$ and $ps - qr$ would presumably also be factors. $(p^2q - r^2s)(pr - qs)(ps - qr)$ has degree 6, as required. It also has the correct number of terms (8). So we try multiplying it out and find that it is the same.

So $(p^2q - r^2s)(pr - qs)(ps - qr) = 0$. But that means that at least one factor must be zero, which gives the result. Note that the only reason for giving us that a^2d is non-zero, is because that implies that none of the roots are zero (their product d is non-zero) and so having got $p^2q = r^2s$, we can divide to get $p/r = s/q$.

Problem B6

A and B are equidistant from O . Given $k > OA$, find the point P in the plane OAB such that $OP = k$ and $PA + PB$ is a minimum.

Solution

Let C be the circle centre O radius k . Take A' on the ray OA such that $OA \cdot OA' = k^2$ and B' on the ray OB such that $OB \cdot OB' = k^2$. If $A'B'$ intersects the circle C , then the points of intersection give the positions where $PA + PB$ is a

minimum. If not then the nearest point of C to A'B' (which is on the perpendicular bisector of AB) gives the minimum.

For given P on C, the triangles OAP and OPA' are similar, so $PA = (AO/PO) PA'$. Similarly, $PB = (BO/PO) PB'$, so $PA + PB = (AO/k)(PA' + PB')$, so minimising $PA + PB$ is equivalent to minimising $PA' + PB'$. If A'B' intersects C, then clearly the points of intersection minimise. If not, let Q be the point on C closest to A'B'. Let L be the tangent to the circle at that point (so that L is parallel to A'B'). Then $QA + QB < RA + RB$ for other points R on L. Given another point P on C, take the perpendicular to L through P. If it intersects L at R, then $PA + PB > RA + RB > QA + QB$.

Problem B7

Show that we can express any irrational number $\alpha \in (0, 1)$ uniquely in the form $\sum_{1}^{\infty} (-1)^{n+1} 1/(a_1 a_2 \dots a_n)$, where a_i is a strictly monotonic increasing sequence of positive integers. Find a_1, a_2, a_3 for $\alpha = 1/\sqrt{2}$.

Solution

Answer: $a_1 = 1, a_2 = 3, a_3 = 8$.

Let s_n be the sum of the first n terms. The terms alternate in sign and decrease in absolute value, so the odd terms of the sequence s_n decrease and the even terms increase. Every odd term exceeds every even term, so the odds and the evens must each converge. But $s_n - s_{n+1} < 1/2^n$ which tends to zero, so they tend to a common limit.

Choose a_n as follows. Take a_1 to be the smallest integer whose inverse exceeds α . Having chosen a_{2n-1} , take a_{2n} to be the largest integer such that $s_{2n} < \alpha$. Having chosen a_{2n} , take a_{2n+1} to be the largest integer such that $s_{2n+1} > \alpha$.

We have to show that these choices are always possible, or, in other words, that they yield a strictly increasing sequence a_n . This depends on the relation $1/k - 1/k(k+1) = 1/(k+1)$ (*). For suppose we have just chosen a_{2n-1} . Then we know that $s_{2n-1} > \alpha$, but that if increased a_{2n-1} by 1, then s_{2n-1} would be $< \alpha$. Hence, using (*), taking $a_{2n} = a_{2n-1} + 1$ certainly gives $s_{2n} < \alpha$. On the other hand, if we take a_{2n} to be sufficiently large, then s_{2n} will be close to s_{2n-1} and hence exceed α (note that α is irrational so it cannot equal any s_m). So the choice of a_{2n} will exceed a_{2n-1} . A similar argument shows that a_{2n+1} exceeds a_{2n} .

So we have established that we can find a sequence a_n such that all the odd partial sums s_n exceed α and all the even partial sums are less than α . But we have also established that s_n tends to a limit, so that limit must be α . That establishes existence.

Suppose there is another expansion so that $\alpha = 1/a_1 - 1/a_1 a_2 + \dots = 1/b_1 - 1/b_1 b_2 + \dots$. As above, we have that $1/(a_1 + 1) < \alpha < 1/a_1$ and also $1/(b_1 + 1) < \alpha < 1/b_1$. But since a_1 and b_1 are both integers that implies that $a_1 = b_1$. Suppose now that we have established that $a_i = b_i$ for $i \leq n$. Then we have that $\beta = (-1)^n a_1 \dots a_n (\alpha - 1/a_1 + 1/a_1 a_2 - \dots + (-1)^n / a_1 \dots a_n) = 1/a_{n+1} - 1/a_{n+1} a_{n+2} + \dots$. But we also have $\beta = 1/b_{n+1} - 1/b_{n+1} b_{n+2} + \dots$. We now argue as before that β lies between $1/(a_{n+1} + 1)$ and $1/a_{n+1}$ and also between $1/(b_{n+1} + 1)$ and $1/b_{n+1}$. Hence $a_{n+1} = b_{n+1}$. That establishes uniqueness.

Finally, consider $\alpha = 1/\sqrt{2}$. We have $1/2 < 1/\sqrt{2} < 1$, so $a_1 = 1$. We must pick a_2 as the largest integer so that $1 - 1/a_2 < 1/\sqrt{2}$, or $a_2 < 2 + \sqrt{2} = 3.4$. So $a_2 = 3$. We must pick a_3 as the largest integer so that $1 - 1/3 + 1/3a_3 > 1/\sqrt{2}$ or $2 + 1/x > 3/\sqrt{2}$ or $x < 3\sqrt{2} + 4 = 8.2$. So $a_3 = 8$.

14th Putnam 1954

Problem A1

Let N be the set $\{1, 2, \dots, n\}$, where n is an odd integer. Let $f : N \times N \rightarrow N$ satisfy: (1) $f(r, s) = f(s, r)$ for all r, s ; (2) $\{f(r, s) : s \in N\} = N$ for each r . Show that $\{f(r, r) : r \in N\} = N$.

Solution

We have a square array of numbers $a_{ij} = f(i, j)$. Each member of N occurs just once in each row (by (2)), so it occurs n times in all. But the array is symmetric (by (1)), so each member occurs an even number of times off the main diagonal. Hence it must occur an odd number of times, and hence at least once, on the main diagonal. But the main diagonal only has n entries, so if each of n numbers occurs at least once, then each must occur exactly once.

Problem A2

Given any five points in the interior of a square side 1, show that two of the points are a distance apart less than $k = 1/\sqrt{2}$. Is this result true for a smaller k ?

Solution

Let the square be $ABCD$ and O its midpoint. Let the midpoints of AB, BC, CD, DA be P, Q, R, S respectively. Then $ABCD$ is the union of the 4 smaller squares: $APOS, BQOP, CROQ, DSOR$, each with diameter k . At least one of the smaller squares must contain two points. So the two points are a distance apart $\leq k$. However, they cannot be a distance k apart, because the only pairs of points in a square realizing the diameter distance are opposite corners and they are not in the interior of $ABCD$. Hence the two points are a distance $< k$. Take one point at O and the others on the main diagonals a distance ϵ from the corners. Then shortest distance between two points is $k - \epsilon$ which can be made arbitrarily close to k .

Problem A3

Let S be the set of all curves satisfying $y' + a(x)y = b(x)$, where $a(x)$ and $b(x)$ are never zero. Show that if $C \in S$, then the tangent at the point $x = k$ on C passes through a point P_k which is independent of C .

Solution

Let C_h be the curve with $y(k) = h$. Then the gradient of C_h at (k, h) is $b(k) - a(k)h$, so the tangent at (k, h) is $(y - h) = (b(k) - a(k)h)(x - k)$. This may be written as $(a(k)x - a(k)k - 1)h = (b(k)(x - k) - y)$. Evidently, this always passes through the point $(k + 1/a(k), b(k)/a(k))$ whatever the value of h .

Problem A4

A uniform rod length $2a$ is suspended in midair with one end resting against a smooth vertical wall at X and the other end attached by a string length $2b$ to a point on the wall above X . For what angles between the rod and the string is equilibrium possible?

Solution

Answer: always 0; additionally, if $b > a > b/2$, $\cos^{-1}((b^2 + 2a^2)/(3ab))$.

Let the other end of the rod be Y . Let the string be ZY (with Z on the wall) and M its midpoint. The line along which the rod's weight acts passes through the midpoint of the rod and hence also through M . So a necessary condition for equilibrium is that the normal force at X also passes through M , in other words that MX is normal to the wall.

By the cosine rule: $MX^2 = b^2 + 4a^2 - 4ab \cos \theta$ (where $\theta = \text{angle } XYZ$). Similarly, $XZ^2 = 4b^2 + 4a^2 - 8ab \cos \theta$. But $MX^2 + XZ^2 = b^2$, so solving, $\cos \theta = (b^2 + 2a^2)/(3ab)$ (*).

Hence this is also a sufficient condition for equilibrium, because the geometry is now fixed. So we can solve for the tension in terms of the weight (resolving vertically) and then for the normal force (resolving horizontally). Hence the three forces have vector sum zero and the rod is in equilibrium.

Returning to (*), we know that $0 < a < b$. Put $x = a/b$, then $\cos \theta = (1 + 2x^2)/(3x)$. But it is easy to see that for $(1 + 2x^2)/(3x) < 1$, we require $x > 1/2$.

Problem A5

R is the reals. $f : (0, 1) \rightarrow R$ satisfies $\lim_{x \rightarrow 0} f(x) = 0$, and $f(x) - f(x/2) = o(x)$ as $x \rightarrow 0$. Show that $f(x) = o(x)$ as $x \rightarrow 0$.

Solution

We wish to show that given $\delta > 0$, $|f(x)| < x \delta$ for sufficiently small x . Certainly we can find $\varepsilon > 0$, such that for $x < \varepsilon$, $|f(x/2^r) - f(x/2^{r+1})| < x/2^{r+2} \delta$. Summing, $|f(x)| < x \delta (1/4 + 1/8 + \dots) + |f(x/2^n)| = x \delta/2 + |f(x/2^n)|$. But since $f(x/2^n)$ tends to zero, for any given x we can take n sufficiently large that $|f(x/2^n)| < x \delta/2$ and hence $|f(x)| < x \delta$ as required.

Problem A6

The real sequence a_n satisfies $a_n = \sum_{k=1}^{\infty} a_k^2$. Show $\sum a_n$ does not converge unless all a_n are zero.

Solution

Clearly $a_n \geq 0$. If any $a_n = 0$, then all subsequent a_i must be zero, and, by a trivial induction, all previous a_i . So assume no $a_n = 0$.

Notice that $a_{n+1} = a_n^2 + a_n$. But we have just shown that $a_n^2 > 0$, so $a_{n+1} > a_n$.

If the sum converges, then we can take n sufficiently large that $a_{n+1} + a_{n+2} + a_{n+3} + \dots < 1$. Then $a_n = a_{n+1}^2 + a_{n+2}^2 + a_{n+3}^2 + \dots < a_n(a_{n+1} + a_{n+2} + a_{n+3} + \dots) < a_n$. Contradiction. So the sum does not converge.

Problem A7

Prove that the equation $m^2 + 3mn - 2n^2 = 122$ has no integral solutions.

Solution

If m, n is a solution, then $4m^2 + 12mn - 8n^2 = 488$, so $(2m + 3n)^2 - 17n^2 = 488$, so $(2m + 3n)^2 = 12 \pmod{17}$. But 12 is not a quadratic residue of 17 [check: $0^2, 1^2, 2^2, 3^2, 4^2, 5^2, 6^2, 7^2, 8^2 = 0, 1, 4, 9, 16, 8, 2, 15, 13 \pmod{17}$].

Problem B1

Show that for any positive integer r , we can find integers m, n such that $m^2 - n^2 = r^3$.

Solution

We notice that $m^2 - n^2 = (m + n)(m - n)$. This suggests taking $m + n = r^2$, $m - n = r$. This works: $m = r(r + 1)/2$, $n = r(r - 1)/2$.

Problem B2

Let $a_n = \sum_{i=1}^n (-1)^{i+1}/i$. Assume that $\lim_{n \rightarrow \infty} a_n = k$. Rearrange the terms by taking two positive terms, then one negative term, then another two positive terms, then another negative term and so on. Let b_n be the sum of the first n terms of the rearranged series. Assume that $\lim_{n \rightarrow \infty} b_n = h$. Show that $b_{3n} = a_{4n} + a_{2n}/2$, and hence that $h \neq k$.

Solution

If we simply write out $a_{4n} = (1 - 1/2 + 1/3 - 1/4 + \dots - 1/4n)$ and $a_{2n}/2 = (1/2 - 1/4 + 1/6 - 1/8 + \dots - 1/4n)$ and add term by term we find that $a_{4n} + a_{2n}/2 = b_{3n}$.

Taking the limit, we have immediately that $h = 3k/2$. But clearly $k > 0$, since if we group the terms in pairs, each pair is positive. Hence $h \neq k$.

Problem B3

Let S be a finite collection of closed intervals on the real line such that any two have a point in common. Prove that the intersection of all the intervals is non-empty.

Solution

Let $[A, B]$ be the interval with the largest left-hand endpoint, and let $[a, b]$ be the interval with the smallest right-hand endpoint. Then since $[A, B]$ and $[a, b]$ overlap, we must have $A \leq b$, so $[A, b]$ is non-empty.

Now given any interval $[x, y]$ in S , we have $x \leq A$ and $y \geq b$, so $[A, b] \subseteq [x, y]$.

Problem B4

Let F be a point, and L and D lines, in the plane. Show how to construct the point of intersection (if any) between L and the parabola with focus F and directrix D .

Solution

If L and D are parallel and a distance d apart, then take the circle center F radius d . If it meets L , then the point(s) of intersection are the required points. If it does not, then there are none.

So assume L and D meet at O. If F lies on D, then the (degenerate) parabola is the line through F perpendicular to D. If it meets L, then the point of intersection is the required point. So assume F does not lie on D.

If L and D are perpendicular, then there is just one point. Take any point P (except O) on L and let the circle center P radius PO cut the line OF again at Q. Let the line through F parallel to PQ cut L at G. Then G is the required point (because an expansion center O takes the circle center P radius PO into a circle center G radius GO = GF).

So assume L and D are neither perpendicular nor parallel and meet at O. Take D' through O making the same angle with L as D. D and D' divide the plane into 4 sectors. If F lies in one of the two sectors containing L, then the required points exist. There are two required points unless F lies on D', when there is just one. [Any circle with center on L, touching D, must lie entirely within these two sectors, and hence F must also if the points exist.]

Take any point P on L (except O) and draw a circle center P touching D. Let the circle meet the line OF at Q. Let the line through F parallel to PQ cut L at G. Then G is the required point. If F does not lie on D', then there will be two possibilities for Q and hence two possibilities for G.

Problem B5

Let R be the reals. Let $f: (-1, 1) \rightarrow R$ be a function with a derivative at 0. Let a_n be a sequence in $(-1, 0)$ tending to zero and b_n a sequence in $(0, 1)$ tending to zero. Show that $\lim_{n \rightarrow \infty} (f(b_n) - f(a_n))/(b_n - a_n) = f'(0)$.

Solution

From the definition of derivative, we have that $(f(b_n) - f(0))/b_n \rightarrow f'(0)$, and $(f(0) - f(a_n))/-a_n \rightarrow f'(0)$. So $f(b_n) - f(0)$ lies within $\varepsilon/2$ of $b_n f'(0)$ for sufficiently large n, and $f(0) - f(a_n)$ lies within $\varepsilon/2$ of $-a_n f'(0)$ for sufficiently large n. Hence, adding, $f(b_n) - f(a_n)$ lies within ε of $(b_n - a_n)f'(0)$ for sufficiently large n, which is what we need.

Problem B6

If x is a positive rational, show that we can find distinct positive integers a_1, a_2, \dots, a_n such that $x = \sum 1/a_i$.

Solution

Rather surprisingly, the simplest possible algorithm - the greedy algorithm - works: take a_n to be the smallest integer not already chosen so that the sum does not exceed x. This terminates after a finite number of steps.

Note that the numbers involved can be quite large even in simple cases. For example: $4 = 1/1 + 1/2 + 1/3 + \dots + 1/30 + 1/200 + 1/77706 + 1/16532869712 + 1/3230579689970657935732 + 1/36802906522516375115639735990520502954652700$

The key idea is that the numerator of the difference gets smaller each time and hence the process must terminate.

First, we have to get the difference under 1. So take m such that $1 + 1/2 + \dots + 1/m \leq x < 1 + 1/2 + \dots + 1/(m+1)$. This is possible since the harmonic series diverges. If we have equality, then we are home. If not then the difference $x - (1 + 1/2 + \dots + 1/m) < 1/(m+1)$.

Let the difference $x - (1/a_1 + 1/a_2 + \dots + 1/a_n)$ be $d_n = r_n/s_n$ (where r_n and s_n are integers).

So we may assume that we have picked a_1, \dots, a_n and that $d_n < 1/(a_n + 1)$. We now take a_{n+1} so that $1/a_{n+1} \leq d_n < 1/(a_{n+1} - 1)$. Now $d_{n+1} = d_n - 1/a_{n+1} = (r_n a_{n+1} - s_n)/(s_n a_{n+1})$. This expression may not be in lowest terms, but it is certainly *sufficient* to show that $(r_n a_{n+1} - s_n) < r_n$ or $d_n a_{n+1} - 1 < d_n$. But that is true since $d_n(a_{n+1} - 1) < 1$. Finally, we notice that $d_{n+1} < 1/(a_{n+1} + 1)$ is equivalent to $d_n - 1/a_{n+1} < 1/(a_{n+1} + 1)$, which is true since $1/(a_{n+1} - 1) - 1/a_{n+1} < 1/(a_{n+1} + 1)$, since $a_{n+1} > 1$.

Problem B7

Let α be a positive real. Let $a_n = \sum_{i=1}^n (\alpha/n + i/n)^n$. Show that $\lim a_n \in (e^\alpha, e^{\alpha+1})$.

Solution

The largest term is $(1 + \alpha/n)^n$ which tends to e^α . The next largest is $(1 + (\alpha - 1)/n)^n$ which tends to $e^{\alpha-1}$ and so on. All terms are positive, so the limit is at least $e^\alpha + e^{\alpha-1} > e^\alpha$. Also the limit is at most $e^\alpha(1 + 1/e + 1/e^2 + \dots) < e^\alpha(1 + 1/2 + 1/4 + \dots) = 2e^\alpha < e^{\alpha+1}$.

15th Putnam 1955

Problem A1

Prove that if a, b, c are integers and $a\sqrt{2} + b\sqrt{3} + c = 0$, then $a = b = c = 0$.

Solution

$\sqrt{3}$ is irrational. For if not, let $\sqrt{3} = r/s$ with r and s having no common factor. But now $3s^2 = r^2$, so 3 divides r^2 and hence r . Hence 3 divides s^2 and hence s . So r and s have a common factor 3. Contradiction. Similarly $\sqrt{2}$ is irrational.

If $a\sqrt{2} + b\sqrt{3} + c = 0$, then $2a^2 - 3b^2 - c^2 = 2bc\sqrt{3}$. But $\sqrt{3}$ is irrational, so b or $c = 0$. If $b = 0$, then $a\sqrt{2} + c = 0$. But $\sqrt{2}$ is irrational, so $a = c = 0$ also.

Hence $c = 0$. So $a\sqrt{2} + b\sqrt{3} = 0$. We may take a and b to be relatively prime (divide out any common factor). Squaring: $2a^2 = 3b^2$.

Problem A2

O is the center of a regular n -gon $P_1P_2 \dots P_n$ and X is a point outside the n -gon on the line OP_1 . Show that $XP_1XP_2 \dots XP_n + OP_1^n = OX^n$.

Solution

Evidently it is sufficient to prove the result for the case $OP_1 = 1$. So represent P_k by the complex number ω^k , where $\omega = e^{i2\pi/n}$ an n th root of unity. Represent X by the real number r . Then we have to show that $|r - 1||r - \omega| \dots |r - \omega^{n-1}| = r^n - 1$, but that follows immediately because $r^n - 1 = (r - 1)(r - \omega) \dots (r - \omega^{n-1})$.

Problem A3

a_n is a sequence of monotonically decreasing positive terms such that $\sum a_n$ converges. S is the set of all $\sum b_n$, where b_n is a subsequence of a_n . Show that S is an interval iff $a_{n-1} \leq \sum_{i=n}^{\infty} a_i$ for all n .

Solution

The condition is certainly necessary. For suppose $a_{n-1} = k_2 > k_3 = \sum_{i=n}^{\infty} a_i$. Let $k_1 = \sum_{i=1}^{n-2} a_i$. Then k_1 and $k_1 + k_2$ belong to S , but no number in the non-empty interval $(k_1 + k_3, k_1 + k_2)$ belongs to S , so S cannot be an interval.

Now assume that the condition holds. Let $k = \sum a_n$. We show that $S = [0, k]$. We get the endpoints by taking the subsequence to be the empty set or the whole sequence. So take $h \in (0, k)$. Define b_n to be the earliest member of the sequence not so far chosen such that $\sum_{i=1}^n b_i \leq h$. This is clearly possible since $a_m \rightarrow 0$. If at any point we get equality we are home.

So assume that the resulting $\{b_n\}$ is infinite. Clearly $\sum b_n \leq h$. If $\{b_n\}$ is missing infinitely many members of $\{a_n\}$, then given any $\varepsilon > 0$, we can find $a_m < \varepsilon$ missing from the subsequence. But that means that for some n , we rejected a_m because $b_1 + b_2 + \dots + b_n + a_m > h$. So $\sum b_i \geq b_1 + b_2 + \dots + b_n > h - \varepsilon$. Hence $\sum b_i = h$.

It remains to consider the case where only finitely many members are missing from the subsequence. Let a_m be the largest such. Then for some n we have that $b_1 + b_2 + \dots + b_n + a_m > h$. We also have that $b_1 + b_2 + \dots + b_n + \sum_{i=m+1}^{\infty} a_i \leq h$. But $a_m \leq \sum_{i=m+1}^{\infty} a_i$. So we have a contradiction and this case cannot occur.

Problem A4

n vertices are taken on a circle and all possible chords are drawn. No three chords are concurrent (except at a vertex). How many points of intersection are there (excluding vertices)?

Solution

Answer: nC_4 .

Every 4 vertices correspond to a unique point of intersection (the intersection of the two diagonals defined by the 4 points).

[Actually, I got this the hard way. Take one of the vertices. Consider the diagonals from it. The diagonal to the next vertex but one has one vertex on one side and $(n - 3)$ on the other. So it has $1(n - 3)$ points of intersection on it. The next diagonal has $2(n - 4)$ and so on up to $(n - 3)1$. We repeat for each vertex. That gives a total of $n[1(n - 3) + 2(n - 4) + \dots + (n - 3)1]$ points of intersection. But we count each diagonal twice and then each point of intersection twice. So the number of points of intersection is $n/4 \sum_{i=1}^{n-3} r(n - 2 - r) = n/4 [(n - 2) 1/2 (n - 3)(n - 2) - 1/6 (n - 3)(n - 2)(2n - 5)] = 1/24 n(n - 2)(n - 3) [3(n - 2) - (2n - 5)] = nC4$.

Of course, at this point, one wonders why the answer has such a nice form!]

Problem A5

Given a parabola, construct the focus (with ruler and compass).

Solution

You need to know two facts:

- (1) Rays from infinity come to a focus at the focus. In other words, if a line L parallel to the axis of the parabola meets it at X and the focus is F , then L and XF are equally inclined to the normal at X ;
- (2) The line joining the midpoints of two parallel chords is parallel to the axis.

The construction is then fairly obvious. Take two parallel chords. Join their midpoints and extend to meet the parabola at X . Take a line M through X parallel to the chords. Then M is a tangent. The line perpendicular to M through X is the normal. Hence find the line XF (although not yet F). Repeat for another pair of parallel chords. The two lines intersect at F .

Problem A6

For what positive integers n does the polynomial $p(x) \equiv x^n + (2 + x)^n + (2 - x)^n$ have a rational root.

Solution

Answer: $n = 1$.

There are no roots at all if n is even. If $n = 1$, then -4 is the root. So suppose n is odd and at least 3.

We can use a similar argument to that showing that $\sqrt{2}$ is irrational. Suppose $x = r/s$ with r and s having no common factor is a root. Then $r^n + (2s + r)^n + (2s - r)^n = 0$. The last two terms expand and add to give terms with even coefficients, so r must be even. So set $r = 2t$ and we have $t^n + (s + t)^n + (s - t)^n = 0$. The same argument shows that t must be even. But now we have $t^n + 2(s^n + nC2 s^{n-2} t^2 + \dots) = 0$. Since $n \geq 3$, s must be even also. Contradiction. So there are no rational roots for n odd ≥ 3 .

Problem A7

k is a real constant. y satisfies $y'' = (x^3 + kx)y$ with initial conditions $y = 1, y' = 0$ at $x = 0$. Show that the solutions of $y = 0$ are bounded above but not below.

Solution

For some $x_0, x^3 + kx > 0$ for all $x > x_0$. [For example, if $k \geq 0$, then we may take $x_0 = 0$. If $k < 0$, then we may take $x_0 = -k$.] Now suppose that $a < b$ are two consecutive zeros greater than x_0 . If $y(x) > 0$ on (a, b) , then y'' must be negative for at least part of the interval (a, b) , but that is impossible, since $(x^3 + kx)y > 0$ for the whole interval. Similarly, if $y(x) < 0$ on (a, b) , then y'' must be positive for at least part of the interval (a, b) , but that is impossible, since $(x^3 + kx)y < 0$ for the whole interval. So there is at most one root greater than x_0 and that provides a bound. [For completeness, we note that we require y not to be identically zero on a non-zero interval, but that is not possible because $y(0)$ is non-zero.]

Suppose the second part is false. Then we can find x_2 such that for any $x < x_2, y(x)$ is non-zero. There are two cases. Suppose first that $y(x) > 0$ for all $x < x_2$. We can take $x_3 < x_2$ such that $x^3 + kx < -2$ for all $x < x_3$. Hence, in particular, $y''(x) < 0$ for $x < x_3$. Take arbitrary $a < b < x_3$, then as usual we can find $\xi \in (a, b)$ such that $y(a) = y(b) - (b - a)y'(b) + 1/2(b - a)^2 y''(\xi) < y(b) - (b - a)y'(b)$. If $y'(b) > 0$, then for $(b - a)$ sufficiently large $y(a) < 0$ (contradiction). So $y'(b) \leq 0$. So $y(a) \leq y(b) + 1/2(b - a)^2 y''(\xi)$. But now we use the stronger assumption that $\xi^3 + k\xi < -2$ and hence $y(a) < y(b)(1 - (b - a)^2)$, which becomes negative for sufficiently large $(b - a)$. Contradiction.

The remaining case is $y(x) < 0$ and $y''(x) > -2y(x)$ for all $x < x_3$. In particular, $y''(x) > 0$ for $x < x_3$. So arguing as before, $y'(b) \leq 0$. As before $y(a) \geq -y(b) \left(\frac{b-a}{a} - 1 \right)$, which becomes positive for sufficiently large $(b-a)$. Contradiction.

Problem B1

The lines L and M are horizontal and intersect at O . A sphere rolls along supported by L and M . What is the locus of its center?

Solution

Let the angle between the two lines be 2θ . Evidently the center rolls above the angle bisector. Let its projection onto the angle bisector have length x and let it be a height y above the angle bisector. Suppose the sphere has radius r . Then $(x \sin \theta)^2 + y^2 = r^2$. So the locus is an ellipse with semi-axes r and $r/\sin \theta$.

If we assume the rolling is under gravity, then the sphere falls off when it reaches $x = r/\sin \theta$, so the locus is just the upper half of the ellipse. If we assume the sphere is somehow kept in contact with the lines, then it can roll back underneath and so the locus is the whole ellipse. That is the ambiguity.

The trap is that it can also roll the other way, over the other angle bisector, giving another ellipse (or upper half) with semi-axes r and $r/\cos \theta$.

Problem B2

Let \mathbb{R} be the reals. $f: \mathbb{R} \rightarrow \mathbb{R}$ is twice differentiable, f'' is continuous and $f(0) = 0$. Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) = f(x)/x$ for $x \neq 0$, $g(0) = f'(0)$. Show that g is differentiable and that g' is continuous.

Solution

The only issue is $x = 0$, because elsewhere we have simply $g'(x) = f'(x)/x - f(x)/x^2$.

Given x , we can find ξ such that $f(x) = f(0) + x f'(0) + 1/2 x^2 f''(\xi) = x f'(0) + 1/2 x^2 f''(\xi)$. Hence $\lim_{x \rightarrow 0} (g(x) - f'(0))/x = \lim_{x \rightarrow 0} 1/2 f''(\xi) = 1/2 f''(0)$. So $g'(0)$ exists.

Now $\lim_{x \rightarrow 0} g'(x) = \lim_{x \rightarrow 0} (x f'(x) - f(x))/x^2$. But $f'(x) = f'(0) + x f''(\zeta)$, and $f(x) = x f'(0) + 1/2 x^2 f''(\xi)$ with $\xi, \zeta \in (0, x)$. So $\lim_{x \rightarrow 0} g'(x) = \lim_{x \rightarrow 0} f''(\zeta) - 1/2 f''(\xi) = 1/2 f''(0) = g'(0)$. So g' is continuous at 0.

Problem B3

Let S be a spherical cap with distance taken along great circles. Show that we cannot find a distance preserving map from S to the plane.

Solution

Let O be the center of the cap and C its perimeter. Then C is a circle center O . Its image must also be a circle with the same radius R , since the distance between each point of C and O is preserved. The circumference of the circle is also preserved. But the circumference is not equal to $2\pi R$.

Problem B4

Can we find n such that $\mu(n) = \mu(n+1) = \dots = \mu(n+1000000) = 0$? [The Möbius function $\mu(r) = 0$ iff r has a square factor > 1 .]

Solution

Answer: yes.

Let p_n be the n th prime. We show how to find a_n such that p_1^2 divides a_n , p_2^2 divides $a_n + 1$, p_3^2 divides $a_n + 2$, ..., p_n^2 divides $a_n + n - 1$.

Evidently we can take $a_1 = 4$. We now use induction. $a_n + k p_1^2 \dots p_n^2$ has the same property as a_n . So we need to pick k such that $a_n + k p_1^2 \dots p_n^2 \equiv 0 \pmod{p_{n+1}^2}$. But that is always possible (by Euclid's algorithm, for example) since p_{n+1}^2 is relatively prime to $p_1^2 \dots p_n^2$.

Problem B5

n is a positive integer. An infinite sequence of 0s and 1s is such that it only contains n different blocks of n consecutive terms. Show that it is eventually periodic.

Solution

We use induction on n . It is obvious for $n = 1$. Suppose it is true for n . Let S be a sequence with just $n + 1$ different $(n+1)$ -blocks. Consider the n -blocks formed by taking the first n elements of each $(n+1)$ -block. These evidently form all the possible n -blocks in S . So if there are at most n of them, then we are home by induction.

But if there are $n + 1$ of them (all distinct), then the first n elements of each $(n+1)$ -block determine the last element. So an n -block determines the entire sequence from that point on (because having fixed the $(n + 1)$ th element, we move the n -block along one place and hence fix the $(n + 2)$ nd element and so on). But there are only finitely many possible n -blocks, so there are two the same (in S). That gives a period, since the elements r places after the start of each block must be the same.

Note that the result is the best possible in the sense that we can construct a non-periodic sequence with just $n + 1$ different n -blocks. Take, for example, the n -blocks to be those with no or just one 1. Then if we take 1 followed by $n + 1$ zeros, followed by a 1, followed by $n + 2$ zeros, followed by a 1, followed by $n + 3$ zeros and so on, we have a non-periodic sequence involving only those $n + 1$ n -blocks.

Problem B6

Let \mathbb{N} be the set of positive integers and \mathbb{R}^+ the positive reals. $f : \mathbb{N} \rightarrow \mathbb{R}^+$ satisfies $f(n) \rightarrow 0$ as $n \rightarrow \infty$. Show that there are only finitely many solutions to $f(a) + f(b) + f(c) = 1$.

Solution

We show first that there are only finitely many solutions to $f(a) + f(b) = k (> 0)$. Take a to be $< b$. Since $f(n) \rightarrow 0$ we can find N such that $f(n) < k/2$ for $n > N$. So a must be $\leq N$. Now for each such a_0 , we can find M such that $f(n) < k - f(a_0)$ for $n \geq M$, so there are only finitely many b such that $f(a_0) + f(b) = k$. Hence there are only finitely many solutions to $f(a) + f(b) = k$.

Now consider $f(a) + f(b) + f(c) = 1$. We can find N such that $f(n) < 1/3$ for $n > N$. So at least one of a, b, c must be $\leq N$. But for each such a we then have only finitely many solutions to $f(b) + f(c) = 1 - f(a)$. Hence there are only finitely many solutions in total.

Problem B7

A three-dimensional solid acted on by four constant forces is in equilibrium. No two lines of force are in the same plane. Show that the four lines of force are rulings on a hyperboloid.

Solution

Let the lines be L_1, L_2, L_3, L_4 . If the line L intersects each of L_1, L_2, L_3 , then the moment of the first three forces about L is zero. Hence the moment of the fourth force about L is also zero and so L must also intersect L_4 . This is sufficient to prove that L_1, L_2, L_3, L_4 lie in a ruled quadric.

In fact the union of all the lines which intersect L_1, L_2, L_3 is a ruled quadric. Let P be a point on L_4 . For a point X and a line L not containing it we take XL to be the unique plane containing X and L . If L and L' are skew lines and X does not lie on either, then $XL \cap XL'$ is the unique line through X which meets L and L' . So $PL_1 \cap PL_2$ is the unique line through P which meets L_1 and L_2 . Suppose it meets L_1 at Q . Now $QL_2 \cap QL_3$ meets L_1, L_2 and L_3 so it must also meet L_4 . But there is a unique line through Q meeting L_2 and L_4 , so $QL_2 \cap QL_3 = PL_1 \cap PL_2$. Hence P lies on the line $QL_2 \cap QL_3$ which is a line of the quadric. P was arbitrary, so we have shown that L_4 lies in the quadric.

16th Putnam 1956

Problem A1

$\alpha \neq 1$ is a positive real. Find $\lim_{x \rightarrow \infty} ((\alpha^x - 1)/(\alpha x - x))^{1/x}$.

Solution

$\lim_{x \rightarrow \infty} (1/x)^{1/x} = \lim_{y \rightarrow 0} y^y = 1$. If $\alpha > 1$, then $\lim 1/(\alpha - 1)^{1/x} = 1$, and $\lim (\alpha^x - 1)^{1/x} = \lim (\alpha^x)^{1/x} = \alpha$, so the whole expression tends to α . If $\alpha < 1$, then $(\alpha^x - 1)/(\alpha - 1)$ tends to $1/(1 - \alpha)$, so the whole expression tends to 1.

Problem A2

Given any positive integer n , show that we can find a positive integer m such that mn uses all ten digits when written in the usual base 10.

Solution

Let n have d digits. Take $N = 12345678900 \dots 0$, ending in $d+1$ digits. One of $N+1, N+2, \dots, N+n$ must be a multiple of n and its first 10 digits are all different.

Problem A3

Find the trajectory of a particle which moves from rest in a vertical plane under (constant) gravity and a force kv perpendicular to its velocity v .

Solution

Take the x -axis horizontal and the y -axis vertically downwards. Then the equations of motion are: $y'' = g - kv'$, $x'' = ky'$. If $k = 0$, then we have the usual constant acceleration downwards ($x = 0, y = 1/2 gt^2$). So suppose $k \neq 0$.

By inspection, $x = A(\cos kt - 1) + B(\sin kt - kt)$ (we have chosen the terms $-A$ and $-kbt$ to give $x(0) = x'(0) = 0$). Hence (integrating $x'' = ky'$) we have $y = -A \sin kt + B(\cos kt - 1)$. But $y'(0) = 0$, so $A = 0$. Finally, to satisfy $y'' = g - kv'$, we $B = -g/k^2$. So $x = gt/k - g/k^2 \sin kt, y = g/k^2 (1 - \cos kt)$. This is the equation of a cycloid. Note that the particle is never able to fall more than a finite distance, it keeps being returned to its starting level.

Problem A4

Let $p(x)$ be a real polynomial of degree n with leading coefficient 1 and all roots real. Let R be the reals and $f: [a, b] \rightarrow R$ be an n times differentiable function with at least $n + 1$ distinct zeros. Show that $p(D)f(x)$ has at least one zero on $[a, b]$, where D denotes d/dx .

Solution

The basic tool is Rolle's theorem which tells us that there is a zero of $f'(x)$ between any two zeros of $f(x)$. But we would like a zero of $(D - \alpha)f$ between any two zeros of f . For that, notice that the zeros of $f(x)$ are also zeros of $e^{-\alpha x} f(x)$. So if $f(x)$ has $n + 1$ zeros, then the derivative of $e^{-\alpha x} f(x)$ has at least n zeros. But the derivative of $e^{-\alpha x} f(x)$ is $e^{-\alpha x}$ times $(D - \alpha)f(x)$. $e^{-\alpha x}$ is never zero, so $(D - \alpha)f(x)$ has at least n zeros.

That (more than) proves the result for $p(x)$ of degree 1. We now use induction on n , the degree of $p(x)$. Suppose it is true for n . Let $p(x)$ be a real polynomial with real roots and degree $n + 1$. Then $p(x) = q(x)(x - \alpha)$ for some α and $q(x)$ of degree n . Let f be an $n+1$ times differentiable real function on $[a, b]$ with at least $n + 2$ distinct zeros. Then we proved above that $(D - \alpha)f(x)$ has at least $n + 1$ distinct zeros. It is also at least n times differentiable. So, by the inductive hypothesis, $q(D)(D - \alpha)f(x)$ has at least one zero, which proves the result is true for $n + 1$.

Problem A5

Show that there are just $(n-k+1)C_k$ subsets of $\{1, 2, \dots, n\}$ with k elements and not containing both i and $i+1$ for any i .

Solution

There is a bijection between subsets of $\{1, 2, \dots, n\}$ with k elements not containing both i and $i+1$ for any i and subsets of $\{1, 2, \dots, n - k + 1\}$ with k elements. Namely, associate $r_1 < r_2 < \dots < r_k$ and $r_1, r_2 - 1, r_3 - 2, \dots, r_k - k + 1$. But there are obviously just $(n-k+1)C_k$ such subsets of $\{1, 2, \dots, n - k + 1\}$.

Problem A6

Let \mathbb{R} be the reals. Find $f : \mathbb{R} \rightarrow \mathbb{R}$ which preserves all rational distances but not all distances. Show that if $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ preserves all rational distances then it preserves all distances.

Solution

Let $f(x) = x$ for x rational, $x + 1$ for x irrational. If $x - y$ is rational, then either both x and y are rational in which case $f(x) - f(y) = x - y$, or neither are in which case $f(x) - f(y) = (x + 1) - (y + 1) = x - y$. So f preserves rational distances. But f does not preserve all distances. For example $f(\sqrt{2}) - f(0) \neq \sqrt{2} - 0$.

Given points A and B in the plane a distance k apart, and any $\varepsilon > 0$, we can find a point C such that AC and BC are rational and $|AC - k| < \varepsilon/2$, $|BC| < \varepsilon/2$. Let $A' = f(A)$, $B' = f(B)$, $C' = f(C)$. Then $|A'C'| = |AC|$, $|B'C'| = |BC|$, and hence $|A'B'|$ lies within ε of k . This is true for all $\varepsilon > 0$, so $|A'B'| = k$ and f preserves all distances.

Problem A7

Show that for any given positive integer n , the number of odd nC_m with $0 \leq m \leq n$ is a power of 2.

Solution

$(2n)C_{(2m+1)} = 2n/(2m+1) (2n-1)C_{2m}$, which is even. There are n even factors in the numerator of $2n C_{2m}$ and n even factors in its denominator. If we cancel 2 from each of these even factors, then we get nC_m . So $2n C_{2m}$ has the same parity as nC_m . Hence there are the same number of odd binomial coefficients for $2n$ as for n .

$(2n+1)C_{2m} = 2n C_{(2m-1)} + 2n C_{2m}$, so $(2n+1)C_{2m}$ has the same parity as $2n C_{2m}$. Similarly, $(2n+1)C_{(2m+1)} = 2n C_{2m} + 2n C_{(2m+1)}$, so $(2n+1)C_{(2m+1)}$ has the same parity as $2n C_{2m}$. Hence there are twice as many odd binomial coefficients for $2n+1$ as for $2n$.

The required result now follows by a trivial induction.

Problem B1

The differential equation $a(x, y) dx + b(x, y) dy = 0$ is homogeneous and exact (meaning that $a(x, y)$ and $b(x, y)$ are homogeneous polynomials of the same degree and that $\partial a/\partial y = \partial b/\partial x$). Show that the solution $y = y(x)$ satisfies $x a(x, y) + y b(x, y) = c$, for some constant c .

Solution

$$d(x a + y b) = (a dx + b dy) + x(\partial a/\partial x dx + \partial a/\partial y dy) + y(\partial b/\partial x dx + \partial b/\partial y dy).$$

Using exactness, this becomes: $(a dx + b dy) + x(\partial a/\partial x dx + \partial b/\partial x dy) + y(\partial a/\partial y dx + \partial b/\partial y dy)$. Homogeneity implies that for some integer k we have $x \partial a/\partial x + y \partial a/\partial y = k a$, and $x \partial b/\partial x + y \partial b/\partial y = k b$. So we get finally $d(x a + y b) = (a dx + b dy) + k(a dx + b dy)$. So if y is a solution, then $d(x a + y b) = 0$ and hence $x a + y b = c$ for some constant c .

Problem B2

Let \mathcal{P} be the set of all subsets of the plane. $f : \mathcal{P} \rightarrow \mathcal{P}$ satisfies $f(X \cup Y) \supseteq f(f(X)) \cup f(Y) \cup Y$ for all $X, Y \in \mathcal{P}$ (*). Show that (1) $f(X) \supseteq X$, (2) $f(f(X)) = f(X)$, (3) if $X \supseteq Y$, then $f(X) \supseteq f(Y)$, for all $X, Y \in \mathcal{P}$. Show conversely that if $f : \mathcal{P} \rightarrow \mathcal{P}$ satisfies (1), (2), (3), then f satisfies (*).

Solution

Taking $Y = X$ in (*) gives immediately that $f(X) = f(X \cup X) \supseteq f(f(X)) \cup f(X) \cup X \supseteq X$, which proves (1).

It also shows that $f(X) \supseteq f(f(X))$. Now put $Y = f(X)$ in (*). Then $X \cup Y = f(X)$ (by (1)), so (*) gives $f(f(X)) \supseteq Y = f(X)$. So we have proved (2).

Finally, if $X \supseteq Y$, then $X \cup Y = X$, so $f(X) = f(X \cup Y) \supseteq f(Y)$, which proves (3).

Now assume (1), (2) and (3) and take any X, Y . Then $X \cup Y \supseteq Y$, so $f(X \cup Y) \supseteq f(Y)$ (using (3)). But $f(Y) \supseteq Y$ (using (1)), so also $f(X \cup Y) \supseteq Y$. Similarly, $f(X \cup Y) \supseteq f(X)$. But $f(X) = f(f(X))$ (using (2)), so $f(X \cup Y) \supseteq f(f(X))$, which establishes (*).

Problem B3

ABCD is an arbitrary tetrahedron. The inscribed sphere touches ABC at S, ABD at R, ACD at Q and BCD at P. Show that the four sets of angles $\{\text{ASB}, \text{BSC}, \text{CSA}\}$, $\{\text{ARB}, \text{BRD}, \text{DRA}\}$, $\{\text{AQC}, \text{CQD}, \text{DQA}\}$, $\{\text{BPC}, \text{CPD}, \text{DPB}\}$ are the same.

Solution

The key is to notice that the two angles in the sets subtended by the same side of the tetrahedron are the same. For example $\angle \text{ASC} = \angle \text{AQC}$. Let O be the centre of the sphere and r its radius. Then $\text{AQ}^2 + r^2 = \text{AO}^2 = \text{AS}^2 + r^2$, so $\text{AQ} = \text{AS}$. Similarly, $\text{CQ} = \text{CS}$. So the triangles AQC, ASC are similar.

Now the sum of the angles in each set is the same. But $\angle \text{ASB}$ in the first set equals $\angle \text{ARB}$ in the second set, so $\angle \text{BSC} + \angle \text{CSA} = \angle \text{BRD} + \angle \text{DRA}$. Similarly, $\angle \text{CQD}$ in the third set equals $\angle \text{CPD}$ in the fourth, so $\angle \text{AQC} + \angle \text{DQA} = \angle \text{BPC} + \angle \text{DPB}$. Adding these two equations and using $\angle \text{BSC} = \angle \text{BPC}$, $\angle \text{CSA} = \angle \text{AQC}$, $\angle \text{BRD} = \angle \text{DPB}$, $\angle \text{DRA} = \angle \text{DQA}$ gives $\angle \text{CSA} = \angle \text{BRD}$. In other words, the angles in the set subtended by opposite sides of the tetrahedron are also the same.

That gives us all we need: $\angle \text{ASC} = \angle \text{BRD} = \angle \text{AQC} = \angle \text{BPD}$; $\angle \text{ASB} = \angle \text{ARB} = \angle \text{CQD} = \angle \text{CPD}$; $\angle \text{BSC} = \angle \text{ARD} = \angle \text{AQD} = \angle \text{BPC}$.

Problem B4

Show that for any triangle ABC, we have $\sin A \cos C + A \cos B > 0$.

Solution

The result is obviously true unless B or C is obtuse. If C is obtuse, then $|\cos C| < \cos B$ (because $|\cos C| = \cos(A + B) < \cos B$). But $\sin A < A$, so the result follows. So assume that B is obtuse.

We have A acute and hence $A < \tan A$. So $A \cos A < \sin A$. But C is acute, so $\cos C$ is positive and $A \cos A \cos C < \sin A \cos C$. Now $A \sin A \sin C$ is positive, so $A \cos(A + C) = A \cos A \cos C - A \sin A \sin C < A \cos A \cos C < \sin A \cos C$. But $\cos(A + C) = -\cos B$, so $-A \cos B < \sin A \cos C$.

Problem B5

Show that a graph with $2n$ points and $n^2 + 1$ edges necessarily contains a 3-cycle, but that we can find a graph with $2n$ points and n^2 edges without a 3-cycle.

Solution

Induction. For $n = 2$, the result is obviously true, because there is only one graph with 4 points and 5 edges and it certainly contains a triangle. Suppose the result is true for n . Consider a graph G with $2n + 2$ points and $n^2 + 2n + 2$ edges. Take any two points P and Q joined by an edge. We now consider two cases. If there are less than $2n + 1$ other edges to P and Q , then if we remove P and Q , we get a graph with $2n$ points and at least $n^2 + 1$ edges, which must contain a triangle, so G does also. If there are at least $2n + 1$ other edges then at least one point must be joined to both P and Q , but that gives a triangle.

Let G and H be disjoint sets each with n points. Join every point of G to every point of H by an edge. Then the resulting graph has n^2 edges, but it does not contain any triangles, because two points of any triangle must belong to G or to H and in either case there is no edge connecting them.

Problem B6

The sequence a_n is defined by $a_1 = 2$, $a_{n+1} = a_n^2 - a_n + 1$. Show that any pair of values in the sequence are relatively prime and that $\sum 1/a_n = 1$.

Solution

We show by induction on k that $a_{n+k} \equiv 1 \pmod{a_n}$. Obviously true for $k = 1$. Suppose it is true for k . Then for some m , $a_{n+k} = m a_n + 1$. Hence $a_{n+k+1} = a_{n+k}(m a_n + 1) + 1 = a_{n+k} m a_n + 1 \equiv 1 \pmod{a_n}$. So the result is true for all k . Hence any pair of distinct a_n are relatively prime.

We show by induction that $\sum_1^n 1/a_r = 1 - 1/(a_{n+1} - 1)$. For $n = 2$, this reduces to $1/2 = 1 - 1/(3 - 1)$, which is true. Suppose it is true for n . Then $\sum_1^{n+1} 1/a_r = 1 - k$, where $k = 1/(a_{n+1} - 1) - 1/a_{n+1} = 1/(a_{n+1}^2 - a_{n+1}) = 1/(a_{n+2} - 1)$, so it is true for $n + 1$. But $a_n \rightarrow \infty$, so $\sum 1/a_n = 1$.

Problem B7

$p(z)$ and $q(z)$ are complex polynomials with the same set of roots (but possibly different multiplicities). $p(z) + 1$ and $q(z) + 1$ also have the same set of roots. Show that $p(z) \equiv q(z)$.

Solution

This is much easier than it looks. We just have to consider $p - q$ and $p' - q'$. Suppose that p has A roots, of which B are distinct, and that $p(z) + 1$ has A roots, of which C are distinct. Without loss of generality we may assume the degree of q is at most A , so that $p - q$ has at most A roots. Clearly the roots of $p(z)$ and $p(z) + 1$ do not overlap. The B distinct roots of $p(z)$ and the C distinct roots of $p(z) + 1$ must all be roots of $p - q$. So $B + C \leq A$. On the other hand, p' has at least $A - B$ roots in common with p and at least $A - C$ in common with $p(z) + 1$, so it has at least $2A - (B + C)$ in total. But its degree is $A - 1$, so $(B + C) \geq A + 1$. Contradiction.

17th Putnam 1957

Problem A1

A surface S in 3-space is such that every normal intersects a fixed line L . Show that we can find a surface of revolution containing S .

Solution

Unfortunately, this is false.

It is not hard to paste together pieces of different surfaces of revolution. A simple example is as follows. Take a cylinder terminated at one end by a circle C . For part of the circumference of C continue the cylinder. Leave a gap either side and then on the rest join a portion of a cone (so that the surface is angled outwards away from the central axis).

It is possible, but hard, certainly too hard for a qu. 1, to prove that S is locally a surface of revolution. For details see Gleason et al.

Problem A2

k is a real number greater than 1. A uniform wire consists of the curve $y = e^x$ between $x = 0$ and $x = k$, and the horizontal line $y = e^k$ between $x = k - 1$ and $x = k$. The wire is suspended from $(k - 1, e^k)$ and a horizontal force applied at the other end, $(0, 1)$ to keep it in equilibrium. Show that the force is directed towards increasing x .

Solution

If d is the density per unit length, then the total mass of wire is $d(1 + \int_0^k \sqrt{1 + e^{2x}} dx)$. The moment about the y -axis is $d(k - 1/2) + d \int_0^k x \sqrt{1 + e^{2x}} dx$. We require that the x -coordinate of the centre of mass exceeds $k - 1$. In other words, $k - 1/2 + \int_0^k x \sqrt{1 + e^{2x}} dx > (k - 1) (1 + \int_0^k \sqrt{1 + e^{2x}} dx)$.

Set $f(k) = \int_0^k x \sqrt{1 + e^{2x}} dx - (k - 1) \int_0^k \sqrt{1 + e^{2x}} dx$. Then we require $f(k) > -1/2$ for all $k > 1$. Clearly $f(0) = 0$. We try differentiating, and get $f'(k) = \sqrt{1 + e^{2k}} - \int_0^k \sqrt{1 + e^{2x}} dx$.

One possibility is to evaluate the integral explicitly (substitute $y = e^{2x}$, then $z^2 = 1 + y$, we end up with a messy but doable integral). But it is maybe easier to approximate. We have $1 + e^{2x} < 1 + e^{2x} + e^{-2x}/4 = (e^x + e^{-x}/2)^2$. So $\int_0^k \sqrt{1 + e^{2x}} dx < \int_0^k (e^x + e^{-x}/2) dx = (e^x - e^{-x}/2) \Big|_0^k = e^k - e^{-k}/2 - 1/2 < e^k < \sqrt{1 + e^{2k}}$. So $f'(k) > 0$. Hence $f(k) > f(0) > 0 > -1/2$.

Problem A3

A and B are real numbers such that $\cos A \neq \cos B$. Show that for any integer $n > 1$, $|\cos nA \cos B - \cos A \cos nB| < (n^2 - 1) |\cos A - \cos B|$.

Solution

It is not clear how to use induction, expanding $\cos(n+1)A = \cos nA \cos A - \sin nA \sin A$ gives a profusion of sines and it is not clear how to get rid of them. So some other approach is needed.

An ingenious manipulation puts the expression entirely in terms of sines. Put $x = (A + B)/2$, $y = (A - B)/2$. Then $A = x + y$, $B = x - y$. Now $2 \cos nA \cos B = \cos(nA + B) + \cos(nA - B)$. Similarly, $2 \cos A \cos nB = \cos(nB + A) + \cos(nB - A)$. Switch to x and y and then go back to products of cosines. We have $\cos(nA + B) - \cos(nB + A) = \cos((n+1)x + (n-1)y) - \cos((n+1)x - (n-1)y) = -2 \sin(n+1)x \sin(n-1)y$. Similarly, $\cos(nA - B) - \cos(nB - A) = -2 \sin(n-1)x \sin(n+1)y$. Hence $|\cos nA \cos B - \cos A \cos nB| = |\sin(n+1)x \sin(n-1)y + \sin(n-1)x \sin(n+1)y|$. Obviously $|\cos A - \cos B| = |2 \sin x \sin y|$, so we have to prove that $|\sin(n+1)x \sin(n-1)y + \sin(n-1)x \sin(n+1)y| < 2(n^2 - 1) |\sin x \sin y|$.

It would evidently be sufficient to show that $|\sin mx| < m |\sin x|$. Unfortunately, that is not true. We have equality if $\sin x = 0$ or if $m = 1$. However, we can easily prove by induction that we have strict inequality in all other cases.

For $m = 2$, we have $\sin 2x = 2 \sin x \cos x$, which establishes the result since $|\cos x| < 1$ for $\sin x$ non-zero. Suppose it is true for m . Then $\sin(m+1)x = \sin mx \cos x + \sin x \cos mx$, so $|\sin(m+1)x| \leq |\sin mx| |\cos x| + |\sin x| |\cos mx| \leq |\sin mx| + |\sin x| < (m+1) |\sin x|$, so it is true for $m+1$.

Now if $\sin x = 0$, then $A + B$ is a multiple of 2π , so $\cos A = \cos B$. Similarly if $\sin y = 0$, then $A - B$ is a multiple of 2π , so $\cos A = \cos B$. But we are told that $\cos A \neq \cos B$, so $\sin x$ and $\sin y$ are both non-zero. Also we are given

that $n > 1$, so we have strict inequality on $|\sin(n+1)x| > (n+1) |\sin x|$ and $|\sin(n+1)y| > (n+1) |\sin y|$, whilst $|\sin(n-1)x| \geq (n-1) |\sin x|$ and $|\sin(n-1)y| \geq (n-1) |\sin y|$. Thus we get the required strict inequality.

Problem A4

$p(z)$ is a polynomial of degree n with complex coefficients. Its roots (in the complex plane) can be covered by a disk radius r . Show that for any complex k , the roots of $n p(z) - k p'(z)$ can be covered by a disk radius $r + |k|$.

Solution

Let the roots of $p(z)$ be a_1, a_2, \dots, a_n . Suppose they all lie in the disk centre c , radius r . Then $|c - a_n| \leq r$. Suppose that $|c - w| > r + |k|$. We show that w is not a root of $n p(z) - k p'(z)$. We have $|w - a_i| \geq |w - c| - |c - a_i| > r + |k| - r = |k|$.

Now $p'(z)/p(z) = \sum 1/(z - a_i)$ (note that this is still true if we have repeated roots), so $|p'(w)/p(w)| < n/|k|$ and hence $|k p'(w)/p(w)| < n$. So $|n - k p'(w)/p(w)| > 0$. But $|p(w)| > 0$ (since w lies outside the disk containing all the roots of $p(z)$), so $|n p(w) - k p'(w)| = |p(w)| |n - k p'(w)/p(w)| > 0$.

Problem A5

Let S be a set of n points in the plane such that the greatest distance between two points of S is 1. Show that at most n pairs of points of S are a distance 1 apart.

Solution

Induction on n . Obviously true for $n \leq 3$. Suppose it is true for n . Take $n+1$ points. If no point is a distance 1 from more than 2 points, then we are done. So assume that A, B and C are all a distance 1 from P . wlog the largest of the three angles APB, APC, BPC is APB . It must be at most 60° , since $AB \leq 1$. So the ray PC lies between the rays PA and PB .

Now suppose there is another point D (apart from P) such that $CD = 1$. Then CD must intersect PA , because otherwise one of CP, CA, DP and DA would exceed 1. Similarly, it must intersect PB . But that is impossible. So there is no such point D . Hence if we remove C we lose only one realisation of the distance 1. But the remaining n points have at most n realisations, so the result is established.

Problem A6

Define a_n by $a_1 = \ln \alpha$, $a_2 = \ln(\alpha - a_1)$, $a_{n+1} = a_n + \ln(\alpha - a_n)$. Show that $\lim_{n \rightarrow \infty} a_n = \alpha - 1$.

Solution

Unfortunately, this does not quite work as stated. If α is too small then $a_2 > \alpha$ and hence a_3 is undefined. The limit is just under 0.3442. So assume that α is sufficiently large that $a_2 < \alpha$.

For all non-zero x we have $e^x > 1 + x$. Hence for all positive x not equal to 1 we have $\ln x < x - 1$.

Suppose that $a_2 = \alpha - 1$. Then $a_n = \alpha - 1$ for all $n > 2$, which gives the result. So assume a_2 is not equal to $\alpha - 1$. Then for all $n \geq 2$ we have $\ln(\alpha - a_n) < \alpha - a_n - 1$, so $a_{n+1} = a_n + \ln(\alpha - a_n) < \alpha - 1$.

Also for $n \geq 3$, we have that $\alpha - a_n > 1$, so $\ln(\alpha - a_n) > 0$, so $a_{n+1} > a_n$. Thus for $n \geq 3$, a_n is a monotonic increasing sequence bounded above by $\alpha - 1$. So it must converge.

But $a_{n+1} - a_n = \ln(\alpha - a_n)$, so $\ln(\alpha - a_n)$ tends to 0 and hence a_n tends to $\alpha - 1$.

Problem A7

Show that we can find a set of disjoint circles such that given any rational point on the x -axis, there is a circle touching the x -axis at that point. Show that we cannot find such a set for the irrational points.

Solution

Two disjoint circles touching the x -axis at A and B and each with radius r cannot have $AB < 2r$. Now suppose a circle radius $R \geq r$ touches at A and a disjoint circle radius r touches at B . The circle radius r touching at A does not extend outside the circle radius R , so the circle at B must also be disjoint from it. Hence we still have $AB \geq 2r$. Hence there can only be countably many disjoint circles radius r or more touching the x -axis. But a countable set of countable sets is still countable, so there can only be countably many disjoint circles touching the x -axis (any such circle has radius $> 1/n$ for some n). There are uncountably many irrational points, so we cannot have disjoint circles touching at all the irrational points.

For the rational points take the circle touching at m/n (in lowest terms) to have radius $1/(3n^2)$. Now suppose the circles at m/n and a/b have centres P and Q . We have $PQ^2 = (m/n - a/b)^2 + (1/(3n^2) - 1/(3b^2))^2$. If they intersect, then $PQ \leq (1/(3n^2) + 1/(3b^2))$ and hence $(m/n - a/b)^2 \leq (1/(3n^2) + 1/(3b^2))^2 - (1/(3n^2) - 1/(3b^2))^2 = 4/9 \cdot 1/(n^2b^2)$, so $(mb - an)^2 < 4/9$. But $(mb - an)$ must be integral, so $mb - an = 0$ and hence $m/n = a/b$. So distinct circles do not intersect.

Problem B1

Let A be the 100×100 matrix with $a_{mn} = mn$. Show that the absolute value of each of the $100!$ products in the expansion of $\det A$ is congruent to $1 \pmod{101}$.

Solution

Each product is $100! \cdot 100!$. But 101 is prime, so the numbers $1, 2, \dots, 100$ can be divided into pairs with the product of each pair being $1 \pmod{101}$.

Problem B2

The sequence a_n is defined by its initial value a_1 , and $a_{n+1} = a_n(2 - k a_n)$. For what real a_1 does the sequence converge to $1/k$?

Solution

Answer: For a_1 strictly between 0 and $2/k$.

Suppose a_n has the opposite sign to k . Then $2 - k a_n$ is positive, and so a_{n+1} also has the opposite sign to k , so we cannot get convergence. Similarly, if $a_n = 0$, then $a_{n+1} = 0$ and we cannot get convergence. So it is a necessary condition that a_1 should be non-zero and have the same sign as k .

If k is positive and $a_1 > 2/k$, then a_2 is negative and so the sequence does not converge to $1/k$ (as above). Similarly, if k is negative and $a_1 < 2/k$, then a_2 is positive and so we do not have convergence. Thus it is a necessary condition for a_1 to lie strictly between 0 and $2/k$.

Suppose $a_n = 1/k + h$. Then $a_{n+1} = (1/k + h)(2 - (1 + kh)) = 1/k - kh^2$ (*). This is all we need. But to spell it out, consider first k positive. If a_1 lies strictly between 0 and $1/k$, then h lies strictly between $-1/k$ and $1/k$. Now (*) shows that for $n \geq 2$, $a_n \leq 1/k$ and is monotonic increasing. Hence it tends to a positive limit. If this limit is L , then referring to the original equation, $L = L(2 - kL)$, so $L = 1/k$. Similarly for k negative.

Problem B3

\mathbb{R}^+ is the positive reals, $f: [0, 1] \rightarrow \mathbb{R}^+$ is monotonic decreasing. Show that $\int_0^1 f(x) dx \int_0^1 x f(x)^2 dx \leq \int_0^1 x f(x) dx \int_0^1 f(x)^2 dx$.

Solution

$(f(x) - f(y))(y - x) \geq 0$. Also $f(x), f(y) \geq 0$, so $\int \int f(x) f(y) (f(x) - f(y))(y - x) dx dy \geq 0$. But $f(x) f(y) (f(x) - f(y))(y - x) = (f(x)^2 y f(y) - f(x) y f(y)^2) + (f(y)^2 x f(x) - f(y) x f(x)^2)$. So if we take the same limits of integration, 0 and 1 , for both integrals, then we have $\int \int f(x) f(y) (f(x) - f(y))(y - x) dx dy = 2 \int f(x)^2 y f(y) - f(x) y f(y)^2 dx dy = 2 \int f(x)^2 dx \int y f(y) dy - 2 \int f(x) dx \int y f(y)^2 dy$.

Problem B4

Show that the number of ways of representing n as an ordered sum of 1 s and 2 s equals the number of ways of representing $n + 2$ as an ordered sum of integers > 1 . For example: $4 = 1 + 1 + 1 + 1 = 2 + 2 = 2 + 1 + 1 = 1 + 2 + 1 = 1 + 1 + 2$ (5 ways) and $6 = 4 + 2 = 2 + 4 = 3 + 3 = 2 + 2 + 2$ (5 ways).

Solution

We can establish a bijection as follows. Given a representation of n as an ordered sum of 1 s and 2 s, proceed as follows. Add a final 2 at the end. Now group together all summands up to and including the first 2 , then all following summands up to and including the next 2 , and so on. Add the members in each group. This gives us an ordered set of integers each at least 2 and summing to $n+2$.

Conversely, given a representation of $n+2$, write a summand m as $m-2$ 1 s followed by a 2 . This gives an ordered sum of 1 s and 2 s. Finally remove the final 2 . This gives us a representation of n . It is clear that these two operations are inverse and hence each is a bijection.

Problem B5

Let S be a set and \mathcal{P} the set of all subsets of S . $f : \mathcal{P} \rightarrow \mathcal{P}$ is such that if $X \subseteq Y$, then $f(X) \subseteq f(Y)$. Show that for some K , $f(K) = K$.

Solution

Let K be the union of all subsets A of S such that $A \subseteq f(A)$. We show that $K \subseteq f(K)$. Take any x in K . Then x belongs to some A which satisfies $A \subseteq f(A)$. But $A \subseteq K$, so $f(A) \subseteq f(K)$. So x belongs to $A \subseteq f(A) \subseteq f(K)$, so x belongs to $f(K)$.

Since $K \subseteq f(K)$, we have $f(K) \subseteq f(f(K))$. Hence by definition $f(K) \subseteq K$.

Problem B6

y is the solution of the differential equation $(x^2 + 9)y'' + (x^2 + 4)y = 0$, $y(0) = 0$, $y'(0) = 1$. Show that $y(x) = 0$ for some $x \in [\sqrt{(63/53)}\pi, 3\pi/2]$.

Solution

We compare the equation given with (1) $z'' = -4/9 z$, $z(0) = 0$, $z'(0) = 1$, and (2) $w'' = -53/63 w$, $w(0) = 0$, $w'(0) = 1$.

For all $x > 0$ we have $(x^2 + 4)/(x^2 + 9) > 4/9$, so $y''z - yz'' = (4/9 - (x^2 + 4)/(x^2 + 9)) yz$. Integrating gives $y'z - yz'|_0^{3\pi/2} = \int (4/9 - (x^2 + 4)/(x^2 + 9)) yz dx$ (*). Now $y(0) = z(0) = 0$, $z = \sin 2x/3$, so $z(3\pi/2) = 0$, $z'(3\pi/2) = -2/3$, hence $y'z - yz'|_0^{3\pi/2} = 2/3 y(3\pi/2)$. Now if y has no zeros on the open interval $(0, 3\pi/2)$, then $y(3\pi/2) \geq 0$ and the integrand $(4/9 - (x^2 + 4)/(x^2 + 9)) yz$ is negative for the entire interval. Hence lhs (*) ≥ 0 and rhs (*) < 0 . Contradiction. So y has at least one zero on $(0, 3\pi/2)$.

Turning to (2), we note that $\pi^2 < 10$, so $(3\pi/2)^2 < 22.5$. Hence on the interval $[0, 3\pi/2]$ we have $(x^2 + 4)/(x^2 + 9) < 26.5/31.5 = 53/63$. Now $w = \sin(\sqrt{(53/63)}x)$, so $w(x) > 0$ on the interval $(0, k)$, where $k = \sqrt{(63/53)}\pi$.

Now suppose that y has a root in $(0, k]$. Let h be the smallest such root, so that $y(x)$ is positive on $(0, h)$ and $y(h) = 0$, $y'(h) \leq 0$. Integrate $w''y - wy'' = ((x^2 + 4)/(x^2 + 9) - 53/63)yw$. The lhs is $w'y - wy'|_0^h = w(h)y'(h) \geq 0$. The rhs is $\int_0^h ((x^2 + 4)/(x^2 + 9) - 53/63)yw dx$. But the integrand is negative on $(0, h)$, so the rhs < 0 . Contradiction. Hence y has no roots on $(0, k]$. So it must have a root on $(k, 3\pi/2)$.

Problem B7

Let P be a regular polygon and its interior. Show that for any $n > 1$, we can find a subset S_n of the plane such that we cannot translate and rotate P to cover S_n but we can translate and rotate P to cover any n points of S_n .

Solution

Let the radius of the inscribed circle in P be 1. Let P have m sides. We take S_n to be a circle of radius slightly greater than 1. Clearly it cannot fit inside P . Place it so that its centre is at the centre of P . Suppose it cuts each side a distance $\tan^{-1}\theta$ either side of the midpoint, where θ is small. Then the arcs lying outside P subtend a total angle $2m\theta$ at the centre. Now rotate the circle about its centre. Let ϕ measure the angle of rotation from the starting point. In order to keep a given point of the circle inside P , ϕ must avoid intervals of total length $2m\theta$. So to keep any given n points of the circle inside P , ϕ must avoid intervals of total length $2mn\theta$. The worst case is that these intervals are all disjoint, but provided $\theta < \pi/mn$ they cannot exhaust the available 2π , so we can find angles ϕ for which all n points are inside P .

18th Putnam 1958

Problem A1

Show that the real polynomial $\sum_0^n a_i x^i$ has at least one real root if $\sum a_i/(i+1) = 0$.

Solution

Integrate from 0 to 1. We get zero. Hence the polynomial has at least one zero between 0 and 1.

Problem A2

A rough sphere radius R rests on top of a fixed rough sphere radius R . It is displaced slightly and starts to roll off. At what point does it lose contact?

Solution

Let the top of the fixed sphere be T and its centre O . Let X be the point of contact. Take angle $XOT = \theta$. Let the mass be M and the normal force between the two spheres N . Resolving radially, $MR(d\theta/dt)^2 = Mg \cos \theta - N$ (1). Taking moments about an axis through X , $MgR \sin \theta = I d^2\theta/dt^2$ (2), where I is the moment of inertia of the sphere about $X =$ the moment about a diameter $+ MR^2 = 7/5 MR^2$.

Integrating equation (2) gives $g(1 - \cos \theta) = 7/10 R (d\theta/dt)^2$. Substituting in (1) gives $N = g \cos \theta - 10/7 (1 - \cos \theta)$. Contact is lost when N becomes 0. In other words at the angle $\theta = \cos^{-1} 10/17$.

Problem A3

A sequence of numbers $\alpha_i \in [0, 1]$ is chosen at random. Show that the expected value of n , where $\sum_1^n \alpha_i > 1$, $\sum_1^{n-1} \alpha_i \leq 1$ is e .

Solution

We can consider the possible values of $\alpha_1, \alpha_2, \dots, \alpha_n$ as points of the n -cube. The points corresponding to sum at most 1 are those in the corner at the origin, bounded by the hyperplane through $(1, 0, \dots, 0), (0, 1, 0, \dots, 0), \dots, (0, \dots, 0, 1)$. By an easy induction this has volume $1/(n-1)!$. Let $p_n =$ the prob that the sum of the first n numbers is at most 1. We have just shown that $p_n = 1/(n-1)!$.

Now the required expected value is $(1/1! - 1/2!) 2 + (1/2! - 1/3!) 3 + (1/3! - 1/4!) 4 + \dots = 2 + (3 - 2)/2! + (4 - 3)/3! + (5 - 4)/4! + \dots = e$.

Problem A4

z_1, z_2, \dots, z_n are complex numbers with modulus $a > 0$. Let $f(n, m)$ denote the sum of all products of m of the numbers. For example, $f(3, 2) = z_1 z_2 + z_2 z_3 + z_3 z_1$. Show that $|f(n, m)| / a^m = |f(n, n-m)| / a^{n-m}$.

Solution

It is obviously sufficient to take $a = 1$. Then if $z = e^{i\theta}$ we have $1/z = e^{-i\theta}$ = the complex conjugate of z . Now $f(n, m)/(z_1 z_2 \dots z_n) =$ the sum of the inverses of the terms in $f(n, n-m) =$ the sum of the complex conjugates of the terms in $f(n, n-m) =$ complex conjugate of $f(n, n-m)$. But a number and its complex conjugate have the same modulus.

Problem A5

Let R be the reals. Show that there is at most one continuous function $f: [0, 1]^2 \rightarrow R$ satisfying $f(x, y) = 1 + \int_0^x \int_0^y f(s, t) dt ds$.

Solution

If there are two, then their difference $d(x, y)$ satisfies $d(x, y) = \int_0^x \int_0^y d(s, t) dt ds$ (*).

The domain is compact and d is continuous, so $|d|$ has some upper bound k . Now applying (*) gives $|d(x, y)| \leq k xy$. Applying (*) again gives $|d(x, y)| \leq k x^2/2 y^2/2$. So by a simple induction $|d(x, y)| \leq k x^n/n! y^n/n!$. Hence $|d(x, y)| \leq k/(n! n!)$ for all n , and so $d(x, y) = 0$.

Problem A6

Assume that the interest rate is r , so that capital of k becomes $k(1+r)^n$ after n years. How much do we need to

invest to be able to withdraw 1 at the end of year 1, 4 at the end of year 2, 9 at the end of year 3, 16 at the end of year 4 and so on (in perpetuity)?

Solution

To meet the withdrawal at the end of year n we need to invest $n^2/(1+r)^n$. Thus the total investment required is $1/(1+r) + 4/(1+r)^2 + 9/(1+r)^3 + 16/(1+r)^4 + \dots$. Now we know that $1 + x + x^2 + \dots = 1/(1-x)$. Differentiating, we get: $x/(1-x)^2 = x + 2x^2 + 3x^3 + \dots$. Differentiating again: $x(1+x)/(1-x)^3 = 1^2x + 2^2x^2 + 3^2x^3 + \dots$. Substituting $x = 1/(1+r)$ gives $(1+r)(2+r)/r^3$ as the required investment.

Problem A7

Show that we cannot place 10 unit squares in the plane so that no two have an interior point in common and one has a point in common with each of the others.

Solution

Note that we can place 9 squares - arrange them in a regular 3×3 array. Then the centre square touches 4 at the corners and the other 4 along the sides.

Consider first placing squares to touch a line. If neither square has a side in contact with the line, then it is easy to see that the two points of contact are more than 1 apart. If square U has a side in contact and square V does not, then together they occupy more than a length 1 of the line (from the far side of U to the point of contact of V).

So the only way to get three squares in contact with AB , a side of a fixed square $S = ABCD$, is to place one square so that its side is AB . One can then place a second square with its corner touching A and a third with its corner touching B .

So to get more than $8 = 4 \times 2$ squares touching S , we must get three squares to touch one side as above. Call them T_2 in contact with AB , T_1 touching at A and T_3 touching at B . Now consider the side BC . T_3 has a side BD . The most favourable case is that angle $CBD = 90^\circ$ - T_3 may be placed so that angle CBD is less than 90° . But certainly any square touching BC (apart from T_3) cannot extend beyond the ray BD which is perpendicular to BC . But it is easy to see that this means we can get only one additional square to touch BC unless we place T_4 with so that its side is BC and T_5 touching C at a corner, which gives 2.

In particular, if 3 squares touch AB , then at most 2 (additional) squares touch each of BC and DA .

Hence at most 2 sides, which must be opposite, can have three squares touching. But in that case it is easy to see that the other two sides each have at most 1 additional square touching, so we get only 8 in total. Hence the only way to get 9 is to have 3 squares touching AB and 2 (additional) squares touching each of the other three sides. But those touching BC and DA must then be placed as indicated above and it is easy to see that we then get at most 1 (additional) square touching CD .

Problem B1

Do both (1) and (2):

(1) Given real numbers a, b, c, d with $a > b, c, d$, show how to construct a quadrilateral with sides a, b, c, d and the side length a parallel to that length b . What conditions must a, b, c, d satisfy?

(2) H is the foot of the altitude from A in the acute-angled triangle ABC . D is any point on the segment AH . BD meets AC at E , and CD meets AB at F . Show that $\angle AHE = \angle AHF$.

Solution

(1) A necessary and sufficient condition is that we should be able to form a triangle with sides $c, d, (a-b)$. Hence we require $b + c + d \geq a$, $a + d \geq b + c$, and $a + c \geq b + d$. Construct two such triangles, one at each end of the segment a , then join the two tops to get the required quadrilateral.

(2) Take a line through A parallel to BC and project CF, HF, HE, BE to meet it in W, X, Y, Z . Then using similar triangles, we deduce that $AY/CH = AZ/BC = BH \cdot AW/CH (1/BC) = BH/CH \cdot AW/BC = BH/CH \cdot AX/BH = AX/CH$, so $AX = AY$. But AH is perpendicular to XY , so the result follows.

Problem B2

Let n be a positive integer. Prove that $n(n+1)(n+2)(n+3)$ cannot be a square or a cube.

Solution

$(n+1)(n+2) = n^2 + 3n + 2$ and $n(n+3) = n^2 + 3n$. So their product is $(n^2 + 3n + 1)^2 - 1$. Hence $n(n+1)(n+2)(n+3)$ is 1 less than a square, so it cannot be a square.

One of $n+1, n+2$ must be odd. Suppose it is $n+1$. Then $n+1$ has no factor in common with $n(n+2)(n+3)$, so $n(n+2)(n+3) = n^3 + 5n^2 + 6n$ must be a cube. But $(n+1)^3 = n^3 + 3n^2 + 3n + 1 < n^3 + 5n^2 + 6n < n^3 + 6n^2 + 12n + 8 = (n+2)^3$, so $n^3 + 5n^2 + 6n$ cannot be a cube.

Similarly, suppose $n+2$ is odd. Then it has no factor in common with $n(n+1)(n+3) = n^2 + 4n^2 + 3n$, so $n^2 + 4n^2 + 3n$ must be a cube. But for $n \geq 2$, $(n+1)^3 < n^2 + 4n^2 + 3n < (n+2)^3$, so $n^2 + 4n^2 + 3n$ cannot be a cube for $n \geq 2$. The case $n = 1$ is checked by inspection: 24 is not a cube.

Problem B3

In a tournament of n players, every pair of players plays once. There are no draws. Player i wins w_i games. Prove that we can find three players i, j, k such that i beats j , j beats k and k beats i iff $\sum w_i^2 < (n-1)n(2n-1)/6$.

Solution

Suppose there are no such three players. Let A be the (or one of the) top-scoring players. We claim that A has $n-1$ wins. Suppose not, then he is beaten by some B . Now if every player beaten by A is also beaten by B , then B would have a higher score than A , so we must be able to find C who is beaten by A , but who beats B (who beats A). But we assumed there were no such triads. So A has $n-1$ wins. But we can now consider the remaining players. The same argument shows that the top scoring player amongst them beat $n-2$ of them. He did not beat A (who beat everyone), so his score is $n-2$. The argument repeats to show that the scores are $n-1, n-2, \dots, 1, 0$. The sum of the squares is thus $(n-1)n(2n-1)/6$.

Conversely, it is clear that if the scores are $n-1, n-2, \dots, 1, 0$ then there can be no triad. For the player with $n-1$ beats everyone, so he cannot be part of a triad. But now the player with $n-2$ beats everyone else, so he cannot either and so on.

Thus we have shown that there is a triad iff the scores are not $n-1, n-2, \dots, 1, 0$. It remains to show that if the scores are not $n-1, n-2, \dots, 1, 0$, then sum of the squares is less than $(n-1)n(2n-1)/6$. But if the scores are not $n-1, n-2, \dots, 1, 0$, then two players must have the same score m . Changing the scores to $m-1, m+1$ (by changing the result of the match between the two players) increases the sum of the squares by 2. After a finite number of repetitions we must arrive at $n-1, n-2, \dots, 1, 0$, so the starting sum must be strictly less.

Problem B4

Let S be a spherical shell radius 1. Find the average straight line distance between two points of S . [In other words S is the set of points (x, y, z) with $x^2 + y^2 + z^2 = 1$].

Solution

Answer: $4/3$.

It is sufficient to fix one point and to find the average distance of the other points from it. Take the point as P and the centre as O . Now consider a general point Q on the surface. Let angle $POQ = \theta$. The distance PQ is $2 \sin \theta/2$ and this is the same for all points in a band angular width $d\theta$ at the angle θ . The band has radius $\sin \theta$. Hence the average distance is $1/4\pi \int_0^\pi (2\pi \sin \theta) (2 \sin \theta/2) d\theta = \int 4 \sin^2 \theta/2 d(\sin \theta/2) = 4/3$.

Problem B5

S is an infinite set of points in the plane. The distance between any two points of S is integral. Prove that S is a subset of a straight line.

Solution

Suppose not. Take three non-collinear points A, B, C . Suppose $AB = n$. Any point P has $PA + AB \geq PB$, so $PB - PA \leq n$. Similarly $PB - PA \geq -n$. But $PB - PA$ is integral, so it must take one of the values $-n, -(n-1), \dots, 0, 1, \dots, n$. So P must lie on one of a finite number of hyperbolae $|PB - PA| = k$ (regarding the allowed line pair as a degenerate

hyperbola). Similarly it must lie on one of a finite number of hyperbolae $|PA - PC| = h$. But each pair of hyperbolae intersect in at most 4 points, so the number of points in addition to A, B, C is finite. Contradiction.

Problem B6

A particle of unit mass moves in a vertical plane under the influence of constant gravitational force g and a resistive force which is in the opposite direction to its velocity and with magnitude a function of its speed. The particle starts at time $t = 0$ and has coordinates (x, y) at time t . Given that $x = x(t)$ and is not constant, show that $y(t) = -g x(t) \int_0^t ds/x'(s) + g \int_0^t x(s) / x'(s) ds + a x(t) + b$, where a and b are constants.

Solution

We can write the equations of motion as $x'' = -f(x', y') x'$, $y'' = -f(x', y') y' - g$. If $x'(t) = 0$ at some t , then a possible solution to the equations would be $x = \text{constant}$ (and y found by integrating $y'' = -f(0, y') y' - g$). But solutions are unique, so this would be *the* solution. But we are told that x is not constant. So $x'(t)$ is never zero.

We do not know anything about f , so we resolve perpendicular to it to get $x'' \sin \theta - y'' \cos \theta = g \cos \theta$, or $x'' \tan \theta - y'' = g$. But $\tan \theta = dy/dx = y'/x'$, so $x'' y'/x' - y'' = g$ (*).

Dividing by x' gives $x'' y'/(x')^2 - y''/x' = g/x'$. Integrating, $y'/x' = A - g \int dt/x'$. So $y' = Ax' - g x' \int dt/x'$. Integrating again, $y(T) = A x + B - g \int_0^T x' \int_0^t ds/x' dt$. Now we can integrate the last integral by parts: $\int_0^T x' \int_0^t ds/x' dt = \int_{t=0}^T \int_0^t ds/x' dx = x(T) \int_0^T ds/x' - \int_0^T x/x' dt$. So we get finally the expression in the question.

Problem B7

\mathbb{R} is the reals. $f : [a, b] \rightarrow \mathbb{R}$ is continuous and $\int_a^b x^n f(x) dx = 0$ for all non-negative integers n . Show that $f(x) = 0$ for all x .

Solution

We deduce immediately that $\int_a^b x^n f(x) p(x) dx = 0$ for any polynomial $p(x)$. We now need the Weierstrass approximation theorem: we can uniformly approximate any continuous function on a compact set by polynomials. In other words, we can find a polynomial $p(x)$ such that $|f(x) - p(x)| < \epsilon$ for all x in $[a, b]$. Also, since $[a, b]$ is compact, f must be bounded, say by k , so $|f(x)| < k$ for all x in $[a, b]$. Now we have that $|\int_a^b f(x)^2 dx| = |\int_a^b f(x) (f(x) - p(x)) dx| \leq \int_a^b |f(x)| |f(x) - p(x)| dx \leq (b - a) k \epsilon$, which can be made arbitrarily small by choosing ϵ sufficiently small. Hence $\int_a^b f(x)^2 dx = 0$, so $f(x) = 0$ throughout the interval.

19th Putnam 1958

A1. Define $f(i, j)$ as follows: $f(i, 1) = f(1, j) = 1$, $f(i+1, j+1) = f(i, j+1) + f(i+1, j) + f(i, j)$. Let $d(n) = f(1, n-1) + f(2, n-2) + \dots + f(n-1, 1)$. Show that $d(n+2) = d(n) + 2d(n+1)$.

A2. Define $a_1 = 1$, $a_{n+1} = 1 + n/a_n$. Show that $\sqrt{n} \leq a_n < 1 + \sqrt{n}$.

A3. Assuming that there is a unique function $f(x)$ satisfying $f(0) = 1$, $f'(x) = f(x) + \int_0^1 f(t) dt$, find it.

A4. Find the general solution in real numbers a, b, c, d to the inequalities $2a > a + b > a + c > 2b > b + c > a + d > 2c > b + d > c + d > d + d$. Find the smallest solution in positive integers.

A5. Let $A = (a_{ij})$ be the $n \times n$ matrix with $a_{ij} = 1$ if $i \neq j$, and $a_{ii} = 0$. Show that the number of non-zero terms in the expansion of $\det A$ is $n! \sum_{i=0}^{n-1} (-1)^i / i!$.

A6. \mathbb{R} is the reals. $\alpha \in [0, 1)$. $f : [0, 1] \rightarrow [0, \alpha]$ and $g : [0, 1] \rightarrow \mathbb{R}$ are continuous. β satisfies $\beta = \max (g(x) + f(x) - \beta)$. When do we have $\beta = \max g(x) / (1 - f(x))$?

A7. m, n are relatively prime positive integers with n even. Given any positive integer r , define $f(r)$ to be the integer which minimizes $|f(r)/r - m/n|$. Show that $\lim_{k \rightarrow \infty} \sum_{r=1}^k |f(r)/r - m/n| r/k = 1/4$.

B1. Let $a_n = \sum_{r=0}^n 1/nCr$. Show that $a_n = 1 + a_{n-1} (n+1)/(2n)$. Deduce that $\lim a_n = 2$.

B2. Let X be the set $\{1, 2, 3, \dots, 2n\}$, take $Y \subseteq X$ with $|Y| = n+1$. Show that we can find $a, b \in Y$ with a dividing b .

B3. Show that if X is a square side 1 and $X = A \cup B$, then A or B has diameter at least $\sqrt{5}/2$. Show that we can find A and B both having diameter $\leq \sqrt{5}/2$.

B4. Let \mathbb{R} be the reals. $f : \mathbb{R} \rightarrow \mathbb{R}$ is three times differentiable. As $x \rightarrow \infty$, $f(x)$ tends to a finite limit, and $f'''(x)$ tends to zero. Show that $f'(x)$ and $f''(x)$ also tend to zero.

B5. A sequence of points P_n in the plane is defined by: P_0 is at the origin; P_1 is at $(1, 0)$; and $P_{n-1}P_n$ is length $1/n$ and at an angle θ to the previous segment. Find the coordinates of $\lim P_n$.

B6. A graph has n vertices $\{1, 2, \dots, n\}$ and a complete set of edges. Each edge is oriented, as either $i \rightarrow j$ or $j \rightarrow i$. Show that we can find a permutation of the vertices a_i so that $a_1 \rightarrow a_2 \rightarrow a_3 \rightarrow \dots \rightarrow a_n$.

B7. Let $N = \{1, 2, \dots, n\}$. Given a permutation $f : N \rightarrow N$, define $d(f) =$ no. of i such that $f(i) > f(j)$ for all $j > i$. Find the mean of $d(f)$ over all permutations f on N .

20th Putnam 1959

Problem A1

Prove that we can find a real polynomial $p(y)$ such that $p(x - 1/x) = x^n - 1/x^n$ (where n is a positive integer) iff n is odd.

Solution

Take n odd. For $n = 1$ the result is obvious. Now suppose the result is true for odd $n \leq 2m - 1$. Expand $(x - 1/x)^{2m+1}$ by the binomial theorem. Since $(2m+1)C_0 = (2m+1)C_{2m+1}$, $(2m+1)C_1 = (2m+1)C_{2m}$, $(2m+1)C_2 = (2m+1)C_{2m-1}$, ..., $(2m+1)C_m = (2m+1)C_{m+1}$, we may group these pairs of terms together to get: $(x - 1/x)^{2m+1} = (x^{2m+1} - x^{-(2m+1)}) - (2m+1)C_1 (x^{2m-1} - x^{-(2m-1)}) + \dots + (-1)^m (2m+1)C_m (x - 1/x)$. This gives the polynomial for $n = 2m+1$ in terms of the lower order polynomials.

Note that $x - 1/x$ has the same value $3/2$ for $x = -1/2$ and $x = 2$. But $x^{2m} - 1/x^{2m}$ is negative for $x = -1/2$ and positive for $x = 2$, so we cannot express $x^{2m} - 1/x^{2m}$ as a function of $(x - 1/x)$, polynomial or otherwise.

Problem A2

Let $\omega^3 = 1$, $\omega \neq 1$. Show that $z_1, z_2, -\omega z_1 - \omega^2 z_2$ are the vertices of an equilateral triangle.

Solution

Let A be the point z_1 , B the point z_2 . Then $z_2 - z_1$ represents the vector from A to B . Now $-\omega^2$ has unit length and makes an angle $\pm\pi/3$ with the positive real axis, so multiplying $z_2 - z_1$ by it rotates AB through an angle $\pi/3$ (clockwise or counterclockwise). Adding the result to $z_2 - z_1$ gives a point C such that AC is at an angle $\pi/3$ to AB . In other words ABC is equilateral. It is easily checked that C is then $z_1 - \omega^2(z_2 - z_1) = -\omega z_1 - \omega^2 z_2$ (since $1 + \omega + \omega^2 = 0$).

Problem A3

Let C be the complex numbers. $f : C \rightarrow C$ satisfies $f(z) + z f(1 - z) = 1 + z$ for all z . Find f .

Solution

Putting $z = 1 - w$ we have $f(1 - w) + (1 - w) f(w) = 2 - w$. Hence $w f(1 - w) + (w - w^2) f(w) = 2w - w^2$. Hence $(1 + w - f(w)) + (w - w^2) f(w) = 2w - w^2$, giving $(w^2 - w + 1) f(w) = (w^2 - w + 1)$. So provided w is not $(1 \pm i\sqrt{3})/2$, we have $f(w) = 1$.

But at these two values f can be different. We can take one of them to be arbitrary. For example, take $f((1 + i\sqrt{3})/2) = k$, any complex number. Then $f((1 - i\sqrt{3})/2) = 1 + (1 - k)(1 - i\sqrt{3})/2$.

Problem A4

R is the reals. $f, g : [0, 1] \rightarrow R$ are arbitrary functions. Show that we can find x, y such that $|xy - f(x) - g(y)| \geq 1/4$.

Solution

It is enough to consider the values at 0 and 1. If the pairs $(0, 0)$, $(0, 1)$, $(1, 0)$ do not work for (x, y) , then we have $|f(0) + g(0)| < 1/4$, $|f(0) + g(1)| < 1/4$, and $|f(1) + g(0)| < 1/4$. Hence $f(1) + g(1) \leq f(1) + g(0) + f(0) + g(1) - (f(0) + g(0)) < 1/4 + 1/4 + 1/4$. So $1 - f(1) - g(1) > 1 - 3/4 = 1/4$. Hence the pair $(1, 1)$ does work.

Problem A5

At a particular moment, A, T and B are in a vertical line, with A 50 feet above T , and T 100 feet above B . T flies in a horizontal line at a fixed speed. A flies at a fixed speed directly towards B , B flies at twice T 's speed, also directly towards T . A and B reach T simultaneously. Find the distance traveled by each of A, B and T , and A 's speed.

Solution

Answer: A travels $25(3 + \sqrt{73})/3 = 96.2$ ft, B travels $400/3$ ft = 133.3 ft, T $200/3$ ft = 66.7 ft, A 's speed is $(3 + \sqrt{73})/8 = 1.443$ times T 's speed.

Take the x -axis vertical and the y -axis horizontal, so that at $t = 0$, the target is at $(0, 0)$ and the pursuer is at $(a, 0)$. Assume the target (T) has speed v and the pursuer (A or B) has speed kv with $k > 1$. At time t , the target is at $(0,$

vt). The equations of motion are: $(dx/dt)^2 + (dy/dt)^2 = kv$ and $y' = (y - vt)/x$. The first equation gives $kv/(dx/dt) = -\sqrt{(1 + y'^2)}$ (negative because x decreases with time).

Differentiating the second equation gives $xy'' = -v/(dx/dt)$. So $kxy'' = \sqrt{(1 + y'^2)}$. Integrating, $\ln(y' + \sqrt{(1 + y'^2)}) = \ln x + \text{const}$. At $x = a$, $y = 0$, so $(x/a)^{1/k} = y' + \sqrt{(1 + y'^2)}$. Hence $2y' = (x/a)^{1/k} - (x/a)^{-1/k}$. Integrating again, $2y = a/(1+1/k) (x/a)^{1+1/k} - a/(1-1/k) (x/a)^{1-1/k} - a/(1+1/k) + a/(1-1/k)$, since $y = 0$ at $x = a$. When pursuit ends, $x = 0$ and hence $y = ak/(k^2 - 1)$.

For B, $k = 2$, so $y = 2a/3 = 200/3$ ft. That is the distance travelled by T. B is travelling at twice the speed, so B travels a distance $400/3$ ft. For A reaches T at the same y , hence $200/3 = 50 k/(k^2 - 1)$, so $4k^2 - 3k - 4 = 0$, so $k = (3 + \sqrt{73})/8$.

Problem A6

Given any real numbers $\alpha_1, \alpha_2, \dots, \alpha_m, \beta$, show that for $m, n > 1$ we can find m real $n \times n$ matrices A_1, \dots, A_m such that $\det A_i = \alpha_i$, and $\det(\sum A_i) = \beta$.

Solution

Start by setting A_i to be the matrix with $1, 1, \dots, 1, \alpha_i$ down the main diagonal and zeros elsewhere. Modify A_1 by changing the $n, n-1$ element to 1 . Modify A_m by changing the $n-1, n$ element to $m(\alpha_1 + \dots + \alpha_m) - \beta/m^{n-2}$. It is clear that this gives $\det A_i = \alpha_i$.

$A_1 + \dots + A_m$ has $m, m, \dots, m, (\alpha_1 + \dots + \alpha_m)$ down the main diagonal. The only other non-zero elements are $n, n-1$, which is 1 and $n-1, n$, which is $m(\alpha_1 + \dots + \alpha_m) - \beta/m^{n-2}$. Hence its determinant evaluates to β .

Problem A7

Let R be the reals. Let $f : [a, b] \rightarrow R$ have a continuous derivative, and suppose that if $f(x) = 0$, then $f'(x) \neq 0$. Show that we can find $g : [a, b] \rightarrow R$ with a continuous derivative, such that $f(x)g'(x) > f'(x)g(x)$ for all $x \in [a, b]$.

Solution

We note that the derivative of f/g is $-(fg' - f'g)/g^2$ so if we take g so that f/g is decreasing, then we are almost home. Not quite, because of the difficulty that f may be zero. For example, if we take $g(x) = x f(x)$, then $fg' - f'g$ is zero whenever $f(x) = 0$.

f has only finitely many zeros, for otherwise the zeros would have a limit point c and then by continuity we would have $f(c) = f'(c) = 0$. So we may take a polynomial $p(x)$ such that $p(x) f'(x) = -1$ at each zero of f . Now consider $g(x) = x f(x) + k p(x)$. We have $f(x) g'(x) - f'(x) g(x) = f(x)^2 + k(f(x) p'(x) - f'(x) p(x))$. Now $f(x) p'(x) - f'(x) p(x) = 1$ at each zero of f , so we can find an open set K containing all these zeros such that $f(x) p'(x) - f'(x) p(x) > 1/2$ on K . Now $[a, b] - K$ is compact and contains no zeros of $f(x)^2$, which is continuous, so we can find $\epsilon > 0$, so that $f(x)^2 > \epsilon$ on $[a, b] - K$. But $[a, b]$ is compact, so $|f(x) p'(x) - f'(x) p(x)| < \text{some } M$ on $[a, b]$. Take $k < \epsilon/M$. Then on $[a, b] - K$, $f(x)^2 > \epsilon$ and $k|f(x) p'(x) - f'(x) p(x)| < \epsilon$, so $f(x) g'(x) - f'(x) g(x) > 0$. On K , $f(x)^2$ is non-negative and $k(f(x) p'(x) - f'(x) p(x))$ is positive, so $f(x) g'(x) - f'(x) g(x) > 0$.

Problem B1

Join each of m points on the positive x -axis to each of n points on the positive y -axis. Assume that no three of the resulting segments are concurrent (except at an endpoint). How many points of intersection are there (excluding endpoints)?

Solution

Answer: $mn(m-1)(n-1)/4$.

Let us call the points on the y -axis $y_1 < y_2 < \dots < y_n$ and the points on the x -axis $x_1 < x_2 < \dots < x_m$. Let us join first y_1 to each x_i , then y_2 and so on. Joining y_1 to each of the x_i creates no intersections. Joining y_2 to x_1 creates $m-1$ intersections. Joining it to x_2 creates $m-2$ intersections and so on. So, in all, joining y_2 gives $1 + 2 + \dots + m-1 = m(m-1)/2$ intersections.

Similarly joining y_3 to x_1 gives $2(m-1)$ intersections. Joining it to x_2 gives $2(m-2)$ and so on. So, in all, y_3 gives $2 \times m(m-1)/2$. Similarly, y_{r+1} gives $r \times m(m-1)/2$. So in total we get $m(m-1)/2 (1 + 2 + \dots + n-1) = mn(m-1)(n-1)/4$.

Problem B2

Show that any positive real can be expressed in infinitely many ways as a sum $\sum 1/(10 a_n)$, where $a_1 < a_2 < a_3 < \dots$ are positive integers.

Solution

$\sum 1/n$ diverges, so $\sum 1/(10n)$ diverges. Let k be any positive real. Take N such that $1/(10N) < k$. Let M be any integer $> N$. We show how to find an expression $k = \sum 1/(10a_n)$ with $a_1 = M$. Take enough terms $a_2 = 1/(10(M+1))$, $a_3 = 1/(10(M+2))$, ... so that $\sum 1/(10a_n) < k$, but adding another term would give a sum $\geq k$. Let the difference $k - \sum 1/(10a_n)$ be $k_1 > 0$. Now take M_1 so that $1/(10 M_1) < k_1$ and repeat. [In other words, take as many terms $1/(10 M_1) + 1/(10(M_1+1)) + \dots$ as we can whilst keeping the sum $< k_1$.] Let k_2 be the new difference, and so on. This process gives a sum which converges to k . Each such sum is different because it has a different starting term.

Problem B3

Find a continuous function $f : [0, 1] \rightarrow [0, 1]$ such that given any $\beta \in [0, 1]$, we can find infinitely many α such that $f(\alpha) = \beta$.

Solution

We use an adapted space-filling curve. Let $g(x)$ be a piecewise linear function with period 2: $g(x) = 0$ on $[0, 1/3]$, 1 on $[2/3, 4/3]$ and 0 on $[5/3, 2]$ with linear connecting pieces. For t any real in the range $0 \leq t \leq 1$ let $f(t) = g(t)/2 + g(9t)/4 + g(81t)/8 + g(729t)/16 + \dots$. The series is obviously uniformly convergent, so f is continuous.

Now suppose that in base 3 we have $t = 0.c_1c_2c_3 \dots$ with all $c_n = 0$ or 2. Then $9^n t = \text{even integer} + 0.c_{2n+1}c_{2n+2}\dots$. Now $0.c_{2n+1}c_{2n+2}\dots$ lies in $[0, 1/3]$ if $c_{2n+1} = 0$ and in $[2/3, 1]$ if $c_{2n+1} = 2$, so $g(9^n t) = 0$ if $c_{2n+1} = 0$, 1 if $c_{2n+1} = 2$. Thus $f(t)$ has the binary expansion $0.a_1a_2a_3 \dots$, where $a_n = c_{2n-1}/2$. So given any s in $[0, 1]$ we can find infinitely many t with $f(t) = s$, for odd n pick c_n to give the correct binary expansion and for even n pick c_n arbitrarily.

Problem B4

A is the 5 x 5 array:

11	17	25	19	16
24	10	13	15	3
12	5	14	2	18
23	4	1	8	22
6	20	7	21	9

Pick 5 elements, one from each row and column, whose minimum is as large as possible (and prove it so).

Solution

Answer: 25, 23, 20, 18, 15.

We can only pick one of 16, 17, 19, 25; only one of 23, 24; only one of 18, 22; and only one of 20, 21. That only gives 4 elements, so the minimum cannot be larger than 15. The answer shows that 15 can be achieved.

Problem B5

L_1 is the line $\{ (t + 1, 2t - 4, -3t + 5) : t \text{ real} \}$ and L_2 is the line $\{ (4t - 12, -t + 8, t + 17) : t \text{ real} \}$. Find the smallest sphere touching L_1 and L_2 .

Solution

The sphere has diameter PQ, where $P(s + 1, 2s - 4, -3s + 5)$ is on L_1 , and $Q(4t - 12, -t + 8, t + 17)$ is on L_2 . PQ is as short as possible. It can also be characterised as perpendicular to L_1 and L_2 .

$PQ^2/2 = 7s^2 + st + 9t^2 + 25s - 52t + 457/2$. At the minimum the two partial derivatives must be zero, so $14s + t + 25 = 0$, $s + 18t - 52 = 0$. Solving $s = -2$, $t = 3$. This gives $P(-3, -12, 11)$, $Q(0, 5, 20)$, the radius ($= PQ/2$) as $1/2 \sqrt{251}$ and the centre as $(-3/2, -7/2, 31/2)$.

Problem B6

α and β are positive irrational numbers satisfying $1/\alpha + 1/\beta = 1$. Let $a_n = [n\alpha]$ and $b_n = [n\beta]$, for $n = 1, 2, 3, \dots$. Show that the sequences a_n and b_n are disjoint and that every positive integer belongs to one or the other.

Solution

β positive implies $\alpha > 1$, so $[\alpha(n+1)] > [\alpha n]$ and the sequence does not contain any integers twice. Similarly for $[\beta n]$.

If k appears in both sequences, then for some m, n we have: $k < n\alpha < k+1$, $k < m\beta < k+1$. Hence $k/\alpha < n < (k+1)/\alpha$, $k/\beta < m < (k+1)/\beta$. Adding gives $k < m+n < k+1$. Contradiction. So an integer appears in at most one sequence.

Suppose k does not appear in the sequence $[n\alpha]$. Then for some n we have $n\alpha < k$, and $k+1 < (n+1)\alpha$. The first inequality implies that $n < k/\alpha = k - k/\beta$. Hence $(k-n)\beta > k$. The second inequality implies that $n+1 > (k+1)/\alpha = k+1 - (k+1)/\beta$. Hence $(k-n)\beta < k+1$. So $[(k-n)\beta] = k$.

Problem B7

Given any finite ordered tuple of real numbers X , define a real number $[X]$, so that for all x_i, α :

(1) $[X]$ is unchanged if we permute the order of the numbers in the tuple X ;

(2) $[(x_1 + \alpha, x_2 + \alpha, \dots, x_n + \alpha)] = [(x_1, x_2, \dots, x_n)] + \alpha$;

(3) $[(-x_1, -x_2, \dots, -x_n)] = -[(x_1, x_2, \dots, x_n)]$;

(4) for $y_1 = y_2 = \dots = y_n = [(x_1, x_2, \dots, x_n)]$, we have $[(y_1, y_2, \dots, y_n, x_{n+1})] = [(x_1, x_2, \dots, x_{n+1})]$.

Show that $[(x_1, x_2, \dots, x_n)] = (x_1 + x_2 + \dots + x_n)/n$.

Solution

It is convenient to write $[(x_1, \dots, x_n)]$ simply as $[x_1, \dots, x_n]$. We use induction on n . For $n = 1$, we have $[0] = -[0]$, so $[0] = 0$. Then (3) gives that $[x] = x$, which establishes the result for $n = 1$.

$[x, 0] = -[-x, 0] = -[0, -x] = -[x, 0] + x$, so $[x, 0] = x/2$. Hence $[x, y] = y + [x - y, 0] = y + (x - y)/2 = (x + y)/2$, which establishes the result for $n = 2$. Now suppose it is true for n .

We have that $[x, -x, 0, 0, \dots, 0] = -[-x, x, 0, 0, \dots, 0]$ by (3), $= -[x, -x, 0, 0, \dots, 0]$ by (1), so $[x, -x, 0, 0, \dots, 0] = 0$. Now suppose $x_1 + x_2 + \dots + x_{n+1} = 0$, then the n numbers x_1, x_2, \dots, x_n and the n numbers $0, 0, \dots, 0, -x_{n+1}$ have the same sum, so by (4) we have for any z that $[x_1, \dots, x_n, z] = [0, 0, \dots, 0, -x_{n+1}, z]$. Take $z = x_{n+1}$, then $[0, \dots, 0, -x_{n+1}, x_{n+1}] = 0$, so $[x_1, \dots, x_n, x_{n+1}] = 0$.

Finally, given arbitrary x_1, \dots, x_{n+1} , let k be their mean, then $x_1 - k, x_2 - k, \dots, x_{n+1} - k$ have zero sum, so $[x_1 - k, x_2 - k, \dots, x_{n+1} - k] = 0$. Hence $[x_1, \dots, x_{n+1}] = k$, as required.

21st Putnam 1960

Problem A1

For n a positive integer find $f(n)$, the number of pairs of positive integers (a, b) such that $ab/(a+b) = n$.

Solution

$ab = n(a+b)$ implies $a(b-n) = nb > 0$, so a and b must both exceed n . Let $b = n+d$, then $a = n(n+d)/d = n + n^2/d$. This is a solution iff d divides n^2 . So $f(n) =$ the number of divisors of n^2 .

Problem A2

Let S be the set consisting of a square with side 1 and its interior. Show that given any three points of S , we can find two whose distance apart is at most $\sqrt{6} - \sqrt{2}$.

Solution

Suppose we can find three points of S so that their three distances exceed $k = \sqrt{6} - \sqrt{2}$. If one of the points is not on the top side of S and another on the bottom side, we can expand S by a combination of translation and expansion in the direction perpendicular to those sides to get three points with larger minimum distance. Similarly, if one of the points is not on each of the other two sides of S , we can expand S by a combination of expansion and translation to get three points with larger minimum distance. So we end up with a point on each side. But there are only three points, so one point must be at a vertex of the square. The other points on the two sides meeting at the vertex are all a distance $\leq 1 < k$ from the vertex, so the other two points must be on the other two sides. Let the vertices be A, B, C, D (in order). Assume the vertex point is at A . Let X be the point on BC a distance $2 - \sqrt{3}$ from B . Then $AX = k$, so the point on BC must lie on the segment XC . Similarly, let Y be the point on DC a distance $2 - \sqrt{3}$ from D . Then $AY = k$, so the point on DC must lie on the segment YC . But $XY = k$, so the two points will be closer than k unless one is at X and the other at Y but in that case the distances AX, AY do not exceed k . So it is not possible to arrange for the three points to have all distances $> k$.

Problem A3

Let $\alpha, \beta, \gamma, \delta, \varepsilon$ be arbitrary reals. Show that $(1-\alpha)e^\alpha + (1-\beta)e^{\alpha+\beta} + (1-\gamma)e^{\alpha+\beta+\gamma} + (1-\delta)e^{\alpha+\beta+\gamma+\delta} + (1-\varepsilon)e^{\alpha+\beta+\gamma+\delta+\varepsilon} \leq k_4$, where $k_1 = e, k_2 = k_1^e, k_3 = k_2^e, k_4 = k_3^e$ (so k_4 is 10^k with k approx 1.66 million).

Solution

Differentiating shows that the maximum of $(1-x)e^x$ is 1 and occurs at $x=0$. Now consider $(1-\alpha)e^\alpha + (1-\beta)e^{\alpha+\beta} = e^\alpha(1-\alpha + (1-\beta)e^\beta)$. For any given α , the maximum over β is 1, so we have to maximise $e^\alpha(2-\alpha)$.

Differentiating shows that this has maximum $k_1 = e$ at $\alpha = 1$.

Now consider $(1-\alpha)e^\alpha + (1-\beta)e^{\alpha+\beta} + (1-\gamma)e^{\alpha+\beta+\gamma} = e^\alpha(1-\alpha + (1-\beta)e^\beta + (1-\gamma)e^{\beta+\gamma})$. We have just shown that the maximum of $(1-\beta)e^\beta + (1-\gamma)e^{\beta+\gamma}$ is k_1 at $\beta = 1, \gamma = 0$. So we have to maximise $(1-\alpha + k_1)e^\alpha$.

Differentiating shows that this is k_2 at $\alpha = k_1$.

Similarly, the maximum of $(1-\alpha)e^\alpha + (1-\beta)e^{\alpha+\beta} + (1-\gamma)e^{\alpha+\beta+\gamma} + (1-\delta)e^{\alpha+\beta+\gamma+\delta}$ is k_3 at $\alpha = k_2, \beta = k_1, \gamma = 1, \delta = 0$. Finally, the maximum of the desired expression is k_4 at $\alpha = k_3, \beta = k_2, \gamma = k_1, \delta = 1, \varepsilon = 0$.

Problem A4

Given two points P, Q on the same side of a line l , find the point X which minimises the sum of the distances from X to P, Q and l .

Solution

wlog take P to the right of Q and Q to be closer to the line than P . Take A on the line so that PA makes a 30° angle with the line and A is to the left of P . Similarly, take B on the line so that QB makes a 30° angle with the line and B is to the right of Q . There are three cases to consider.

(1) If the two segments PA and QB intersect, then their point of intersection X is the required point.

(2) If A is to the right of B , then join P to Q' the reflection of Q in the line. X is the point at which PQ' meets the line.

(3) If PA and QB do not meet, but A is to the left of B , then take $X = Q$.

We must now prove these statements. It is convenient to take the line l to be the x -axis. Take Q to be (a, b) and P to be (c, d) . We assume that $d > b > 0$. Notice first that the sum of the three distances from the point X with

coordinates (x, y) , which we will write as $f(x, y)$, is a continuous and differentiable function of x and y . Also it tends to infinity as x or y tends to infinity. Hence the minimum must occur either (A) at a point where $\text{grad } f = 0$, or (B) on the boundary of the allowed region, in other words on $y = 0$, or (C) at a point where $\text{grad } f$ does not exist.

We have $f(x, y) = y + \sqrt{(x-a)^2 + (y-b)^2} + \sqrt{(x-c)^2 + (y-d)^2}$. Hence $\text{grad } f = ((x-a)/QX + (x-c)/PX, 1 + (y-b)/QX + (y-d)/PX) = \mathbf{r} + \mathbf{q} + \mathbf{p}$, where \mathbf{r} is the unit vector along the perpendicular line from the line l to X , \mathbf{q} is the unit vector from Q to X , and \mathbf{p} is the unit vector from P to X . By resolving along the line perpendicular to \mathbf{r} we find that the angles between \mathbf{q} and \mathbf{r} and between \mathbf{p} and \mathbf{r} must be equal. Similarly, the other pairs of angles. Hence the angle between each pair of $\mathbf{p}, \mathbf{q}, \mathbf{r}$ is 120° . But it is clear that the only possible such point is the intersection H of the segments PA and QB (if it exists). H is thus the only candidate of type (A).

It is obvious that for points X on the line l , $f(x, y)$ is minimised at K , the intersection of the line with PQ' (joining P to the reflection of Q), so that is the only candidate of type (B). Finally, the only points where the existence of $\text{grad } f$ is doubtful are P and Q and indeed it is easy to see that $\text{grad } f$ is not well defined there (for example, the limits as we approach parallel to the two axes are different). But clearly the distance sum is lower at Q than at P , so Q is the only candidate of type (C).

We now look separately at cases (1), (2) and (3). In case (1) all three candidates H, K and Q are available. We show first that H is better than Q . Let the perpendicular from Q to l meet l at C and the perpendicular from H to l meet l at D . Then angle HQC is 60° , so $QC = HD + QH/2$. By the cosine rule we have $PQ^2 = PH^2 + QH^2 + PH \cdot QH > (PH + QH/2)^2$. Hence $PQ + QH/2 > PH + QH$ and so $PQ + QC = PQ + QH/2 + HD > PH + QH + HD$. Next we show that H is better than K . As before let HD be the perpendicular to l . It is sufficient to show that $PH + QH + HD < PD + QD$, since we know that $PD + QD \leq PK + QK$. Now $PD^2 = PH^2 + HD^2 + PH \cdot HD > (PH + HD/2)^2$, so $PD > PH + HD/2$. Similarly, $QD > QH + HD/2$. Adding gives the required result. That shows that H minimises the distance sum for case (1).

In case (2) only candidates K and Q are available. Take QC and PE as the perpendiculars to l . Then $PQ^2 = CE^2 + (PE - QC)^2$, $PQ'^2 = CE^2 + (PE + QC)^2$. We wish to show that $PQ' < PQ + QC$, or $CE^2 + (PE + QC)^2 < CE^2 + (PE - QC)^2 + QC^2 + 2QC\sqrt{CE^2 + (PE - QC)^2}$, or $(4PE - QC)^2 < 4(CE^2 + PE^2 - 2PEQC + QC^2)$, or $4CE^2 > 12PE^2 - 3QC^2$. That is certainly true since we have $CE > (PE + QC)\sqrt{3}$.

Finally, in case (3) the only candidates are K and Q . So as in the previous paragraph we wish to show that $4CE^2 < 12PE^2 - 3QC^2$. Now we have $CE < \sqrt{3}(PE - QC)$ from which the inequality follows.

Problem A5

The real polynomial $p(x)$ is such that for any real polynomial $q(x)$, we have $p(q(x)) = q(p(x))$. Find $p(x)$.

Solution

Take $q(x) = x + k$ and set $x = 0$. Then we have $p(k) = p(0) + k$. This is true for all k , so $p(x)$ must be $x + c$ for some c . Now take $q(x) = x^2$. We get $x^2 + c = (x + c)^2 = x^2 + 2cx + c^2$. Hence $c = 0$ and $p(x) = x$.

Problem A6

A player throws a fair die (prob $1/6$ for each of $1, 2, 3, 4, 5, 6$ and each throw independent) repeatedly until his total score $\geq n$. Let $p(n)$ be the probability that his final score is n . Find $\lim p(n)$.

Solution

Answer: $2/7$.

For $i = 1, 2, 3, 4, 5$, let $p_i(n) = \text{prob that the final score is } n+i \text{ if the player stops when his total score is at least } n$. We note that $p(n)$ is also the probability that the player's total equals n on some throw if he throws repeatedly. Now we can see that $p_5(n) = 1/6 p(n-1)$, because the only way to achieve a final score of $n+5$ without passing through $n, n+1, n+2, n+3, n+4$ is to reach $n-1$ and then throw a 6 . Similarly, $p_4(n) = 1/6 p(n-1) + 1/6 p(n-2)$, because to reach $n+4$ without passing through $n, n+1, n+2, n+3$ you must either go through $n-1$, which requires reaching $n-1$ and then throwing a 5 , or not, in which case you must reach $n-2$ and then throw a 6 . Similarly, $p_3(n) = 1/6 p(n-1) + 1/6 p(n-2) + 1/6 p(n-3)$, $p_2(n) = 1/6 p(n-1) + 1/6 p(n-2) + 1/6 p(n-3) + 1/6 p(n-4)$, and $p_1(n) = 1/6 p(n-1) + 1/6 p(n-2) + 1/6 p(n-3) + 1/6 p(n-4) + 1/6 p(n-5)$. Adding, we get: $1 = p(n) + p_1(n) + p_2(n) + p_3(n) + p_4(n) + p_5(n) = p(n) + 5/6 p(n-1) + 4/6 p(n-2) + 3/6 p(n-3) + 2/6 p(n-4) + 1/6 p(n-5)$.

In the limit, $p(n-5) = p(n-4) = \dots = p(n)$. Hence we get: $p_1(n) = 5/6 p(n)$, $p_2(n) = 4/6 p(n)$, $p_3(n) = 3/6 p(n)$, $p_4(n) = 2/6 p(n)$, $p_5(n) = 1/6 p(n)$. But they sum to 1, so $p(n) = 2/7$.

We have (as above) $p(n) = 1/6 p(n-1) + 1/6 p(n-2) + 1/6 p(n-3) + 1/6 p(n-4) + 1/6 p(n-5) + 1/6 p(n-6)$ (*). Let $m(n) = \min\{p(n-1), p(n-2), p(n-3), p(n-4), p(n-5), p(n-6)\}$. Then (*) establishes that $p(n) \geq m(n)$, and so $m(n+1) \geq m(n)$. Similarly, let $M(n) = \max\{p(n-1), p(n-2), p(n-3), p(n-4), p(n-5), p(n-6)\}$. Then (*) shows that $p(n) \leq M(n)$, so $M(n+1) \leq M(n)$. Thus $m(n)$ is a monotonic increasing sequence and $M(n)$ is a monotonic decreasing sequence. But $m(n)$ is obviously bounded above by any $M(m)$, and $M(n)$ is bounded below by any $m(m)$. So both sequences converge. Suppose they converged to different limits. So $m(n)$ converges to m and $M(n)$ converges to M with $M - m > 36k > 0$. Take n sufficiently large that $m(n) > m - 6k$. At least one of the terms on the rhs of (*) must equal $M(n)$ and the others are at least $m(n)$, so $p(n) \geq 5/6 m(n) + 1/6 M(n) > 5/6 (m - 6k) + 1/6 M > 5/6 m - 5k + 1/6 (m + 36k) = m + k$. But that means that $m(n) > m + k$ for all sufficiently large n . Contradiction. Hence M and m are the same and $p(n)$ must have the same limit.

Problem A7

Let $f(n)$ be the smallest integer such that any permutation on n elements, repeated $f(n)$ times, gives the identity. Show that $f(n) = p f(n-1)$ if n is a power of p , and $f(n) = f(n-1)$ if n is not a prime power.

Solution

A permutation on n elements can be written as a product of disjoint cycles of length $\leq n$. So $f(n)$ is the smallest number divisible by 2, 3, ..., n .

If n is not a power of a prime, then we can write $n = rs$, with r and s relatively prime and each greater than 1. Then $r, s < n - 1$, so r and s divide $f(n-1)$ and hence n divides $f(n-1)$. So $f(n) = f(n-1)$. If $n = p^{m+1}$, then p^m divides $f(n-1)$, but not p^{m+1} , so $f(n) = p f(n)$.

Problem B1

Find all pairs of unequal integers m, n such that $m^n = n^m$.

Solution

Answer: (2, 4), (4, 2), (-2, -4), (-4, -2).

Suppose first that m and n are both positive. Assume $m > n$. Then we can put $m = n + k$ with $k > 0$. Hence $(1 + k/n)^n = n^k$. But for $x > 1$ we have $1 + x < e^x$ (the derivative of $f(x) = e^x - x - 1$ is positive and $f(0) = 0$) and hence $(1 + k/n)^n < e^k$. So there are no solutions for $n > 2$. If $n = 1$, then $n^m = 1$ and hence $m = 1$, contradicting the fact that m and n are unequal. If $n = 2$, then m must be a power of 2. Suppose $m = 2^h$. Then we find $h = 1$ or 2. $h = 1$ is invalid (because m and n are unequal), so $m = 4$. There is also the corresponding solution with $m < n$.

If $n < 0$ and $m > 0$, then $n^m = 1/m^{-n}$. So m divides 1 and hence $m = 1$. But m must be even for to make n^m positive. Contradiction. So there are no solutions of this type. If m and n are both negative, then $-m, -n$ is a solution, so the only possibilities are (-2, -4) and (-4, -2) and it is readily checked that these are indeed solutions.

Problem B2

Let $f(m, n) = 3m+n+(m+n)^2$. Find $\sum_0^\infty \sum_0^\infty 2^{-f(m, n)}$.

Solution

Answer: 4/3.

The obvious approach is to sum over n and then over m (or vice versa). But it is not obvious how to sum x^N where N runs through the squares. So we wonder just what values $f(m, n)$ can take. A little experimentation suggests that it takes each even value just once.

In fact, it is clear that $r(r+1)$ is always even and that the difference between $r(r+1)$ and the next number in the sequence, $(r+1)(r+2)$ is $2r+2$, so any positive even number can be uniquely expressed as $r(r+1) + 2s$ with $0 \leq s \leq r$. But taking $m = s$, $n = r-s$, this is equivalent to the statement that any positive even number can be uniquely expressed as $3m+n+(m+n)^2$.

Hence the sum is just $1 + 1/4 + 1/4^2 + \dots = 4/3$.

Problem B3

Fluid flowing in the plane has the velocity $(y + 2x - 2x^3 - 2xy^2, -x)$ at (x, y) . Sketch the flow lines near the origin. What happens to an individual particle as $t \rightarrow \infty$?

Solution

The key insight is that the unit circle centre the origin is one possible trajectory. [The velocity is $(y + 2x - 2xr^2, -x) = (y, -x)$ on $r = 1$.] Hence fluid starting inside this circle remains inside it and fluid starting outside it remains outside. Differentiating $r^2 = x^2 + y^2$ gives $r \, dr/dt = 2x^2(1 - r^2)$. Hence if $r < 1$, then dr/dt is positive, so points inside the unit circle move outwards towards it. Similarly, if $r > 1$, then dr/dt is negative, so points outside the unit circle move inwards towards it.

To find the behaviour close to the origin, we drop the cubic terms. The equations are then linear and can be solved to get $x = (at + b)e^t$, $y = (-at + a - b)e^t$. This gives S shaped curves centred on the origin at a 45° angle touching the line $y = -x$.

Problem B4

Show that if an (infinite) arithmetic progression of positive integers contains an n th power, then it contains infinitely many n th powers.

Solution

Let the difference between adjacent terms of the progression be d . Suppose that N^n is an n th power in the progression. Then $(N+d)^n$ is a larger n th power in the progression (it is obvious from the binomial expansion that it has the form $N + kd$).

Problem B5

Define a_n by $a_0 = 0$, $a_{n+1} = 1 + \sin(a_n - 1)$. Find $\lim (\sum_0^n a_i)/n$.

Solution

Answer: 1.

Note that $1 - a_{n+1} = \sin(1 - a_n)$. Put $c_n = 1 - a_n$. Note that c_0 belongs to the interval $(0, 1]$. Now for x in $(0, 1]$ we have that $\sin x < x$ and $\sin x$ is also in $(0, 1]$. So it follows (by a trivial induction) that $1 = c_0 > c_1 > c_2 > \dots > c_n > 0$. So c_n is a monotonically decreasing sequence bounded below by 0. Hence it must tend to a limit $c \geq 0$. But c must satisfy $c = \sin c$. Hence $c = 0$. Hence a_n converges to 1. Hence $(\sum_0^n a_i)/n$ converges to 1 also.

Problem B6

Let $2^{f(n)}$ be the highest power of 2 dividing n . Let $g(n) = f(1) + f(2) + \dots + f(n)$. Prove that $\sum \exp(-g(n))$ converges.

Solution

If n is odd, then $f(n) = 0$. If n is even, then $f(n) \geq 1$. Hence $g(n+2) \geq g(n) + 1$. So the series is dominated by $1 + (1/e + 1/e) + (1/e^2 + 1/e^2) + \dots < 2/(1 - 1/e)$. So the sums are bounded above. All terms are positive, so it must converge.

Problem B7

Let R' be the non-negative reals. Let $f, g : R' \rightarrow R$ be continuous. Let $a : R' \rightarrow R$ be the solution of the differential equation: $a' + f a = g$, $a(0) = c$. Show that if $b : R' \rightarrow R$ satisfies $b' + f b \geq g$ for all x and $b(0) = c$, then $b(x) \geq a(x)$ for all x . Show that for sufficiently small x the solution of $y' + f y = y^2$, $y(0) = d$, is $y(x) = \max (d e^{-h(x)} - \int_0^x e^{-(h(x)-h(t))} u(t)^2 dt)$, where the maximum is taken over all continuous $u(t)$, and $h(t) = \int_0^t (f(s) - 2u(s)) ds$.

Solution

Put $F(x) = \int_0^x f(t) dt$. Put $A(x) = a(x) e^{F(x)}$, $B(x) = b(x) e^{F(x)}$. Then $A' = (a' + f a) e^F = g e^F$, and $B' = (b' + f b) e^F \geq g e^F$, since $e^F > 0$. Hence $B' \geq A'$ for all x . But $B(0) = A(0)$, so $B(x) \geq A(x)$ for all x and hence $b(x) \geq a(x)$ for all x . Let u be any continuous function on R' . Put $U(x) = \int_0^x u(t) dt$. Put $h(x) = F(x) - 2U(x) = \int_0^x f(t) - 2u(t) dt$. Then $h(0) = 0$. We have $(y - u)^2 \geq 0$, so $y^2 - 2uy \geq -u^2$. Now put $z = y e^h$, so that $z(0) = y(0) e^{h(0)} = d$.

$z' = (y' + h'y) e^h = (y' + (f - 2u)y) e^h = (y^2 - 2uy) e^h \geq -u^2 e^h$. Hence $z(x) = z(0) + \int_0^x z'(t) dt \geq d - \int_0^x u(t)^2 e^{h(t)} dt$. Hence $y(x) = z(x) e^{-h(x)} \geq d e^{-h(x)} - \int_0^x e^{-(h(x)-h(t))} u(t)^2 dt$. So $z(x) \geq$ the maximum over all u .

If we put $u = y$ (the solution), then we get equality.

22nd Putnam 1961

Problem A1

The set of pairs of positive reals (x, y) such that $x^y = y^x$ form the straight line $y = x$ and a curve. Find the point at which the curve cuts the line.

Solution

Put $y = kx$. Then we get $x = k^{1/(k-1)}$, $y = k^{k/(k-1)}$. This gives a curve which cuts $y = x$ at $x = \lim k^{1/(k-1)}$, where the limit is taken as k tends to 1. Put $k = 1 + 1/n$, then the limit is $(1 + 1/n)^n$, which is e . So the point of intersection is (e, e) .

Problem A2

For which real numbers α, β can we find a constant k such that $x^\alpha y^\beta < k(x + y)$ for all positive x, y ?

Solution

Answer: $\alpha, \beta \geq 0$ and $\alpha + \beta = 1$.

If $\alpha < 0$, then taking x sufficiently small and $y = 1$ gives $x^\alpha y^\beta > k(x + y)$ for any k . Similarly for $\beta < 0$. If $\alpha + \beta > 1$, then taking $x = y$ sufficiently large gives $x^\alpha y^\beta > k(x + y)$ for any k . On the other hand, if $\alpha + \beta < 1$, then taking $x = y$ sufficiently small gives $x^\alpha y^\beta > k(x + y)$ for any k . Suppose, on the other hand that $\alpha, \beta \geq 0$ and $\alpha + \beta = 1$. Take $x \geq y$. Then $x^\alpha y^\beta \leq x^{\alpha+\beta} = x$, so the result holds with $k = 1$.

Problem A3

Find $\lim_{n \rightarrow \infty} \sum_{i=1}^n n/(N + i^2)$, where $N = n^2$.

Solution

As usual, we try integration. We can write the sum as $1/n \sum_{i=1}^n 1/(1 + (i/n)^2)$. This is a Riemann sum for the integral $\int_0^1 1/(1 + x^2) dx$ and hence tends to $\int_0^\infty 1/(1 + x^2) dx = \tan^{-1} x \Big|_0^\infty = \pi/2$.

Problem A4

If $n = \prod p^r$ be the prime factorization of n , let $f(n) = (-1)^{\sum r}$ and let $F(n) = \sum_{d|n} f(d)$. Show that $F(n) = 0$ or 1 . For which n is $F(n) = 1$?

Solution

Answer: $F(n) = 1$ for n square.

If some r is odd, then the factors of n can be grouped into pairs $p^a m, p^{r-a} m$ with $f(p^a m), f(p^{r-a} m)$ having opposite signs. So in this case $F(n) = 0$.

If all r are even, so that n is a square, then we may use induction on the number N of prime factors. For $N = 1$, we have $n = p^{2m}$, so there are $m+1$ factors p^{2r} with $f = 1$ and m factors p^{2r+1} with $f = -1$. So the result is true for $N = 1$. Suppose it is true for N . Take $n = p^{2m} P$, where P has m prime factors. Each divisor of n has the form $p^{2r} s$ or $p^{2r+1} s$, where s is a divisor of P . For the first type $f(p^{2r} s) = f(s)$, so summing over s gives 1. Then summing over r gives $m+1$. For the second type $f(p^{2r+1} s) = -f(s)$, so summing over s gives -1 . Then summing over r gives $-m$. Hence $F(n) = 1$ and the result is true for $N+1$.

Problem A5

Let X be a set of n points. Let P be a set of subsets of X , such that if $A, B \in P$, then $X - A, A \cup B, A \cap B \in P$. What are the possible values for the number of elements of P ?

Solution

Answer: $2, 4, 8, \dots, 2^n$.

Take $0 \leq r < n$. Let $A_1 = \{1\}, A_2 = \{2\}, \dots, A_r = \{r\}, A_{r+1} = \{r+1, r+2, \dots, n\}$. Consider the collection of all unions of sets A_i . This collection has 2^{r+1} elements and satisfies the conditions. So this shows that we can achieve the values $2, 4, 8, \dots, 2^n$.

To prove the converse we use induction on n . It is obviously true for $n = 1$. Suppose it is true for all $n < N$. Let X have N points and P be a set of subsets of X satisfying the condition. If $|P| = 2$, then there is nothing to prove. Otherwise we can find a member Y of P other than the empty set and X which has no non-trivial subsets in P . Now consider the collection Q of subsets of $X - Y$ which are in P . It is easy to check that (1) P is just the collection of all A and $A \cap Y$, where A is in Q , so that $|P| = 2|Q|$, and (2) Q satisfies the conditions, so $|Q|$ is a power of 2 by induction.

Problem A6

Consider polynomials in one variable over the finite field F_2 with 2 elements. Show that if $n + 1$ is not prime, then $1 + x + x^2 + \dots + x^n$ is reducible. Can it be reducible if $n + 1$ is prime?

Solution

Let $n+1 = ab$, then $1 + x + x^2 + \dots + x^n = (1 + x + x^2 + \dots + x^{a-1})(1 + x^a + x^{2a} + \dots + x^{ab-a})$. Note that this does not depend upon the field having two elements.

Yes. For example, $(1 + x + x^3)(1 + x^2 + x^3) = 1 + x + x^2 + x^3 + x^4 + x^5 + x^6$.

Problem A7

S is a non-empty closed subset of the plane. The disk (a circle and its interior) $D \supseteq S$ and if any disk $D' \supseteq S$, then $D' \supseteq D$. Show that if P belongs to the interior of D , then we can find two distinct points $Q, R \in S$ such that P is the midpoint of QR .

Solution

The existence of D imposes strong restrictions on S . In fact, we will show that S must contain the entire perimeter C of D . For suppose it does not contain a point X on C . Then since S is closed, it does not contain a neighbourhood of X . So we can find a small circle C' centre X so that the disk D' enclosed by C' lies entirely outside S . Let C' meet C at A and B . Take a circle C'' through A and B with centre on the same side of AB as the centre of C , but further away (so that it has a larger radius than C). Then C'' contains all points of $D - D'$ and hence all points of S . But it does not contain all points of D . Contradiction.

Finally, note that any point in the interior of D lies on the midpoint of some chord of C .

Problem B1

a_n is a sequence of positive reals. $h = \lim (a_1 + a_2 + \dots + a_n)/n$ and $k = \lim (1/a_1 + 1/a_2 + \dots + 1/a_n)/n$ exist. Show that $h k \geq 1$.

Solution

Apply the arithmetic-geometric mean theorem to each sum. We get that $(a_1 + a_2 + \dots + a_n)/n \geq (a_1 a_2 \dots a_n)^{1/n}$, $(1/a_1 + 1/a_2 + \dots + 1/a_n)/n = 1/(a_1 a_2 \dots a_n)^{1/n}$. Hence their product is at least 1. Hence the product of the limits also.

Problem B2

Two points are selected independently and at random from a segment length β . What is the probability that they are at least a distance α ($< \beta$) apart?

Solution

Consider which points (x, y) of the square $x = 0$ to β , $y = 0$ to β have $|x - y| \geq \alpha$. Evidently the acceptable points lie in the two right-angled triangles: $(0, \alpha)$, $(0, \beta)$, $(\beta - \alpha, \beta)$ and $(\alpha, 0)$, $(\beta, 0)$, $(\beta, \beta - \alpha)$. These fit together to give a square side $\beta - \alpha$, so the area of the acceptable points is $(\beta - \alpha)^2$ out of a total area of β^2 . Thus the probability is $(\beta - \alpha)^2 / \beta^2$.

Problem B3

A, B, C, D lie in a plane. No three are collinear and the four points do not lie on a circle. Show that one point lies inside the circle through the other three.

Solution

If the convex hull is a triangle, then the fourth point lies inside its circumcircle. So suppose $ABCD$ is convex. One pair of opposite angles must have sum greater than 180° (otherwise the points would lie on a circle). Suppose they are A and C . Then we claim that C lies inside the circle through A, B, D . Take a point C' on the ray DC on the far

side of C from D such that $\angle CBC' = \angle A + \angle C - 180^\circ$. Then $\angle C' = 180^\circ - \angle A$. So C' lies on the circle. Hence C which lies on the segment C'D lies inside the circle.

Problem B4

Given $x_1, x_2, \dots, x_n \in [0, 1]$, let $s = \sum_{1 \leq i < j \leq n} |x_i - x_j|$. Find $f(n)$, the maximum value of s over all possible $\{x_i\}$.

Solution

Answer: $f(2m) = m^2$ (half the points 0, half 1); $f(2m+1) = m^2 + m$ ($m+1$ points 0, m points 1).

It remains to prove that we cannot do better than this. We prove that this is best possible by induction. It is obvious for $n = 2, 3$. Suppose it is true for n . Take $n+2$ points. Let A be the leftmost point and B the rightmost point. Then the new terms in the sum the $n+2$ points are AB and AC, CB for each existing C. But these $2n + 1$ terms sum to at most $n + 1$ ($AC + CB = AB \leq 1$). So $f(n+2) \leq f(n) + n + 1$. For $n = 2m$ this gives $f(2m+2) \leq m^2 + 2m + 1 = (m + 1)^2$. For $n = 2m + 1$, it gives $f(2m+3) \leq m^2 + m + 2m + 1 + 1 = (m+1)^2 + (m+1)$.

Problem B5

Let n be an integer greater than 2. Define the sequence a_m by $a_1 = n$, $a_{m+1} = n$ to the power of a_m . *Either* show that $a_m < n!! \dots!$ (where the factorial is taken m times), *or* show that $a_m > n!! \dots!$ (where the factorial is taken $m-1$ times).

Solution

If it was true that $n^k < k!$, then the upper limit on a_m would be almost trivial. Unfortunately, $n^k > k!$ for small k , which results in a few complications. Let $n_m = n!! \dots!$ (m times). We will prove that $a_m < n_m$ by induction on m . For $m = 1$, the result just states that $n < n!$, which is obviously true. Suppose now that $n \geq 5$. Then $n! \geq n(n-1)3 > 2n^2$. So, a fortiori, $n_m > 2n^2$ for $m \geq 1$. It is easy to check that for $n = 3$, $a_2 = 27 < n_2 = 3!! = 720$, so the result is true for $m = 2$. Also $n_2 > 2n^2 = 18$. Similarly, for $n = 4$, $a_2 = 256 < n_2 = 4!! = 24!$ which establishes the result for $m = 2$. Also $n_2 > 2n^2 = 32$. So for all n we have a starting case m for which also $n_m > 2n^2$.

If $k \geq 2n^2$, then half the terms in $k!$ are at least n^2 , so their product is at least $n^{2k/2} = n^k$. In other words $k! > n^k$. So suppose that $a_m < n_m$, (*). Then since $n_m > 2n^2$, we may take (*) to the power of n to get $a_{m+1} = n^{a_m} < n^{n_m} < n_{m+1}$. Hence the result is true for all m .

The lower limit is harder because it is not true that $n^k > k!$, so the result for $m+1$ does not follow in an immediate way from that for m .

Let $f_0(n) = n$, $f_1(n) = f_0(n) n!$, $f_2(n) = f_1(n) n!!$, $f_3(n) = f_2(n) n!!!$, We show that $f_{m+1}(n) < n$ to the power of $f_m(n)$. We use induction on m . For $m = 0$, this reads $n n! < n^n$, which is obviously true. Now assume it is true for m **and all** n . Then we have $f_{m+1}(n) = n f_m(n!) < n (n! \text{ to the power of } f_{m-1}(n!)) < (n n!) \text{ to the power of } f_{m-1}(n!) < n^n \text{ to the power of } f_{m-1}(n!) = n$ to the power of $f_m(n)$.

It is now an easy induction that $f_m(n) < a_{m+1}$. For $m = 1$, this just states that $n n! < n^n$. Suppose it is true for m , then $f_{m+1}(n) < n$ to the power of $f_m(n) < n$ to the power of $a_{m+1} = a_{m+2}$. But obviously $n! \dots!$ (! taken m times) $< f_m(n)$ and hence $< a_{m+1}$.

Problem B6

Let y be the solution of the differential equation $y'' = -(1 + \sqrt{x})y$ such that $y(0) = 1$, $y'(0) = 0$. Show that y has exactly one zero for $x \in (0, \pi/2)$ and find a positive lower bound for it.

Solution

Let z be the solution of $z'' = -z$ with $z(0) = 1$, $z'(0) = 0$. Suppose $y > 0$ for $0 \leq x < \pi/2$. Then since $y''z - yz'' = -(1 + \sqrt{x})yz + yz = -\sqrt{x}yz$, we have $\int_0^{\pi/2} -\sqrt{x}yz \, dx = \int_0^{\pi/2} (y''z - yz'') \, dx = (y'z - yz') \Big|_0^{\pi/2} = y(\pi/2) \geq 0$. But $-\sqrt{x}yz < 0$ throughout the range, so $\int_0^{\pi/2} -\sqrt{x}yz \, dx < 0$. Contradiction. Hence y has at least one zero in $(0, \pi/2)$.

Let w be the solution of $w'' = -3w$ with $w(0) = 1$, $w'(0) = 0$. Then $w = \cos(\sqrt{3}x)$. Suppose y had a zero in $(0, \pi/(2\sqrt{3}))$. Then the same argument would show that w had a root in $(0, \pi/(2\sqrt{3}))$ which is false. So $\pi/(2\sqrt{3})$ is a positive lower bound for any zero of y .

Finally, suppose that y has more than one zero in $(0, \pi/2)$. Let the first be at $x = h$ and the second at $x = k$. Take $v(x)$ to be $A \cos(\sqrt{3}x + B)$. We select A and B so that $v(h) = 0$ and $v'(h) = y'(h)$. Now v cannot have another zero in (h, k) , so we can establish a contradiction by a similar argument to the above. We have $v'' = -3v$. Hence $y''v - yv'' = (-(1 + \sqrt{x}) + 3)yv$ which is strictly positive on (h, k) . Hence $\int_h^k (y''v - yv'') dx > 0$. But $\int_h^k (y''v - yv'') dx = (y'v - yv')|_h^k = y'(k)v(k) < 0$. [y' must be positive since y is zero and $y(x) < 0$ for x just less than k , whilst we know that $v(k)$ is negative]. Contradiction. Hence y has exactly one zero.

Problem B7

The sequence of non-negative reals satisfies $a_{n+m} \leq a_n a_m$ for all m, n . Show that $\lim a_n^{1/n}$ exists.

Solution

$a_n \leq a_1 a_{n-1} \leq a_1^2 a_{n-2} \leq \dots \leq a_1^n$. So $a_n^{1/n} \leq a_1$. All a_n are non-negative, so $a_n^{1/n} \geq 0$. Thus we have established that $\{a_n^{1/n}\}$ is bounded.

Fix n . Take any $N > n$. Then we may write $N = nq + r$, with $0 \leq r < n$. We have $a_N \leq a_n^q a_r$, so $a_N^{1/N} \leq a_n^{q/N} a_r^{1/N}$, where $s = q/N = (1 - r/N)/n$. But $(1 - r/N)$ tends to 1 as N tends to infinity, as does $a_r^{1/N}$. Hence $\limsup a_N^{1/N} \leq a_n^{1/n}$. This is true for all n , so the sequence cannot have more than one limit point and hence converges.

23rd Putnam 1962

Problem A1

5 points lie in a plane, no 3 collinear. Show that 4 of the points form a convex quadrilateral.

Solution

If the convex hull has 4 or 5 vertices, then we are done. If not, then two of the points, say D and E, must lie inside the triangle ABC formed by the other three. DE must intersect just two of AB, BC and CA. Suppose that it does not intersect AB. Then the four points A, B, D, E form a convex quadrilateral.

Problem A2

Let \mathbb{R} be the reals. Find all $f : K \rightarrow \mathbb{R}$, where K is $[0, \infty)$ or a finite interval $[0, a)$, such that $(1/k \int_0^k f(x) dx)^2 = f(0) f(k)$ for all k in K .

Solution

Answer: for $[0, \infty)$ $f(x) = b/(cx + 1)^2$ with c non-negative. For $[0, a)$, $f(x) = b/(cx + 1)^2$ with $c \geq -1/a$.

Put $y = \int_0^x f(t) dt$. Then the given equation becomes $y^2/x^2 = by'$, where $b = f(0)$. Integrating, $b/y = c + 1/x$, or $y = bx/(cx + 1)$. Differentiating gives $f(x) = y' = b/(cx + 1)^2$.

$f(-1/c)$ is undefined, so if K is the half-line, then c cannot be negative. If K is the finite interval $[0, a)$, then we require $c \geq -1/a$. Note that we can have equality because k must be strictly less than a . On the other hand, b can have any value.

Problem A3

ABC is a triangle and $k > 0$. Take A' on BC, B' on CA, C' on AB so that $AB' = k B'C$, $CA' = k A'B$, $BC' = k C'A$. Let the three points of intersection of AA' , BB' , CC' be P, Q, R. Show that the area PQR $(k^2 + k + 1) = \text{area ABC} (k - 1)^2$.

Solution

There does not seem to be a geometric solution, so one is faced with a messy algebraic calculation. There are various ways of doing it. The worst is probably to use ordinary Cartesian coordinates.

Let us use points in the Argand diagram. Take A to be the origin and C to be the real point $1+k$. Take B to be $z(1+k)$. Then B' is k and A' is $zk+1$ (check: $((zk+1) - (1+k)) / ((z(1+k)) - (zk+1)) = (zk-k)/(z-1) = k$). We want to find R, the intersection of AA' and BB' . It lies on AA' , so it is $r(zk+1)$ for some $0 < r < 1$. But it lies on BB' , so it is $k + s(zk+z-k)$ for some $0 < s < 1$. After some manipulation, we get $r = (k^2+k)/(k^2+k+1)$.

Now take $k < 1$. Then $\text{area ARB} = (AR/AA') \text{area AA'B} = r \text{area AA'B}$ (this is where we need $k < 1$, because if $k > 1$, then the geometry changes). But $\text{area AA'B} = (A'B/CB) \text{area ABC}$, so $\text{area ARB} = r/(1+k) \text{area ABC} = k/(k^2+k+1) \text{area ABC}$. Similarly, taking P as the intersection of BB' and CC' and Q as the intersection of AA' and CC' , we have that $\text{area AQC} = k/(k^2+k+1) \text{area ABC}$, and $\text{area CPB} = k/(k^2+k+1) \text{area ABC}$. Adding we get $(\text{area ABC} - \text{area PQR})$, so $\text{area PQR}/\text{area ABC} = 1 - 3k/(k^2+k+1) = (1 - k)^2/(k^2+k+1)$.

If $k > 1$, then we get the result with $1/k$ replacing k , but multiplying through by k^2 gives the same formula.

Problem A4

\mathbb{R} is the reals. $[a, b]$ is an interval with $b \geq a + 2$. $f : [a, b] \rightarrow \mathbb{R}$ is twice differentiable and $|f(x)| \leq 1$ and $|f''(x)| \leq 1$. Show that $|f'(x)| \leq 2$.

Solution

Take the interval to be $[-1, 1]$. Taylor's formula gives $f(1) = f(x) + (1-x)f'(x) + (1-x)^2 f''(h)/2$, $f(-1) = f(x) + (-1-x)f'(x) + (-1-x)^2 f''(k)/2$ for some h, k in the interval (note that we are expanding about x).

Subtracting, $f(1) - f(-1) = 2f'(x) + (1-x)^2 f''(h)/2 - (1+x)^2 f''(k)/2$. Hence $2|f'(x)| \leq |f(1)| + |f(-1)| + (1-x)^2 |f''(h)|/2 + (1+x)^2 |f''(k)|/2 \leq 2 + (1-x)^2/2 + (1+x)^2/2 = 3 + x^2 \leq 4$. So $|f'(x)| \leq 2$.

Problem A5

Find $nC_1 1^2 + nC_2 2^2 + nC_3 3^2 + \dots + nC_n n^2$ (where nC_r is the binomial coefficient).

Solution

Answer: $n(n+1)2^{n-2}$.

Differentiate $(1+x)^n$, multiply by x and differentiate again. Set $x = 1$.

Problem A6

X is a subset of the rationals which is closed under addition and multiplication. $0 \notin X$. For any rational $x \neq 0$, just one of $x, -x \in X$. Show that X is the set of all positive rationals.

Solution

Either x or $-x$ belongs to X . X is closed under multiplication, so the square $x^2 = (-x)^2$ belongs to X . In particular, 1 belongs to X . Hence by repeated addition all positive integers must belong to X . Suppose positive rational m/n does not belong to X . Then $-m/n$ does, and hence by repeated addition $-m$. So m does not belong to X . Contradiction. So X contains all positive rationals. But if x is in X , $-x$ is not, so X does not contain any negative rationals and hence X is just the set of all positive rationals.

Problem B1

Define $x^{(n)} = x(x-1)(x-2)\dots(x-n+1)$ and $x^{(0)} = 1$. Show that $(x+y)^{(n)} = nC_0 x^{(0)}y^{(n)} + nC_1 x^{(1)}y^{(n-1)} + nC_2 x^{(2)}y^{(n-2)} + \dots + nC_n x^{(n)}y^{(0)}$.

Solution

Induction on n . It is obviously true for $n = 1$. Suppose it is true for n . We now multiply the rhs by $(x+y-n)$. We now write:

$$\begin{aligned} nC_0 x^{(0)}y^{(n)}(x+y-n) &= (nC_0 x^{(0)}y^{(n)}(y-n)) + (nC_0 x^{(0)}y^{(n)}x); \\ nC_1 x^{(1)}y^{(n-1)}(x+y-n) &= (nC_1 x^{(1)}y^{(n-1)}(y-n+1)) + (nC_1 x^{(1)}y^{(n-1)}(x-1)); \\ nC_2 x^{(2)}y^{(n-2)}(x+y-n) &= (nC_2 x^{(2)}y^{(n-2)}(y-n+2)) + (nC_2 x^{(2)}y^{(n-2)}(x-2)); \\ &\dots \\ nC_n x^{(n)}y^{(0)}(x+y-n) &= (nC_n x^{(n)}y^{(0)}y) + (nC_n x^{(n)}y^{(0)}(x-n)). \end{aligned}$$

Now the first term in the 1st equation gives $(n+1)C_0 x^{(0)}y^{(n+1)}$. The second term in the 1st equation and the first term in the 2nd equation give $(n+1)C_1 x^{(1)}y^{(n)}$. Then the second term in the 2nd equation and the first in the 3rd equation gives $(n+1)C_2 x^{(2)}y^{(n-1)}$ and so on. Finally the second term in the last equation gives $(n+1)C(n+1) x^{(n+1)}y^{(0)}$. So we have the result for $n+1$.

Problem B2

Let R be the reals, let N be the set of positive integers, and let $P = \{X : X \subseteq N\}$. Find $f : R \rightarrow P$ such that $f(a) \subset f(b)$ (and $f(a) \neq f(b)$) if $a < b$.

Solution

Let Q be the rationals. Define $g : Q \rightarrow N$ as follows. For m/n where m and n are positive integers without any common factor, let $g(m/n) = 2^m 3^n$ and let $g(-m/n) = 2^m 5^n$. Now define $f : R \rightarrow P$ by $f(x) = \{g(r) : r \leq x \text{ is rational}\}$. We claim that f has the required property.

Clearly g is injective. Now suppose $x < y$ are reals. We can find a rational r such that $x < r < y$. So $g(r)$ belongs to $f(y)$ but not to $f(x)$. But S belongs to $f(x)$, then $S = g(q)$ for some q in Q with $q < x$. So $q < y$ and hence S belongs to $f(y)$ also. So $f(x)$ is a subset of $f(y)$.

Problem B3

Show that a convex open set in the plane containing the point P , but not containing any ray from P , must be bounded. Is this true for any convex set in the plane?

Solution

Answer: no.

Let S be the convex set. Take any ray R from P . We can find X on R not in S . Since S is open we can take a small segment AB perpendicular to R through P with $AP = PB$ which is entirely contained in S . Take A' on the line AX ,

the opposite side to A, and B' on BX the opposite side to B, with A'B' perpendicular to R. Then no point C' on the segment A'B' can be in S. For if it was, then we could project C'X to meet the segment AB in C and X would lie between two points of S and hence would have to be in S. But the angle A'PB' is greater than zero. So we have found an open interval about the direction PX (measured as a polar angle at P) such that S extends at most a distance AA' for directions within that interval.

The interval $[0, 2\pi]$ is compact and is covered by these open intervals, so a finite number of them cover it. Hence there is a finite distance d such that we can find a point not in S within a distance d of P in *any* direction. That means that no point Q a distance greater than d from P can be in S (otherwise all points on the segment QP would be in S and there would be no point not in S within a distance d in that direction). So S lies inside the disk radius d on P and hence is bounded.

Take the slab $0 < y < 1$ together with the point $(0, 0)$. It is convex and unbounded but does not contain any ray through the origin.

Problem B4

A finite set of circles divides the plane into regions. Show that we can color the plane with two colors so that no two adjacent regions (with a common arc of non-zero length forming part of each region's boundary) have the same color.

Solution

Color points inside an odd number of circles blue and points inside an even number of circles red. Then we change color whenever we cross a circle.

Problem B5

Show that for $n > 1$, $(3n + 1)/(2n + 2) < \sum_1^n r^n/n^n < 2$.

Solution

Suppose $m \leq n$, then expanding by the binomial theorem, $m^n = ((m-1) + 1)^n = (m-1)^n + n(m-1)^{n-1} + \text{further positive terms} > (m-1)^n + n(m-1)^{n-1} > 2(m-1)^n$. Hence $n^n > (n-1)^n + (n-1)^n > (n-1)^n + (n-2)^n + (n-2)^n > \dots > (n-1)^n + (n-2)^n + \dots + 1^n$. So $2n^n > \sum_1^n r^n$, which is one of the two required inequalities.

The other is slightly harder. We approximate the integral of $f(x) = x^n$ between 0 and 1. We have to do slightly better than the usual Riemann sums. We also take into account the small triangles. The curve has increasing gradient, so the area under the curve between $(r-1)/n$ and r/n is less than the area under the chord joining $((r-1)/n, f((r-1)/n))$ and $(r/n, f(r/n))$. Summing those areas gives: $(1/n) \sum_1^n (r/n)^n - (1/2n) \sum_1^n ((r/n)^n - ((r-1)/n)^n) = (1/n)K - (1/2n)(n/n)^n$, where $K = \sum_1^n (r/n)^n$. The integral is just $1/(n+1)$, so we have: $1/(n+1) < K/n - 1/(2n)$, or $K > n/(n+1) + 1/2 = (3n+1)/(2n+2)$.

Problem B6

$f: [0, 2\pi] \rightarrow [-1, 1]$ satisfies $f(x) = \sum_0^n (a_j \sin jx + b_j \cos jx)$ for some real constants a_j, b_j . Also $|f(x)| = 1$ for just $2n$ distinct values in the interval. Show that $f(x) = \cos(nx + k)$ for some k .

Solution

We call $f(x)$ a trigonometric sum of degree n . Let x_1, x_2, \dots, x_{2n} be the points at which $f(x) = 1$ or -1 . Two insights are needed. The first is that $f'(x_i) = 0$. For if the value was non-zero, then $f(x)$ would lie outside $[-1, 1]$ for points sufficiently close (and on the correct side). [Note that there is a minor complication at 0. If $f(0) = 1$ and $f'(0) < 0$, then the aberrant points would be just less than 2π .]

The second is that if a trigonometric sum of degree n has $2n$ roots (counting multiplicities), then it is determined up to a multiplicative constant (and if it has more than $2n$ roots then it is identically zero). We will establish that later. But $1 - f(x)^2$ has the $2n$ roots x_i . Also, they all have multiplicity at least 2 since $f'(x_i) = 0$. So we have at least $4n$ roots (counting multiplicities). Using the familiar formulae $\cos(A + B)$ etc we can write $1 - f(x)^2$ in the same form as $f(x)$ but with the sum from 0 to $2n$.

Similarly, $f'(x)$ is a trigonometric sum of degree n with the $2n$ roots x_i . If we square it, each root is doubled, so $f'(x)^2$ is a trigonometric sum of degree $2n$ with the $4n$ roots x_i (each counted twice). Hence we must have $f'(x)^2 = \text{multiple of } (1 - f(x)^2)$. Both are non-negative, so it must be a positive multiple. Hence we can take the square root

to get $f'(x) = N \sqrt{1 - f(x)^2}$ for some fixed N . Solving, we get $f(x) = \cos(Nx + k)$. But for this to have just $2n$ values ± 1 we must have $N = n$.

To prove the result about trigonometric polynomials put $z = e^{ix}$, then $f(x) = z^{-n} p(z)$, where p is a polynomial in z of degree $2n$. Hence p has at most $2n$ roots and is determined by those roots up to a multiplicative constant. Note that if $f'(x) = 0$ and $f(x) = 0$, then $p(z) = 0$ and since $0 = f'(x) =$ combination of $p(z)$ and $p'(z)$, then $p'(z) = 0$ also, so double roots of f have corresponding roots of multiplicity at least 2 in p .

24th Putnam 1963

Problem A1

Dissect a regular 12-gon into a regular hexagon, 6 squares and 6 equilateral triangles. Let the regular 12-gon have vertices P_1, P_2, \dots, P_{12} (in that order). Show that the diagonals $P_1P_9, P_{12}P_4$ and P_2P_{11} are concurrent.

Solution

Take squares on the inside of alternate sides of the 12-gon. Each angle of the 12-gon is 150° , so the remaining angles after placing the squares are all 60° , so we have equilateral triangles on the inside of the remaining 6 sides of the 12-gon. Now the opposite sides of the squares form a hexagon. (As a check we may notice that if A, B, C, D are adjacent vertices of the 12-gon and we place squares $ABPQ$ on AB and $CDRS$ on CD , then S and P coincide and the equilateral triangle ABP and the two squares take $60^\circ + 90^\circ + 90^\circ$ out of the 360° angle at P leaving 120° , which is the angle between sides of a regular hexagon.)

Let O be the centre of the 12-gon. P_1P_9 and $P_{12}P_4$ have the same length and one is obtained from the other by rotation about O through 90° . Thus they intersect on the perpendicular bisector of $P_{12}P_1$ a distance $P_{12}P_1/2$ from $P_{12}P_1$. $P_{11}P_2$ is parallel to $P_{12}P_1$ and its midpoint lies on the perpendicular bisector of $P_{12}P_1$. The angle $P_{11}P_2P_1$ is $180^\circ - \text{angle } P_{12}P_1P_2 = 30^\circ$, so the distance between of the midpoint along the bisector is $P_{12}P_1 \sin 30^\circ = P_{12}P_1/2$. Hence the three lines intersect at this point.

Problem A2

The sequence a_1, a_2, a_3, \dots of positive integers is strictly monotonic increasing, $a_2 = 2$ and $a_{mn} = a_m a_n$ for m, n relatively prime. Show that $a_n = n$.

Solution

The key observation is that if $a_N = N$ for N odd, then $a_{2N} = 2N$, and hence $a_{N+1} = N+1, a_{N+2} = N+2, \dots, a_{2N-1} = 2N-1$ (because a_n is strictly increasing, so there are only $N-1$ holes for $N-1$ pegs. But we can now repeat using $2N-1$ (provided $2N-1 > N$, or $N > 1$) and the result is established for all n .

So we just need to get started with a_3 . Suppose $a_3 = m$. Then $a_6 = 2m$. So $m+2 \leq a_5 \leq 2m-1$. Hence $a_{10} \leq 4m-2$ and $a_9 \leq 4m-3$. So $a_{18} \leq 8m-6$ and hence $a_{15} \leq 8m-9$. But $a_{15} = a_3 a_5 \geq m^2 + 2m$, so $m^2 + 2m \leq 8m - 9$ or $(m-3)^2 \leq 0$. Hence $m = 3$.

Problem A3

Let D be the differential operator d/dx and E the differential operator $xD(xD - 1)(xD - 2) \dots (xD - n)$. Find an expression of the form $y = \int_1^x g(t) dt$ for the solution of the differential equation $Ey = f(x)$, with initial conditions $y(1) = y'(1) = \dots = y^{(n)}(1) = 0$, where $f(x)$ is a continuous real-valued function on the reals.

Solution

The first step is to try to simplify Ey . Experimenting with small n , we soon suspect that $Ey = x^{n+1}y^{(n+1)}$ and that is easily proved by induction.

So we seek a solution to $x^{n+1}y^{(n+1)} = f(x)$, subject to $y(1) = y'(1) = \dots = y^{(n)}(1) = 0$. The trick is to use the Taylor expansion: $y(x) = y(1) + (x-1)y'(1) + (x-1)^2/2! y''(1) + \dots + (x-1)^n y^{(n)}(1) + \int_1^x g(t) dt$, where $g(t) = (x-t)^n y^{(n+1)}(t)/n! = (x-t)^n f(t)/(n! t^{n+1})$.

Problem A4

Show that for any sequence of positive reals, a_n , we have $\limsup_{n \rightarrow \infty} n((a_{n+1} + 1)/a_n - 1) \geq 1$. Show that we can find a sequence where equality holds.

Solution

Suppose $\limsup < 1$. Then we cannot find an infinite subsequence with $n((a_{n+1} + 1)/a_n - 1) \geq 1$, so we must have $n((a_{n+1} + 1)/a_n - 1) < 1$ for all sufficiently large n . Suppose it is true for all $n \geq N$. Then $(a_{N+1} + 1)/a_N < (N+1)/N$ and hence $a_N/N > (a_{N+1} + 1)/(N+1) = a_{N+1}/(N+1) + 1/(N+1)$. But similarly, $a_{N+1}/(N+1) > a_{N+2}/(N+2) + 1/(N+2)$ and so on. Hence $a_N \geq 1/(N+1) + 1/(N+2) + \dots$, which is impossible since the rhs diverges. So we have established that $\limsup \geq 1$.

We can experiment with various series. $a_n = n$ gives 2. $a_n = n^2$ also gives 2. Higher powers give higher values. So we try looking at exponents between 1 and 2. $a_n = n^{1+k}$ gives $1+k$. So by taking k arbitrarily small we can get close to 1, but we cannot reach it. It is often helpful to think of $\log n$ as n^k with k infinitesimally small. So we try $a_n = n \log n$. That gives $n((a_{n+1} + 1)/a_n - 1) = (1 + n \log(1 + 1/n) + \log(n + 1))/\log n = (1 + n \log(1 + 1/n) + (\log(1 + 1/n))/\log n + 1)$. But $1 + n \log(1 + 1/n) + \log(1 + 1/n)$ is bounded and so $n((a_{n+1} + 1)/a_n - 1)$ has limit 1 (and hence also $\limsup = 1$).

Problem A5

\mathbb{R} is the reals. $f : [0, \pi] \rightarrow \mathbb{R}$ is continuous and $\int_0^\pi f(x) \sin x \, dx = \int_0^\pi f(x) \cos x \, dx = 0$. Show that f is zero for at least two points in $(0, \pi)$. Hence or otherwise, show that the centroid of any bounded convex open region of the plane is the midpoint of at least three distinct chords of its boundary.

Solution

$\sin x > 0$ for all x in $(0, \pi)$, so if f did not change sign, then we could not have $\int_0^\pi f(x) \sin x \, dx = 0$. If there was only one sign change, at k say, then $\sin(x - k)$ would also have only one sign change and $f(x) \sin(x - k)$ would not change sign in the interval, so $\int_0^\pi f(x) \sin(x - k) \, dx = \cos k \int_0^\pi f(x) \sin x \, dx - \sin k \int_0^\pi f(x) \cos x \, dx$ would be non-zero. Contradiction. So f must have at least two sign changes and hence at least two zeros in the interval.

Take polar coordinates with the centroid as origin. Let the boundary be $f(\theta)$ for $0 \leq \theta \leq \pi$ and $-g(\theta)$ for $0 > -\theta > -\pi$.

Taking moments about the x-axis gives $\int_0^\pi f(\theta) \sin \theta \, d\theta = \int_0^\pi g(\theta) \sin \theta \, d\theta = \int_0^\pi h(\theta) \sin \theta \, d\theta$, where $h(\theta) = g(\pi - \theta)$. So $\int (f(\theta) - h(\theta)) \sin \theta \, d\theta = 0$.

Similarly, taking moments about the y-axis gives $\int f(\theta) \cos \theta \, d\theta + \int g(\theta) \cos \theta \, d\theta = 0$ or $\int (f(\theta) - h(\theta)) \, d\theta = 0$. So the result proved in the first part gives that $f(\theta) = g(\pi - \theta)$ for at least two values in $(0, \pi)$. In other words, the centroid is the midpoint of at least two chords. But we could take the x-axis to be one of these chords and then repeat the result to get two more, giving three in total.

Problem A6

M is the midpoint of a chord PQ of an ellipse. A, B, C, D are four points on the ellipse such that AC and BD intersect at M . The lines AB and PQ meet at R , and the lines CD and PQ meet at S . Show that M is also the midpoint of RS .

Solution

The key is to generalise and think of the ellipse and the line pair as conics. Take M as the origin and PQ as the x-axis. The ellipse has equation $ax^2 + bxy + cy^2 + dx + ey + f = 0$. If P is $(k, 0)$, then Q must be $(-k, 0)$, so $ak^2 + dk + f = ak^2 - dk + f = 0$. Hence $d = 0$. The line AC must have equation $y = gx$ for some g and the line BD must have equation $y = hx$ for some h . So the line pair AC, BD is the conic with equation $(y - gx)(y - hx) = 0$. The ellipse and the line pair are two distinct conics through A, B, C, D , so any conic through those four points has equation $r(ax^2 + bxy + cy^2 + ey + f) + s(y - gx)(y - hx)$. This conic meets PQ (which is $y = 0$) at the points given by $r(ax^2 + f) + sghx^2$. This has no x term, so the two points of intersection are equally spaced about M . In particular, this is true for the line pair AB, CD .

Problem B1

Find all integers n for which $x^2 - x + n$ divides $x^{13} + x + 90$.

Solution

Answer: $n = 2$.

If n is negative or zero, then the quadratic has two real roots. But we can easily check that the other polynomial has derivative everywhere positive and hence only one real root. So n must be positive.

If $x^2 - x + n$ divides $x^{13} + x + 90$, then $x^{13} + x + 90 = p(x)(x^2 - x + n)$, where $p(x)$ is a polynomial with integer coefficients. Putting $x = 0$, we see that n must divide 90. Putting $x = 1$, we see that it must divide 92. Hence it must divide $(92 - 90) - 2$. So the only possibilities are 1 and 2. Suppose $n = 1$. Then putting $x = 2$, we have that 3 divides $2^{13} + 92$. But 2^{odd} is congruent to 2 mod 2, so $2^{13} + 92$ is congruent to 1 mod 3. So n cannot be 1.

To see that $n = 2$ is possible, we write explicitly: $(x^2 - x + 2)(x^{11} + x^{10} - x^9 - 3x^8 - x^7 + 5x^6 + 7x^5 - 3x^4 - 17x^3 - 11x^2 + 23x + 45) = x^{13} + x + 90$.

Problem B2

Is the set $\{2^m 3^n: m, n \text{ are integers}\}$ dense in the positive reals?

Solution

Answer: yes.

It is easier to consider the set $X = \{m \log 2 + n \log 3: m, n \text{ integers}\}$. We note first that there is no pair of integers m, n (except $(0, 0)$) such that $m \log 2 + n \log 3 = 0$. For we would then have $2^m 3^n = 1$. That is clearly impossible if m and n have the same sign. If m and n have opposite signs, then it would imply that $2^{|m|} = 3^{|n|}$, but a power of 2 is not divisible by 3.

If X has a least positive member k , then $\log 2$ must be an integral multiple of k (for we can write $\log 2 = qk + r$ for some integer q and $0 \leq r < k$, but then r must be zero). Similarly $\log 3$. So if $\log 2 = nk$ and $\log 3 = mk$, then $m \log 2 - n \log 3 = 0$, contradicting what we just proved.

X contains the positive member $\log 2$. But $\log 2$ cannot be the least member, so it contains another member in $(0, \log 2)$. That cannot be the least member, so we can find another smaller positive member, and so on. So we get a countable infinity of members in $(0, \log 2)$. Hence for any $\varepsilon > 0$, there are infinitely many members in a subinterval of $(0, \log 2)$ of length ε . The difference between any two of these must also lie in X and will lie in the interval $(0, \varepsilon)$. Now the multiples of this number all lie in X and form a grid of mesh less than ε , so at least one of them is within ε of any given real.

That establishes that X is dense in the reals. But the map e^x maps the reals to the positive reals and is continuous, so the set $\{2^m 3^n: m, n \text{ are integers}\}$ is dense in the positive reals.

Problem B3

R is the reals. Find all $f: R \rightarrow R$ which are twice differentiable and satisfy: $f(x)^2 - f(y)^2 = f(x+y)f(x-y)$.

Solution

Differentiate wrt x : $2f(x)f'(x) = f'(x+y)f(x-y) + f(x+y)f'(x-y)$. Differentiate the result wrt y : $0 = f''(x+y)f(x-y) - f'(x+y)f'(x-y) + f'(x+y)f'(x-y) - f(x+y)f''(x-y)$. Put $X = x+y$, $Y = x-y$. Then we have $f''(X)f(Y) = f(X)f''(Y)$.

If $f(X) = 0$ for all X , then we have a solution. So suppose for some X_0 , $f(X_0)$ is non-zero and put $k = f''(X_0)/f(X_0)$, then $f''(Y) = k f(Y)$. Now we consider separately $k = 0$, $k \leq 0$ and $k > 0$. If $k = 0$, then integrating gives $f(Y) = AY + B$. But putting $y = 0$ in the original relation gives immediately that $f(y) = 0$, so $B = 0$ and we have the solution $f(Y) = AY$. This includes the solution $f(Y) = 0$ noticed earlier.

If $k < 0$, put $k = -a_2$. Then $f(Y) = A \sin aY + B \cos aY$. But $f(0) = 0$, so $B = 0$ and we have the solution $f(Y) = A \sin aY$. If $k > 0$, put $k = a_2$. Then $f(Y) = A \sinh aY + B \cosh aY$. Again $B = 0$ and we have the solution $f(Y) = A \sinh aY$.

It remains to check that these are solutions. It is obvious that $f(Y) = AY$ is a solution. If $f(Y) = A \sin aY$, then $f(x+y)f(x-y) = A^2(\sin ax \cos ay + \cos ax \sin ay)(\sin ax \cos ay - \cos ax \sin ay) = A^2(\sin^2 ax \cos^2 ay - \cos^2 ax \sin^2 ay) = A^2(\sin^2 ax - \sin^2 ay) = f(x)^2 - f(y)^2$, as required. Similarly for the sinh expression.

Problem B4

Γ is a closed plane curve enclosing a convex region and having a continuously turning tangent. A, B, C are points of Γ such that ABC has the maximum possible perimeter p . Show that the normals to Γ at A, B, C are the angle bisectors of ABC . If A, B, C have this property, does ABC necessarily have perimeter p ? What happens if Γ is a circle?

Solution

Let L be the external bisector of angle C (so that it is perpendicular to the internal bisector of angle ACB). Let B' be the reflection of B in the line L . If X is any point on the other side of L from ABC , then $BX > B'X$, so $AX + BX >$

$AX + B'X$. But $AX + B'X \geq AB'$ (with equality iff X lies on AB'), and $AB' = AC + CB$. So $AX + BX > AC + CB$. Thus the perimeter of $AXB > p$. Hence X cannot belong to the region. But if L does not coincide with the tangent at C then it will intersect the region in further points and so the region will contain points on both sides of L . So L must be the tangent.

No, this is not a sufficient condition. For example, take the curve Γ to be an equilateral triangle with rounded corners. Then take A, B, C to be the midpoints of the sides. In this case the perimeter is certainly not maximal.

If Γ is a circle, then they achieve the maximum by taking ABC to be equilateral.

Problem B5

The series $\sum a_n$ of non-negative terms converges and $a_i \leq 100a_n$ for $i = n, n+1, n+2, \dots, 2n$. Show that $\lim_{n \rightarrow \infty} na_n = 0$.

Solution

We need to invert the inequality given. We are given a collection of a_i which are less than a fixed a_n . We want to fix a_n and find a collection of a_j such that $a_n \leq 100 a_j$.

Evidently, $a_{2n} \leq 100 a_{2n-1}$, $a_{2n} \leq 100 a_{2n-2}$, $a_{2n} \leq 100 a_{2n-3}$, ... , $a_{2n} \leq 100 a_n$. Adding and multiplying by two, $2n a_{2n} \leq 200 (a_n + a_{n+1} + \dots + a_{2n-1})$. But $\sum a_n$ converges, so $(a_n + a_{n+1} + \dots + a_{2n-1}) < \varepsilon/200$ for all sufficiently large n , and hence $2n a_{2n} < \varepsilon$ for sufficiently large n . Similarly for $(2n+1) a_{2n+1}$. So na_n tends to zero.

Problem B6

Let $S = S_0$ be a set of points in space. Let $S_n = \{ P : P \text{ belongs to the closed segment } AB, \text{ for some } A, B \in S_{n-1} \}$. Show that $S_2 = S_3$.

Solution

We have to show that S_2 is already the convex hull of S . We can define the convex hull of points x_i $i = 1, 2, \dots, n$ as the collection of all points $\sum \lambda_i x_i$ with all λ_i non-negative and $\sum \lambda_i = 1$. Given two such points $P = \sum \lambda_i x_i$ and $Q = \sum \mu_i x_i$, any point on the closed segment PQ can be written as $\lambda P + \mu Q$ with λ, μ non-negative and $\lambda + \mu = 1$, but $\lambda P + \mu Q = \sum (\lambda \lambda_i + \mu \mu_i) x_i$ and $\sum (\lambda \lambda_i + \mu \mu_i) = 1$. So it follows immediately that all S_n are subsets of the convex hull.

Now we claim that any point in the convex hull can be written as a sum $\sum \lambda_i x_i$ with at most 4 non-zero terms. Suppose not. Then some point P requires at least $m \geq 5$ terms. wlog we may take these to be the first m terms, so that $P = \sum_{i=1}^m \lambda_i x_i$. But now we can find μ_i not all zero so that $\sum_{i=1}^m \mu_i x_i = 0$ (3 equations, one for each coordinate) and $\sum \mu_i = 0$ (1 equation). But now $P = \sum_{i=1}^m (\lambda_i + k \mu_i) x_i$ for all k and we still have $\sum (\lambda_i + k \mu_i) = 1$. Take the smallest λ_i/μ_i and set k as its negative. Then $\lambda_i + k \mu_i = 0$ but the other terms $\lambda_j + k \mu_j$ remain non-negative, so we have expressed P as a sum of less than m terms. Contradiction.

So given any P in the convex hull we may write (wlog) $P = \lambda_1 x_1 + \dots + \lambda_4 x_4$ for some non-negative λ_i with sum 1. If two or three of the λ_i are zero, then this shows that P is in S_1 . So suppose $\lambda_1, \lambda_2, \lambda_3$ are all positive. Then $P = (\lambda_1 + \lambda_2) (\lambda_1 x_1 / (\lambda_1 + \lambda_2) + \lambda_2 x_2 / (\lambda_1 + \lambda_2)) + (\lambda_3 + \lambda_4) (\lambda_3 x_3 / (\lambda_3 + \lambda_4) + \lambda_4 x_4 / (\lambda_3 + \lambda_4))$, which shows that P is in S_2 .

25th Putnam 1964

Problem A1

Let $A_1, A_2, A_3, A_4, A_5, A_6$ be distinct points in the plane. Let D be the longest distance between any pair, and d the shortest distance. Show that $D/d \geq \sqrt{3}$.

Solution

Let ABC be a triangle with sides $a = BC \geq b = CA \geq c = AB$. Suppose that angle $A \geq 2\pi/3$. It follows immediately from the cosine formula we have $a^2 = b^2 + c^2 - 2bc \cos A \geq b^2 + c^2 + bc \geq 3c^2$, and hence that $a/c \geq \sqrt{3}$. So it is sufficient to find a triangle with an angle of at least $2\pi/3$.

If the 6 points form a convex hexagon, then the six angles of the hexagon sum to 4π so at least one is at least $2\pi/3$. If not then one point is inside the convex hull of the others. Draw diagonals to triangulate the hull. Then the inside point must lie inside (or on) one of the triangles. But if P lies inside (or on) the triangle ABC , then at least one of the angles APB, BPC, CPA is at least $2\pi/3$.

Problem A2

α is a real number. Find all continuous real-valued functions $f : [0, 1] \rightarrow (0, \infty)$ such that $\int_0^1 f(x) dx = 1$, $\int_0^1 x f(x) dx = \alpha$, $\int_0^1 x^2 f(x) dx = \alpha^2$.

Solution

We have $\int_0^1 (\alpha - x)^2 f(x) dx = \alpha^2 - 2\alpha^2 + \alpha^2 = 0$. But the integrand is positive, except possibly for one point of the range, so the integral must also be positive. Contradiction. So there are no functions with this property.

Problem A3

The distinct points x_n are dense in the interval $(0, 1)$. x_1, x_2, \dots, x_{n-1} divide $(0, 1)$ into n sub-intervals, one of which must contain x_n . This part is divided by x_n into two sub-intervals, lengths a_n and b_n . Prove that $\sum a_n b_n (a_n + b_n) = 1/3$.

Solution

The trick is to notice that $3a_n b_n (a_n + b_n) = (a_n + b_n)^3 - a_n^3 - b_n^3$. The first $n-1$ points x_1, \dots, x_{n-1} divide the interval into n sub-intervals. Let c_n be the sum of the cubes of these sub-intervals. Then $3 \sum_1^{n-1} a_i b_i (a_i + b_i) = 1 - c_n$. So it is sufficient to prove that c_n tends to zero.

Take $\varepsilon > 0$ and < 1 . Since the points x_n are dense, we can take N so that for $n > N$ all the subintervals are smaller than ε . Then $c_n < \varepsilon^2$ times the sum of the sub-interval lengths $= \varepsilon^2 < \varepsilon$.

Problem A4

The sequence of integers u_n is bounded and satisfies $u_n = (u_{n-1} + u_{n-2} + u_{n-3}u_{n-4}) / (u_{n-1}u_{n-2} + u_{n-3} + u_{n-4})$. Show that it is periodic for sufficiently large n .

Solution

This is almost trivial. The u_n are bounded and integral, so there are only *finitely* many possible values. Hence there are only finitely many possible values for the 4-tuples $(u_{n-1}, u_{n-2}, u_{n-3}, u_{n-4})$. So we must eventually get a repeat. Suppose the t -tuples are the same for $n = N$ and $n = N+M$. Then the recurrence relation implies that they must also be the same for $n = N+1$ and $n = N+M+1$, and for $n = N+2$ and $N+M+2$ and so on. In other words the sequence is periodic (with period M) from this point onwards.

Problem A5

Find a constant k such that for any positive a_i , $\sum_1^\infty n / (a_1 + a_2 + \dots + a_n) \leq k \sum_1^\infty 1/a_n$.

Solution

Let us write $s_n = a_1 + a_2 + \dots + a_n$ and $b_n = n/s_n$. If $\sum 1/a_n$ diverges, then there is nothing to prove. So assume it converges. Hence a_n must diverge, so there are only finitely many values less than any given K , so we can arrange the terms in increasing order $c_1 \leq c_2 \leq c_3 \dots$. Since all the terms are positive $\sum 1/a_n$ is absolutely convergent and equal to $\sum 1/c_n$. Also we have $s_n \geq c_1 + c_2 + \dots + c_n$. The trick is to notice that half the c_i are at least $c_{n/2}$. It is convenient to consider separately $n = 2m$ and $n = 2m+1$. So $s_{2m} \geq m c_m$. Hence $b_{2m} < 2/c_m$. Similarly, $s_{2m-1} > m c_m$, so $b_{2m-1} < 2/c_m$. Hence $\sum_1^{2m} b_i \leq 4 \sum_1^m 1/c_i$, which gives the required result with $k = 4$.

Problem A6

S is a finite set of collinear points. Let k be the maximum distance between any two points of S . Given a pair of points of S a distance $d < k$ apart, we can find another pair of points of S also a distance d apart. Prove that if two pairs of points of S are distances a and b apart, then a/b is rational.

Solution

Since we are only interested in ratios, we may take $k = 1$ and the points to have coordinates $x_1 = 0, x_2, \dots, x_{n-1}, x_n = 1$. Let these points generate a vector space V of dimension m over \mathbb{Q} .

If $m = 1$, then every x_i is a rational multiple of x_n and hence rational, so the ratio of any two distances is rational.

Suppose that $m > 1$. Take a basis b_1, b_2, \dots, b_m for V as follows. Take $b_1 = 1$. Let $b_2 = x_t$, for some irrational x_t , then extend $\{b_1, b_2\}$ to a basis. Then each x_i is a unique rational linear combination of the b_j . In particular, $x_1 = 0, x_t = b_2$ and $x_n = b_1$. Now define a linear map f from V to \mathbb{Q} as follows. Let $f(b_2) = r b_2$, where r is rational and sufficiently large that $f(b_2) > 1$, and $f(b_i) = b_i/2$ for all other i .

Now suppose we take two distinct points in S , we can write them as $\sum r_i b_i$ and $\sum s_i b_i$, where all r_i and s_i are rational. Hence their images under f are $r_2 b_2 + \sum r_i b_i/2$ and $s_2 b_2 + \sum s_i b_i/2$, where the summations exclude $i = 2$. These must be unequal, otherwise we would have a linear combination of b_i with rational coefficients which was zero.

So there is a unique point x in S with $f(x)$ maximum and a unique point y in S with $f(y)$ minimum. Then $f(x - y)$ is greater than $f(d)$ where d is any other difference $x_i - x_j$. But every difference except $x_n - x_1$ and $x_1 - x_n$ occurs at least twice, so x and y must be the endpoints. Now $f(x_1) = 0$ and $f(x_n) = 1/2$, so $f(x - y) = 1/2$. But $f(x_t - x_1) = f(x_t) = f(b_2) > 1$. Contradiction.

Problem B1

a_n are positive integers such that $\sum 1/a_n$ converges. b_n is the number of a_n which are $\leq n$. Prove $\lim b_n/n = 0$.

Solution

If not, then for some $k > 0$ we can find arbitrarily large N such that $b_N > kN$. Then at least kN of the integers a_1, \dots, a_N do not exceed N . Let S be the set of indices i from $1, 2, \dots, N$ with $a_i \leq N$. Then since S has at least kN members, the sum over S of $1/a_i$ is at least k .

The idea is to take $N' > N$ with $b_{N'} > kN'$ and to ensure that half the indices i with $a_i \leq N'$ were not included in S . Take S' to be these new indices. Then since S' does not overlap S , the sum over S and S' of $1/a_i$ is at least $k + k/2$. We then repeat always choosing the new set of indices not to overlap any of the previous sets. So at each stage we will add $k/2$ to the sum of $1/a_i$, which must therefore diverge, establishing a contradiction.

We can do this by taking the successor N' to N sufficiently big. In fact, it is sufficient to take $N'/2 > N/k$ because we know that all of the previous index sets are contained in $\{1, 2, \dots, N\}$, so to avoid them we do not have to drop more than $N < kN'/2$ indices, leaving at least $kN'/2$ new indices.

Problem B2

S is a finite set. A set P of subsets of S has the property that any two members of P have at least one element in common and that P cannot be extended (whilst keeping this property). Prove that P contains just half of the subsets of S .

Solution

For any subset X of S , P can contain at most one of X and $S-X$, so P cannot contain more than half the subsets.

Suppose P contains less than half the subsets. Then for some X , neither X nor $S-X$ are in P . Hence there must be Y in P such that X and Y have no elements in common (otherwise we could extend P to include X). So Y is a subset of $S-X$. Similarly, P must contain a subset Z of X (otherwise we could extend P to include $S-X$). But that means Y and Z cannot overlap, contradicting the fact that as members of P they must have an element in common.

Problem B3

\mathbb{R} is the reals. $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous and for any $\alpha > 0$, $\lim_{n \rightarrow \infty} f(n\alpha) = 0$. Prove $\lim_{x \rightarrow \infty} f(x) = 0$.

Solution

Take $\varepsilon > 0$. Let $A_n = \{x : |f(nx)| \leq \varepsilon\}$. Let B_n be the intersection of all A_m with $m \geq n$. The function f is continuous, so each A_n is closed. Hence also each B_n . Now suppose we can show that an open interval (a, b) is contained in some B_N . That means that for any x in (a, b) and any $n > N$ we have $|f(nx)| < \varepsilon$. Hence $|f(x)| < \varepsilon$ for any x in (na, nb) . But the intervals (na, nb) expand in size as they move to the right, so for n sufficiently large they overlap and we have that their union includes all sufficiently large x . In other words, for any sufficiently large x we have $|f(x)| < \varepsilon$. But ε was arbitrary, so we have established that $\lim_{x \rightarrow \infty} f(x) = 0$.

It remains to show that we can find such an open interval (a, b) . Now we are given that for any given x , $\lim_{n \rightarrow \infty} f(nx) = 0$ (as n tends to infinity), so x belongs to some B_n for n sufficiently large. In other words, the union of the B_n is the entire line.

We now need the Baire category theorem which states that if a union of closed sets covers the line, then one of the sets contains an open interval. This is bookwork. [But straightforward. Assume none of the B_n contain an open interval. Take a point not in B_1 . Since B_1 is closed we may take a closed interval C_1 about the point, not meeting B_1 . Having chosen C_n , there must be a point in it not in B_{n+1} (or B_{n+1} would contain an open interval - the interior of C_n). Hence we may take C_{n+1} , a closed subinterval of C_n which does not meet B_{n+1} . Now C_n is a nested sequence of closed intervals, so there must be a point X in all the C_n . But B_n cover the line, so X must be in some B_n . Contradiction.]

Problem B4

n great circles on the sphere are in general position (in other words at most two circles pass through any two points on the sphere). How many regions do they divide the sphere into?

Solution

Answer: $n^2 - n + 2$.

We use the well-known formula $E + 2 = V + F$, where E is the number of edges, V the number of vertices and F the number of faces. It is true for a sphere provided $V > 1$, so certainly for $n > 1$.

Each circle intersects every other circle in 2 vertices, so $V = n(n - 1)$. Each vertex has degree 4, so $E = 2V$. Hence $F = V + 2 = n^2 - n + 2$. It is easy to check that the formula also holds for $n = 1$.

Problem B5

Let a_n be a strictly monotonic increasing sequence of positive integers. Let b_n be the least common multiple of a_1, a_2, \dots, a_n . Prove that $\sum 1/b_n$ converges.

Solution

We need a crude upper bound on the number of divisors for N . N is too crude. But we can do better by noticing that N has at most \sqrt{N} divisors $\leq \sqrt{N}$ and hence at most $2\sqrt{N}$ divisors in total (every divisor $d > \sqrt{N}$ has a matching divisor $N/d < \sqrt{N}$). Now we know that b_n has at least n distinct divisors (namely a_1, \dots, a_n). Hence $b_n \geq n^2/4$. But we know that $\sum 1/n^2$ converges.

Problem B6

D is a disk. Show that we cannot find congruent sets A, B with $A \cap B = \emptyset$, $A \cup B = D$.

Solution

It is easy to find two congruent sets which overlap only at the centre O of the disk, so this suggests that the centre is the key.

Wlog we may assume O is in A and that the radius is 1. Let O' be the corresponding point in B . Let PQ be the diameter perpendicular to OO' . Then PO' and QO' are both greater than 1. But $OX \leq 1$ for any point X in A , so $O'Y \leq 1$ for any point Y in B . So P and Q must both be in A . Let P' and Q' be the corresponding points in B . Then $P'Q' = PQ = 2$, so $P'Q'$ must be a diameter. But $O'P' = OP = 1$ and $O'Q' = OQ = 1$, so O' must be the midpoint of $P'Q'$ and hence the centre of the disk. So $O' = O$. Contradiction.

26th Putnam 1965

Problem A1

The triangle ABC has an obtuse angle at B, and angle A is less than angle C. The external angle bisector at A meets the line BC at D, and the external angle bisector at B meets the line AC at E. Also, $BA = AD = BE$. Find angle A.

Solution

Answer: 12 degrees.

Let angle $BAC = k$. Then since $BA = BE$, angle $BEA = k$. Take B' on BA the opposite side of B to A. Then angle $B'BE = 2k$. Angle $B'BC$ is bisected by BE , so angle $CBE = 2k$. Hence angle $ACB = 3k$. So angle $DBA = 4k$. But $AD = BA$, so angle $BDA = 4k$. But AD is the exterior bisector, so angle $BAD = 90 - k/2$. The angles in BAD must sum to 180 deg, so $k = 12$ deg.

Problem A2

Let $k = [(n - 1)/2]$. Prove that $\sum_0^k \binom{n-2r}{n}^2 \binom{n}{2r}^2 = 1/n \binom{2n-2}{n-1} C(n-1)$ (where nCr is the binomial coefficient).

Problem A3

$\{a_r\}$ is an infinite sequence of real numbers. Let $b_n = 1/n \sum_1^n \exp(i a_r)$. Prove that $b_1, b_2, b_3, b_4, \dots$ converges to k iff $b_1, b_4, b_9, b_{16}, \dots$ converges to k .

Solution

If a sequence converges to k , then any subsequence also converges to k .

So suppose that the square terms converge to k . Let $c_r = \exp(i a_r)$. Take two consecutive squares $N = n^2$ and $N' = n^2 + 2n + 1$ and m between them. Then $|b_N - b_m| \leq |b_N - 1/N \sum c_r| + |(1/N - 1/m) \sum c_r|$, where the sums are over m terms. But $|b_N - 1/N \sum c_r| \leq 1/N \sum_{N+1}^m |c_r| = (m - N)/N \leq 2n/n^2 = 2/n$. Also $|(1/N - 1/m) \sum c_r| \leq (1/N - 1/m) \sum |c_r| < (1/n^2 - 1/(n+1)^2) (n+1)^2 = (2n+1)/n^2 < 3/n$. So the difference $|b_m - b_N|$ is arbitrarily small for n sufficiently large. Thus if we take m sufficiently large, then $|b_m - b_N| < \epsilon$ and $|b_N - k| < \epsilon$. So $|b_m - k| < 2\epsilon$ and the sequence converges.

Problem A4

S and T finite sets. U is a collection of ordered pairs (s, t) with $s \in S$ and $t \in T$. There is no element $s \in S$ such that all possible pairs $(s, t) \in U$. Every element $t \in T$ appears in at least one element of U . Prove that we can find distinct $s_1, s_2 \in S$ and distinct $t_1, t_2 \in T$ such that $(s_1, t_1), (s_2, t_2) \in U$, but $(s_1, t_2), (s_2, t_1) \notin U$.

Solution

Suppose that we cannot find such s_i, t_i . We will establish a contradiction.

Take t in T . Suppose that there are n distinct s in S such that (s, t) is in U . Suppose $n > 0$. Then take a specific s' such that (s', t) is in U . There must be some t' such that (s', t') is not in U . Now consider whether we have (s, t') in U . If (s, t) is not in U , then (s, t') cannot be in U (or we would have found s_i, t_i). But there are at most $n-1$ distinct s such that (s, t') is in U (the only candidates are the cases for which (s, t) is in U , and one of those, namely s' , does not work).

Iterating, we must eventually get some x in T for which there is no s in S with (s, x) in U . Contradiction.

Problem A5

How many possible bijections f on $\{1, 2, \dots, n\}$ are there such that for each $i = 2, 3, \dots, n$ we can find $j < n$ with $f(i) - f(j) = \pm 1$?

Solution

Answer: 2^{n-1} .

Consider the last element $f(n)$. Suppose it is m , not 1 or n . Then the earlier elements fall into two non-empty sets $A = \{1, 2, \dots, m-1\}$ and $B = \{m+1, m+2, \dots, n\}$. But the difference between an element of A and an element of B is at least 2. So if $f(1)$ is in A , then the first time we get an element of B it has only elements of A preceding it. Contradiction. Similarly, if $f(1)$ is in B .

So we conclude that the last element is always 1 or n. We can now prove the result by induction. Clearly given an arrangement for n we can derive one for n+1 by adding n+1 at the end. We can also derive one for n+1 by increasing each element by 1 and adding 1 at the end. Equally it is clear that all these are distinct and that there are no other arrangements for n+1 that end in 1 or n+1. So there are twice as many arrangements for n+1 as for n.

Problem A6

α and β are positive real numbers such that $1/\alpha + 1/\beta = 1$. Prove that the line $mx + ny = 1$ with m, n positive reals is tangent to the curve $x^\alpha + y^\alpha = 1$ in the first quadrant ($x, y \geq 0$) iff $m^\beta + n^\beta = 1$.

Solution

Suppose $mx + ny = 1$ is tangent to the curve. Suppose it touches at (a, b) . Differentiating, we see that the tangent at (a, b) is $a^{\alpha-1}x + b^{\beta-1}y = 1$, so $m = a^{\alpha-1}$, $n = b^{\beta-1}$. Hence, using $\alpha\beta - \beta = \alpha$, we have that $m^\beta + n^\beta = a^\alpha + b^\alpha = 1$.

Conversely, suppose that $m^\beta + n^\beta = 1$. Take $a = m^{\beta/\alpha}$, $b = n^{\beta/\alpha}$. Then $a^\alpha + b^\alpha = 1$, so (a, b) lies on the curve in the first quadrant. Its tangent is $Mx + Ny = 1$, where $M = a^{\alpha-1}$, $N = b^{\beta-1}$. But $a = m^{\beta/\alpha}$ and $\beta/\alpha(\alpha - 1) = 1$, so $M = m$. Similarly, $N = n$. Thus we have established that $mx + ny = 1$ is tangent to the curve in the first quadrant as required.

Problem B1

X is the unit n -cube, $[0, 1]^n$. Let $k_n = \int_X \cos^2(\pi(x_1 + x_2 + \dots + x_n)/(2n)) dx_1 \dots dx_n$. What is $\lim_{n \rightarrow \infty} k_n$?

Solution

Let $y_i = 1 - x_i$ and change variables from x_i to y_i . The sum $(x_1 + x_2 + \dots + x_n)/(2n)$ becomes $1/2 - (y_1 + y_2 + \dots + y_n)/(2n)$, so the integrand becomes $\sin^2(\pi(y_1 + y_2 + \dots + y_n)/(2n))$. So the integral is identical to the original except that \cos has been changed to \sin . Thus we can add it to the original to get $2k_n = \int_X dx_1 dx_2 \dots dx_n = 1$. So $k_n = 1/2$ for all n .

Problem B2

Every two players play each other once. The outcome of each game is a win for one of the players. Player n wins a_n games and loses b_n games. Prove that $\sum a_n^2 = \sum b_n^2$.

Solution

Suppose there are N players in total. Each player plays $N-1$ games, so $b_n = N - 1 - a_n$. Hence $\sum b_n^2 = \sum (N - 1)^2 - 2(N - 1) \sum a_n + \sum a_n^2 = N(N - 1)^2 - 2(N - 1) \sum a_n + \sum a_n^2$.

Each game is won by just one player, so $\sum a_n = \text{no. of games} = N(N - 1)/2$. Hence $\sum b_n^2 = N(N - 1)^2 - 2(N - 1)N(N - 1)/2 + \sum a_n^2 = \sum a_n^2$.

Problem B3

Show that there are just three right angled triangles with integral side lengths $a < b < c$ such that $ab = 4(a + b + c)$.

Solution

Answer: 12, 16, 20; 10, 24, 26; 9, 40, 41.

We need the result that for some integral d, m, n we have $c = d(m^2 + n^2)$ and $b = 2dmn$, $a = d(m^2 - n^2)$ or $b = d(m^2 - n^2)$, $a = 2dmn$ (*).

It follows that $4(a + b + c) = 4d(2m^2 + 2mn)$, $ab = 2d^2mn(m^2 - n^2)$. Hence, $4 = dn(m - n)$. So we must have $n = 1, 2$ or 4 and $(d, m, n) = (1, 5, 1), (2, 3, 1), (4, 2, 1), (1, 4, 2), (2, 3, 2)$ or $(1, 5, 4)$, giving the three answers above.

It remains to prove (*). Let d be the gcd of a and b . It follows that d also divides c . Put $a = dA$, $b = dB$, $c = dC$, so that A, B, C are relatively prime in pairs. C cannot be even, for then A and B would both be odd and hence $A^2 + B^2$ would be congruent to $2 \pmod{4}$, which is impossible, since C is a square. So C must be odd and just one of A, B must be even. Assume A is odd. Then $A^2 = (C - B)(C + B)$. If an odd prime p divides A , then it must divide $C - B$ or $C + B$. It cannot divide both, for then it would also divide B and C . So $C - B$ and $C + B$ must both be odd squares. Say $C + B = h^2$, $C - B = k^2$. Then $A = hk$, $B = (h^2 - k^2)/2$, $C = (h^2 + k^2)/2$, with h and k odd. Put $m = (h + k)/2$, $n = (h - k)/2$, then $h = m + n$, $k = m - n$ and $A = m^2 - n^2$, $B = 2mn$, $C = m^2 + n^2$, so we have put a, b, c , in the form (*). On the other hand, it is obvious that if a, b, c have this form then they satisfy $a^2 + b^2 = c^2$.

Problem B4

Define $f_n : \mathbb{R} \rightarrow \mathbb{R}$ by $f_n(x) = (nC_0 + nC_2 x + nC_4 x^2 + \dots) / (nC_1 + nC_3 x + nC_5 x^2 + \dots)$, where nC_m is the binomial coefficient. Find a formula for $f_{n+1}(x)$ in terms of $f_n(x)$ and x , and determine $\lim_{n \rightarrow \infty} f_n(x)$ for all real x .

Solution

It is almost obvious that $f_{n+1} = (f_n + x)/(f_n + 1)$ (*). So if $f_n(x)$ tends to a limit $k(x)$, then $k(x) = (k(x) + x)/(k(x) + 1)$, and hence $k(x) = \sqrt{x}$.

Obviously, $f_n(0) = 1/n$, so $k(0) = 0$. We notice also that $f_n(x) = (\sqrt{x}) N/D$, where $N = (1 + \sqrt{x})^n + (1 - \sqrt{x})^n$ and $D = (1 + \sqrt{x})^n - (1 - \sqrt{x})^n$.

Suppose $0 < x \leq 1$. Then put $y = (1 - \sqrt{x})/(1 + \sqrt{x})$. We have $0 \leq y < 1$ and $N/D = (1 + y^n)/(1 - y^n)$ which tends to 1, so in this case also $k(x) = \sqrt{x}$.

Similarly if $x > 1$, put $y = (\sqrt{x} - 1)/(1 + \sqrt{x})$ and we again get $k(x) = \sqrt{x}$.

It is clear from (*), that for $x < 0$, $f_n(x)$ does not tend to a limit.

Problem B5

Let S be a set with $n > 3$ elements. Prove that we can find a collection of $\lfloor n^2/4 \rfloor$ 2-subsets of S such that for any three distinct elements A, B, C of the collection, $A \cup B \cup C$ has at least 4 elements.

Solution

This is just another form of finding a graph without triangles. If $n = 2m$, then take S to be the disjoint union of U and V , where U and V each have m elements. Now take K to be the collection all sets $\{u, v\}$ with u in U and v in V . Evidently K has $m^2 = \lfloor n^2/4 \rfloor$ members. Suppose we have three distinct elements A, B, C of K . $A \cup B \cup C$ cannot have less than three elements, for then two of A, B, C would be identical. So if it has less than 4 elements, then it has just three elements. But that means that for some a, b, c , we have $A = \{b, c\}$, $B = \{a, c\}$, $C = \{a, b\}$. Suppose a is in U . Then, considering B and C , c and b must be in V , but then A has two elements in V . Contradiction. Similarly if a is in V . So K has the required property.

Similarly, if $n = 2m+1$, take S to be the disjoint union of U and V , where U has $m+1$ elements and V has m elements. Then K , the set of all $\{u, v\}$ with u in U and v in V has $m(m+1) = \lfloor n^2/4 \rfloor$ elements and has the required property by the same argument.

Problem B6

Four distinct points A_1, A_2, B_1, B_2 have the property that any circle through A_1 and A_2 has at least one point in common with any circle through B_1 and B_2 . Show that the four points are collinear or lie on a circle.

Solution

The trick is to take *concentric* circles. If A_1A_2 is not parallel to B_1B_2 then their perpendicular bisectors must intersect at some point O . Take circles centre O through A_1, A_2 and through B_1, B_2 . These must either coincide, in which case the 4 points lie on a circle, or have no points in common.

In the case where A_1A_2 is parallel to B_1B_2 . Assume they do not coincide (otherwise the 4 points would be collinear). Then take a point O on the perpendicular bisector of A_1A_2 on the opposite side of A_1A_2 to B_1B_2 and sufficiently distant from A_1A_2 that the circle through A_1, A_2 centre O only extends less than halfway towards the line B_1B_2 . Similarly, take a circle through B_1, B_2 which extends less than halfway towards A_1A_2 . Then these circles will not meet.